THE MORPHOLOGY AND SYSTEMATICS OF SOME CRETACEOUS CRIBRIMORPH POLYZOA (PELMATOPORINAE)

BY

GILBERT POWELL LARWOOD

BRITISH MUSEUM
12 JUN 62
NATURAL HISTORY

Pp. 1–285 ; *Plates* 1–23 ; 132 *Text-figures*

BULLETIN OF
THE BRITISH MUSEUM (NATURAL HISTORY)
GEOLOGY Vol. 6 No. 1
LONDON : 1962

THE BULLETIN OF THE BRITISH MUSEUM (NATURAL HISTORY), *instituted in* 1949, *is issued in five series corresponding to the Departments of the Museum, and an Historical series.*

Parts will appear at irregular intervals as they become ready. Volumes will contain about three or four hundred pages, and will not necessarily be completed within one calendar year.

This paper is Vol. 6, No. 1 *of the Geological (Palaeontological) series.*

© Trustees of the British Museum, 1962

PRINTED BY ORDER OF THE TRUSTEES OF
THE BRITISH MUSEUM

Issued June, 1962 *Price Six Pounds*

THE MORPHOLOGY AND SYSTEMATICS OF SOME CRETACEOUS CRIBRIMORPH POLYZOA (PELMATOPORINAE)

By G. P. LARWOOD

CONTENTS

Page

I. INTRODUCTION AND ACKNOWLEDGMENTS 5

II. GENERAL STRATIGRAPHY 7

  Subdivisions of the English Chalk 8

  Foreign Upper Cretaceous Horizons 14

III. NOTES ON THE COLLECTIONS STUDIED 19

IV. PRESERVATION OF THE MATERIAL 20

V. TECHNIQUES 20

VI. MORPHOLOGY OF CRETACEOUS CRIBRIMORPH POLYZOA . . . 21

VII. SYSTEMATIC DESCRIPTIONS 51

  Genus *Aeolopora* Lang 54

    Aeolopora distincta Lang 55

    ,, *nebulosa* Lang 62

  Genus *Lagynopora* Lang 65

    Lagynopora pustulosa sp. nov. 67

    ,, *horsleyensis* Lang 68

    ,, *praecursor* (Lang) nov. comb. 71

    ,, *saltdeanensis* Lang 73

    ,, *birlingensis* Lang 75

    ,, *amphora* Lang 77

    ,, *ampulla* Lang 79

    ,, *furcifera* (Brydone) 81

    ,, *ollula* Lang 83

    ,, *lagena* Lang 86

    ,, *urceolus* Lang 90

    ,, *vasculum* Lang 93

  Species of uncertain systematic position previously assigned to *Lagynopora* 95

  Genus *Hexacanthopora* Lang 97

    Hexacanthopora sexspinosa Lang 98

    ,, *viginticostata* Voigt 102

  Genus *Leptocheilopora* Lang 102

    Leptocheilopora magna Lang 103

    ,, *tenuilabrosa* Lang 107

    ,, *filliozati* (Brydone) 112

    ,, *vulnerata* (Brydone) 116

  Species of uncertain systematic position previously assigned to *Leptocheilopora* 118

Page
Genus *Pelmatopora* Lang 119
Pelmatopora calceata Lang 123
,, *crepidaria* Lang 125
,, *solearis* Lang 127
,, *gasteri* Lang 130
,, *chrysalis* (d'Orbigny) 132
,, *larva* Lang 134
,, *striata* (d'Orbigny) 136
,, *fragilis* (d'Orbigny) 138
,, *d'orbignyi* Lang 140
,, *quadrata* Lang 143
,, *fecampensis* Lang 145
,, *pero* Lang 148
,, *suffulta* (Brydone) 151
,, *simplex* Lang 157
,, *plantaris* Lang 159
,, *quadrivolucris* Lang 161
,, *brydonei* Lang 164
,, *coryli* (Lang) 168
,, *marsupitorum* Lang 171
,, *gregoryi* (Brydone) 175
,, *somptingensis* Lang 188
,, *palmata* Lang 196
,, sp. 199
Species of uncertain systematic position previously assigned to *Pelmatopora* 200
Genus *Castanopora* Lang 203
Castanopora armata sp. nov. 206
,, *labiata* (Levinsen) 208
,, *retrorsa* Lang 209
,, *castanea* Lang 211
,, *dibleyi* (Brydone) 215
,, *guascoi* (Ubaghs) 219
,, *spooneri* (Butler & Cheetham) 220
,, *multicostata* Voigt 221
,, *jurassica* (Gregory) 223
,, *glandulosa* Lang 225
,, *aviculosa* Lang 226
,, *magnifica* (d'Orbigny) 228
,, *faujasi* (von Hagenow) 236
,, *voigti* sp. nov. 236
Species of uncertain systematic position previously assigned to *Rhiniopora* and *Castanopora* 237
Genus *Ubaghsia* Jullien 238
Ubaghsia ranunculus (Lang) 242
,, *aurita* (Lang) 244
,, *gasteri* sp. nov. 245
,, *langi* sp. nov. 247
,, *crassa* (Lang) 250
,, *coaxans* (Lang) 252
,, *perforata* (Marsson) 254
,, *ocellata* (Jullien) 255
,, *meudonensis* (Jullien) 256

Page
Ubaghsia demorgani (Jullien) 256
,, *reticulata* (Ubaghs) 257
,, *ornata* (Goldfuss) 260
Species of uncertain systematic position previously assigned to *Batrachopora* 264
Genus *Andriopora* Lang 266
Andriopora minor sp. nov. 266
,, *major* sp. nov. 268
VIII. SYSTEMATIC BASIS, STRATIGRAPHY AND EVOLUTION OF THE GENERA AND SPECIES REVISED 269
IX. REFERENCES 275

SYNOPSIS

Ten genera of Cretaceous cribrimorph Polyzoa, previously described by Lang, are considered to belong to the Pelmatoporinae. Some of these genera, and numerous species, are regarded as synonymous and all are diagnosed and revised. Seven new species are diagnosed and described.

The morphology of Cretaceous cribrimorph Polyzoa is discussed and comparison made with Recent genera. Stratigraphical subdivisions of the English Chalk are reviewed and the distribution of the specimens from foreign Upper Cretaceous horizons is given. The stratigraphical range of the revised genera and species is described and their probable evolution reviewed.

I. INTRODUCTION AND ACKNOWLEDGMENTS

SINCE the two catalogues on the Cretaceous cribrimorph Polyzoa by Dr. W. D. Lang were published in 1921 and 1922, the British Museum (Natural History) has received two very large collections of Chalk Polyzoa from Mr. C. T. A. Gaster. This has given the opportunity to examine and revise some of those results in the light of this added material, to re-assess the value of the cribrimorph Polyzoa as stratigraphical indices, and to reconsider the evidence for their evolution. It was found that, as in Recent cheilostomatous Polyzoa, there was much morphological variation in single species of the Cretaceous cribrimorphs, and far more in fact than Lang allowed for. As a result, many of his systematic subdivisions were found to be excessive.

The study of Cretaceous cribrimorph Polyzoa is made difficult by the complexity of their structure. Confusion has also arisen from the unique terminology introduced by Lang, and from his attempts to assign systematic and stratigraphical positions to species known only from earlier literature, where they were often poorly described and inadequately figured. Consequently I have not used Lang's terminology, but have used instead the more widely accepted terms which are generally applied to Recent and fossil Polyzoa. I have also rejected, as of uncertain systematic position, the ill-defined species just mentioned. Further, I have also rejected some of the hypothetical origins to which Lang attributed the development of certain structures, to which he gave evolutional significance, but to which the evidence appears opposed.

All this has led to a general reduction in the number of genera and species recognized, to a revision of their stratigraphical ranges, and to a great modification of the evolutionary sequences enunciated by Lang.



FIG. 1. Key to the abbreviations :

Ap = zone of *Actinocamax plenus*.
Apl = subzone of *A. plenus*.
A.p.m. = *Actinocamax plenus* Marls.
Aq = zone of *Actinocamax quadratus*.
Aqu = subzone of *A. quadratus*.
G[A]pM = *Gonioteuthis* [*Actinocamax*] *plenus* Marls.
G[A]q = zone of *Gonioteuthis* [*A.*] *quadrata*.
G[A]qu = subzone of *G.* [*A.*] *quadrata*.
Bl(Ol) = zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk).

Bm = zone of *Belemnitella mucronata*.
Ec = subzone of *Echinocorys scutata* var. *cincta*.
Ed = subzone of *Echinocorys scutata* var. *depressula*.
Hp = zone of *Holaster planus*.
Hsg = zone of *Holaster subglobosus*.
Il = zone of *Inoceramus labiatus*.
M = zone of *Marsupites testudinarius*.
Ma = subzone of *M. testudinarius*.
MAS = Maastrichtian Stage.

By using modern binocular microscopes I have been able to study structures in the Cretaceous cribrimorph Polyzoa which must have been invisible to Lang or less well defined with the instruments available to him. I wish to record my high appreciation of the great value of the work done by Lang in his extensive studies of this group of Cretaceous Polyzoa.

The present work[1] has been almost wholly carried out in the British Museum (Natural History) and my thanks are due to the Keepers of Palaeontology and Zoology for the facilities provided for my researches.

I wish to record particularly my warm appreciation of the considerable assistance given to me by Dr. Anna B. Hastings and Dr. H. Dighton Thomas to whom I am indebted for their invaluable advice, interest, suggestions and criticisms. I am also much indebted to Dr. H. Dighton Thomas for reading a large part of my typescript. I also wish to acknowledge the help of the library staffs at the British Museum (Natural History) and to thank Miss P. L. Cook and Mr. R. F. Wise for their general assistance, Mr. M. G. Sawyers for the photography and Miss C. M. P. von Hayeck and Mr. A. C. Townsend for preparing translations.

I am most grateful to Dr. J. F. Kirkaldy, Queen Mary College, University of London, for his supervision of my studies and for the criticisms he has given during the course of my work.

My gratitude is also due to Mr. A. G. Brighton, Curator of the Sedgwick Museum, Cambridge, for the loan of material and for information on the R. M. Brydone Collection ; to Mr. R. R. Clarke, Curator of Norwich Castle Museum, for the loan of material ; to Mons. E. Buge, of the Laboratoire de Paléontologie, Muséum National d'Histoire Naturelle, Paris, for photographs of, and information on, specimens from the d'Orbigny Collection and to the Editor of the *Bulletin of the Zoological Society of France* for permission to reproduce Plate 22, figs. 1–8.

Finally, I record my gratitude to my wife for help and encouragement with many problems associated with this work.

II. GENERAL STRATIGRAPHY

Subdivisions of the English Chalk which have been established or adopted by previous authors are discussed below. The subdivisions followed here (column 7, Text-fig. 1, p. 6) are based on a consideration of the extensive work of the authors mentioned.

---

[1] A major part of the subject of a thesis for the degree of Doctor of Philosophy in the University of London.

Mca = zone of *Micraster cor-anguinum.*
Mct = zone of *Micraster cortestudinarium.*
Olu = subzone of *Ostrea lunata.*
Op = zone of *Offaster pilula.*
Opi = subzone of *O. pilula.*
Pl = zone of *Plocoscyphia labrosa.*
Rc = zone of *Rhynchonella cuvieri.*
Rm = zone of *Rhynchonella martini.*
Sc = zone of *Stauronema carteri.*
Sca = subzone of *S. carteri.*
Scr = subzone of *Saccacoma cretacea.*
Sv = zone of *Schloenbachia varians.*
T.B. = *Trochiliopora Bed.*
Tg = zone of *Terebratulina gracilis.*
Tl = zone of *Terebratulina lata.*
U = subzone of *Uintacrinus westfalicus.*
1 = Lower belt of zone of *Offaster pilula.*
2 = Belt of *Echinocorys scutata* var. *cincta.*
3 = Upper belt of zone of *Offaster pilula.*
I = Horizon of *Hagenowia rostrata.*
II = Horizon of abundant *Offaster pilula.*

The *Ostrea lunata* Chalk is contained well within the zone of *Belemnella lanceolata* and is Maastrichtian in age. Gaster's (1924–51) main subdivisions of the zones of *Gonioteuthis* [*Actinocamax*] *quadrata* and *Offaster pilula* are followed, but Brydone's (1912) upward extension of the zone of *Micraster cor-anguinum* to include the *Uintacrinus* Chalk (usually placed in the zone of *Marsupites testudinarius*) has not been generally accepted and is not followed here.

Subdivisions of the English Chalk

Cenomanian

Zone of *Schloenbachia varians*

The glauconitic, sandy marl, usually about one foot thick, which occurs at the base of this zone over much of southern England is here termed the subzone of *Stauronema carteri*; this agrees with Brydone's and Gaster's general use of the term. Griffith & Brydone (1911) and Withers (1935) gave the unit zonal status; it is not, however, sufficiently widespread or thick enough to be regarded as a true zone. Griffith & Brydone (1911) also suggested *Plocoscyphia labrosa* as an alternative name form, but apparently this sponge is only locally significant.

The Chalk Marl, with the Totternhoe Stone at the top, succeeds the *Stauronema carteri* subzone giving a general thickness of about 160 ft. for the zone of *Schloenbachia varians*. The thickness may be much less; at Hunstanton, Norfolk, it is only 18–20 ft. The zone of *Schloenbachia varians*, as determined for southern England, is not recognized in the Chalk of Lincolnshire and Yorkshire. Bower & Farmery's (1910) division of the Lincolnshire Cenomanian was adopted for that of Yorkshire by C. W. & E. V. Wright (1942). They divided the stage into a lower zone of *Holaster subglobosus* and an upper zone of *H. trecensis*, and estimated an inland thickness of 70 ft. for the whole stage.

Zone of *Holaster subglobosus*

In southern England this zone has a general thickness of about 100 ft. Griffith & Brydone (1911) and Withers (1935) excluded from it the *Actinocamax plenus* Marls, which they regarded as a separate zone. Subsequently both Brydone and Gaster have described these Marls as a subzone of the *Holaster subglobosus* Zone. C. W. & E. V. Wright (1951 : 3, 4) also regarded the *Actinocamax plenus* Marls as a distinct subzone, but stated " Their cephalopod fauna is of Turonian affinities, and it therefore seems best to treat them as a subzone of the *Inoceramus labiatus* zone ". The Marls vary in thickness from a few to about 12 ft. and are here regarded only as a lithological marker of the base of the Turonian in some areas.

Turonian

Zone of *Inoceramus labiatus* and " *Rhynchonella* " *cuvieri*[1]

Both name forms occur commonly in the zone which is about 70–90 ft. thick in southern England. In Yorkshire the zone is very thin—only 11 ft. of gritty Chalk with broken *Inoceramus labiatus* shells is present. As mentioned above, the *Actinocamax plenus* Marls are a lithological marker of the base of the zone.

---

[1] Pettit (1954 : 29) placed the species in *Orbirhynchia*.

Zone of *Terebratulina lata*

Withers's (1935 : 54) estimate of the general thickness of this zone in southern England as about 160 ft. is probably a maximum. Dines and others (1954) recorded up to 140 ft. of *T. lata* Chalk in the Chatham district, Kent, and Wright (1947) estimated only 58 ft. for the zone in Dorset. In Yorkshire it is very much thicker (210 ft.).

Zone of *Holaster planus*

Although Brydone and Gaster excluded this zone from the Turonian (see below), it is generally regarded as the topmost zone of that stage.

The base of the zone of *H. planus* is often well marked by the Chalk Rock. This hard, somewhat nodular, glauconitic, yellowish limestone, which is about 6 ft. thick, contains a shallow water fauna with *Hyphantoceras reussianum.* In the localities where the actual rock band is not developed, the fauna, although somewhat modified, is often present. This basal bed of the *Holaster planus* Zone is distinguished as the *Hyphantoceras reussianum* Subzone (see Withers, 1935 : 55).

The zone is generally thin ; the maximum thickness given by Withers is about 60 ft., the minimum 14 ft. However, the zone thickens northwards, Dines and others (1954) recorded up to 70 ft. of *Holaster planus* Chalk in the Chatham area of Kent, Peake & Hancock (1961) estimate 75 ft. for the zone in north Norfolk and Rowe (1904 : 193) estimated 125 ft. as the thickness of the zone in Yorkshire.

Wright (1947 : 202) drew attention to " a 2-foot band near the base of the zone [in Dorset] crowded with the polyzoan *Bicavea rotaformis* Gregory ". This very restricted occurrence of *Bicavea* contrasts with that of *B. radiata* Gaster and *B. striata* Gaster which are distributed throughout the zone of *Gonioteuthis* [*Actinocamax*] *quadrata* in the Senonian.

SENONIAN

Zone of *Micraster cortestudinarium*

As mentioned above, Griffith & Brydone (1911), Brydone (1912) and Gaster (1924–51) placed the base of the Senonian at the bottom of the *Holaster planus* Zone. The first two authors also regarded the lowest third of the generally accepted zone of *Micraster cor-anguinum* as the uppermost unit of the *Micraster cortestudinarium* Zone. Withers (1935 : 57) estimated a general thickness of about 60 ft. for the *M. cortestudinarium* Chalk.

Zone of *Micraster cor-anguinum*

Griffith & Brydone (1911) and Brydone (1912) placed the lower boundary of this zone about 85 ft. above the generally accepted level (e.g., Rowe, 1900), and raised the upper boundary by about 60 ft. to include the *Uintacrinus* Chalk ; this arrangement has not been adopted by later authors.

Peake & Hancock (1961) record 170 ft. of *M. cor-anguinum* chalk in Norfolk.

The upper boundary of the zone, as commonly defined at present, is the base of the *Uintacrinus* Chalk, its lower limit being placed about 260 ft. below that level.

Gaster (1920 : 526) first drew attention to the *Trochiliopora* Bed near the base of the zone of *Micraster cor-anguinum* in Sussex. He stated " the exact position in which it occurs . . . is as follows :

" Lower fourth of zone of *Micraster coranguinum*	Strong *M. coranguinum* tabular flint band.
	Chalk, about 35 feet.
	Chalk with *Trochiliopora* sp., 10 feet.
	Chalk, about 17 feet.

" Chalk of zone of *Micraster cortestudinarium* ".

This distinctive horizon in Sussex was not mentioned by Wright (1947) as recognizable in the Chalk of the *M. cor-anguinum* Zone of Dorset.

The " strong *Micraster cor-anguinum* tabular [flint band] " of Rowe (1900 : 301) was mentioned previously by Whitaker (1872 : 50). In his work on the Sussex Chalk Gaster (e.g., 1937 : 361) also recognized this marker flint band near the base of the zone. C. W. & E. V. Wright (1942) adopted, for Yorkshire, Rowe's (1904) suggestion of a zone of *Hagenowia rostrata* which is abundant throughout Chalk equivalent to the *Micraster cor-anguinum* Zone in southern England.

Zone of *Marsupites testudinarius*

This zone, about 125 ft. thick, has been recognized generally, e.g., Rowe (1900), Gaster (1924–51), Withers (1935) as comprising two " bands " or subzones—that of *Marsupites testudinarius* above (65 ft. thick) and that of *Uintacrinus westphalicus* below (60 ft. thick). As mentioned, Griffith & Brydone (1911) and Brydone (1912) did not include the *Uintacrinus* Chalk in their zone of *Marsupites testudinarius*. C. W. & E. V. Wright (1942 : 118) pointed out that Brydone's scheme avoids the anomaly of a " *Marsupites* Zone in one subzone of which *Marsupites* does not occur ". In the case of the Yorkshire Chalk they recognized a zone of *Uintacrinus* as have Peake and Hancock (1961) in Norfolk.

Brydone (1915 : 13–15) added a few feet of Chalk, containing unusual crinoid plates, to the top of the *Marsupites* Zone ; this change was also adopted by Gaster (1937 : 362).

Withers (1935 : 58) stated that " The *Marsupites* zone probably represents the maximum depth attained during the whole of the English Cretaceous period ".

Zone of *Offaster pilula*[1]

This zone, proposed by Brydone (1912 : 14), was established originally in the Chalk of Hampshire and later traced by him in Norfolk, Suffolk, Sussex, Dorset and the Isle of Wight (see Brydone, 1914*a*, 1932*a*, *b*). Blackmore was able to trace the zone in Wiltshire, and other workers have established its presence throughout the rest of the English Chalk.

---

[1] The original spelling of the specific name " *pilula* " was " *pillula* ", as *Ananchytes pillula* Lamarck (1816 : 27). Brydone (1912 : 7, footnote) stated with reference to the spelling *pilula*, " this name was deliberately spelt thus in ' The Zones ' (Griffith & Brydone, 1911) in deference to Mr. Griffith's scruples as a classical scholar against countenancing the spelling ' *pillula* ' . . ".

Brydone divided the zone into a lower subzone of *Echinocorys scutata* var. *depressula* and an upper subzone of *Offaster pilula*. The *depressula* Subzone was not modified by Gaster, but for Brydone's *Offaster pilula* Subzone he (1924 : 94) substituted " subzone of *Echinocorys scutata* var. *cincta* ". In this subzone Gaster included the 60 ft. of Chalk comprising his " horizon of *Hagenowia rostrata* " ; the remainder of the subzone below it he termed the " horizon of abundant *Offaster pilula* ", almost equivalent to the three " belts " of the " subzone of *Offaster pilula* " of Brydone (1914*a*).

Brydone (1912 ; 1914*a*) and particularly Gaster (1924 : 103–107 ; 1930 : 329 ; 1937 : 326) established a sequence of " belts " characterized by different varieties of *Echinocorys scutata*. These are not shown in columns 4 and 5 of Text-fig. 1 but are summarized in tabular form by Gaster as mentioned above. The thickness of the zone of *Offaster pilula* was estimated by Withers (1935 : 59) as about 160 ft.

In Yorkshire, Chalk equivalent to the combined zones of *Offaster pilula* and *Gonioteuthis* [*Actinocamax*] *quadrata*, is characterized by the zonal form *Inoceramus lingua* with, at the top of the zone, a subzone of *Scaphites binodosus* regarded as the equivalent of the *Gonioteuthis* [*Actinocamax*] *quadrata* Zone of south-east England (C. W. & E. V. Wright, 1942 ; 1951 : 4). Peake & Hancock (1961) have not established this zone in Norfolk.

Zone of *Gonioteuthis* [*Actinocamax*] *quadrata*

Rowe's (1900) zone of *A. quadratus* was divided by Griffith & Brydone (1911) into three subzones. In 1912 Brydone combined the two lower subzones as the zone of *Offaster pilula* and restricted the *Actinocamax quadratus* Zone to the Chalk between his new zone and the base of the *Belemnitella mucronata* Zone. He included at the base of his restricted zone of *Actinocamax quadratus* about 60 ft. of Chalk which, in Sussex, Gaster (1924 : 93) regarded as the uppermost part of the underlying zone of *Offaster pilula* [i.e., the horizon of *Hagenowia rostrata*]. Gaster's arrangement was disputed by Brydone (1939) who re-stated his earlier opinion that the boundary between the zones of *Offaster pilula* and *Actinocamax quadratus* should be placed at the top of his " *Planoconvexus* Bed ".

Gaster (1924 : 93) recognized a subzone of *Saccacoma cretacea* for the upper 125 ft. of his restricted zone of *Actinocamax quadratus* in Sussex. *Saccacoma cretacea* is not here regarded as a true subzonal form : the general thickness of the restricted zone of *Gonioteuthis* [*Actinocamax*] *quadrata* was given by Withers (1935 : 59) as about 150 ft., so that *Saccacoma cretacea* extends through most of the zone. Similarly, the polyzoan species *Bicavea radiata* Gaster and *B. striata* Gaster occur throughout the whole zone and are not confined to separate horizons within it.

Zone of *Belemnitella mucronata*

The varying opinions on the position of the upper boundary of this zone are mentioned in the discussion of the overlying *Ostrea lunata* Chalk.

Apart from the inclusion, by some authors, of the *O. lunata* Chalk (e.g., Withers, 1935 : 60), and the Parapachydiscan divisions proposed by Spath (1926), the zone of *Belemnitella mucronata* has not been generally subdivided.[1] The position of the lower boundary of the zone of *Belemnitella mucronata* as established, for example, by Rowe (1900), has not been queried.

Variations in thickness are very marked : in a general estimate Withers (1935 : 60) gave the thickness as 250 ft. (excluding the *Ostrea lunata* Chalk) in Norfolk, 70 ft. in Hampshire, 250 ft. in Dorset, and 475 ft. in the Isle of Wight. The thickness in Norfolk has been more accurately estimated by Peake & Hancock (1961) as 325 ft.

Uneven pre-Tertiary or recent erosion probably accounts for much of this variation, but the thickness which remains is great and more general subdivision would be of value.

Maastrichtian

Zone of *Belemnella lanceolata*

In his original publication on the stratigraphy and fauna of the Trimingham Chalk, Brydone (1900 : 11) concluded that " the great majority of forms . . . which appear in this [faunal] list . . . are characteristic of the Maastrichtian Stage on the Continent ". Subsequently, Brydone (1908*a*, *b*) described the Trimingham Chalk in detail. Since that date the Trimingham Chalk has been placed, by various authors, either as highest Senonian or as Maastrichtian in age.

Boswell (1929 : 406, 408A) recognized Jukes-Browne's (1904 : 5, 12) zone of *Ostrea lunata* and regarded it as comprising " higher beds than those of the *B. mucronata* zone ".

Withers (1935 : 40) discussed the limits of the " Maestrichtian and Upper Senonian (Campanian) " and some of the previous uses of the term " Maestrichtian ". He stated that it " was originally applied by Dumont . . . to include the Tuffeau de Maestricht as exposed at Maestricht, Holland, and [that] this term corresponds to the Dordonian of Coquand ". He also mentioned Haug's (1910) extension of the term Maastrichtian to include the " Maestrichtian (*s. str.*) and the zone of *B. mucronata* ", and Spath's (1926) modification which included with the Maastrichtian only " the uppermost part of the *B. mucronata* zone ".

Withers himself (1935 : 41) applied the term " Maestrichtian . . . in its original sense to beds of the same age as the Tuffeau de Maestricht ", and retained as " Upper Senonian (Campanian) " the " Norwich Chalk ", the Skrivekridt of Denmark and the Schreibkreide of Rügen. The Chalk of Trimingham, Norfolk, he regarded (p. 60) as being " a little higher in the sequence of the *B. mucronata* zone . . . than that occurring in the neighbourhood of Norwich."

---

[1] Wright (1947) mentioned " Lower, Middle and Upper Divisions " of the *Belemnitella mucronata* Zone in Dorset, the Middle Division being characterized by distinctive asteroid remains. C. W. & E. V. Wright (1951) adopted, in a slightly modified form, Spencer's (1913 : 145) division of the " *mucronata* " Chalk into Upper, Middle and Lower zones. They stated (1951 : 3) that of these divisions, " the first [topmost] is now the zone of *Belemnella lanceolata*, and the other two are treated as subzones, as yet unnamed, of a restricted zone of *Belemnitella mucronata* ".

The " Maestrichtian " was suggested by Brydone (1938 : 16) as a " possible continental equivalent " of the " Trimingham General Chalk ", i.e., his upper division of the zone of *Ostrea lunata.*

More recently, Jeletzky (1951), in a comprehensive study of the use of the name Maastrichtian, considered that the Trimingham Chalk is of Lower Maastrichtian age, an opinion based largely on a study of the belemnite faunas of Trimingham and other Maastrichtian horizons on the Continent. The highest *Belemnitella mucronata* Chalk he regarded as that of the Norwich district, which he retained in the Senonian (see Jeletzky, 1951, table facing p. 208).

Jeletzky's interpretation of Dumont's (1849) use of the term Maastrichtian has been questioned by van der Heide (1954). He stated (p. 509) that Dumont " considered the Maastrichtian Chalk and its equivalents as belonging to his Système Maestrichtien : older beds were not included ". After a careful and detailed assessment of the evidence from Dumont's pre-1849 publications, his maps, and Mourlon's publication (1878) based on Dumont's notes and manuscripts (Dumont died in 1857), van der Heide concluded that " the Système Maestrichtian of Dumont (1849) *is equivalent to the Maastricht Chalk* " of the type locality only, and that the newer interpretations, which enlarge the Maastrichtian (e.g. Jeletzky, 1951) are incorrect. Finally, van der Heide stated (p. 511) " the only conclusion at which we can arrive at the moment is that a detailed study of the stratigraphy and palaeontology of the type area of the Maastrichtian is indispensable ".

It is desirable to establish, as van der Heide has done, the original meaning of a stratigraphical term, but this should not then preclude its use in a wider sense—provided that the extended use is also clearly defined.

Recently Jeletzky's enlarged use of the term Maastrichtian has been applied, with various modifications, by several continental authors. For example, Voigt (1951) has described polyzoan faunas of Lower and Upper Maastrichtian age from Holland and Germany ; the zone of *Belemnitella mucronata* being retained in the Upper Campanian. This is in agreement with the extensive work of Hiltermann & Koch (1950), Hiltermann (1952) and Wicher (1949, 1953*a*, 1953*b*) which has shown that the top part of the previously so-called " Upper Senonian " of Denmark, Germany and other areas, may be regarded as a widespread boreal facies of the Maastrichtian (*s. lato*) characterized by a distinctive foraminiferal fauna.

In a study of the foraminifera of the English Chalk Williams-Mitchell (1948) indicated that the foraminifera of the Trimingham Chalk differ considerably from those which characterize the underlying Senonian horizons. Barnard & Banner (1953 : 207) listed 24 species of arenaceous foraminifera which occur in the zone of *Belemnitella mucronata* ; of these, 6 are long-range forms occurring from the Cenomanian to the zone of *Ostrea lunata*, 5 range from the base of the zone of *Belemnitella mucronata* into the *Ostrea lunata* Chalk, and the remaining 13 species are all confined to the zone of *Belemnitella mucronata*, again indicating a change at the top of the latter zone.

A sample collected in 1954 from a foreshore exposure of white chalk with *Ostrea lunata*, just north of Mundesley, was prepared and examined for its foraminiferal content by the late Mr. A. G. Davis. In a letter (20th September, 1954) he wrote

" the foraminiferal content of this sample is quite unlike any from the zones below —which are much better known. I find the foraminifera fauna is characterized by some species of *Bolivina*, species much like *B. incrassata*, *B.* " *gigantea* " and *B. decussens*. There are also some arenaceous agglomerated forms—as also mentioned by Williams-Mitchell.

The species of *Bolivina* present in the sample are characteristic of the Maastrichtian of Hemmoor and Lüneburg. They are rare in the *mucronata* Zone, but do make an appearance and only become abundant higher up. They are abundant in the Mundesley sample.

Other foraminifera are present, but not abundant, and they are of long range. The well-known Chalk foraminifera, Nodosarids, *Globotruncana*, etc., do not seem to be present."

This evidence of a faunal change in the foraminifera is further supported by the distribution of species of *Castanopora* (see p. 274).

Boswell (1923) estimated the total thickness of the *Ostrea lunata* Chalk at about 110 ft., and this is confirmed by Peake & Hancock (1961 : 320) in their precise study of the Trimingham Chalk. The boundary between it and the underlying zone of *Belemnitella mucronata* has not been seen.

C. W. & E. V. Wright (1951 : 3) have stated that " *Ostrea lunata*, the fossil used by Jukes-Browne as an index for the uppermost zone of the Chalk (at Trimingham), is not entirely appropriate [as a zonal name form], as it does not occur at all in a large part of the Trimingham Chalk. Now that Jeletzky has shown that the vast majority of the Trimingham belemnites are *Belemnella lanceolata* (Schlotheim) and not *Belemnitella mucronata* the former species can appropriately be used as a zone fossil, as it is over most of northern Europe ". Thus the authors referred (1951 : 3) to a zone of *Belemnella lanceolata* as an alternative to " zone of *Ostrea lunata* ". They also used the terms " Upper " and " Lower Senonian " as stages embracing the zones of *Ostrea lunata* down to, and including, the zone of *Offaster pilula*, and from the zone of *Marsupites testudinarius* down to, and including, the zone of *Micraster cortestudinarium* respectively. They stated that the " Upper Senonian includes Campanian and Maastrichtian. Lower Senonian includes Coniacian and Santonian ".

Foreign Upper Cretaceous Horizons

Most of the Polyzoa described here are from the English Chalk, but a number of specimens from various horizons of the Upper Cretaceous abroad have also been examined in order to complete the systematic revision of some genera.

The following list of genera and species from such localities and horizons are included in the systematic revision of the Cretaceous cribrimorph Polyzoa described here. When no specimens are available, either in the collections of the British Museum (Natural History), or in the other collections studied, the species have been assessed from published works only. In these cases the name of the species is followed in the list by the reference to its original description [e.g., *Castanopora* [*Rhiniopora*] *labiata* (Levinsen) (1907 : 155, 156, 158, pl. opp. p. 160, fig. 1, 1*a*–*f*). Møen, Aalborg, Denmark. Lower Maastrichtian (as Senonian, zone of *Belemnitella mucronata*)]. Where specimens of species from foreign localities are available the name of the species is marked by an asterisk and is followed directly by its locality and horizon [e.g., **Castanopora glandulosa* Lang. Rügen. Probably boreal Maastrichtian (as Senonian, zone of *Belemnitella mucronata*)]. If a photograph only is available this is stated. Published records of the foreign occurrences of other species are also mentioned.

AEOLOPORA

Aeolopora nebulosa Lang. Also recorded by Voigt (1930 : 498, 542, 560, pl. 27, fig. 1) from Ifö, Sweden. " Mammillatensenon."

CASTANOPORA

Castanopora [*Rhiniopora*] *aviculosa* (Lang). Photograph only of the holotype. Maastrichtian, Maastricht, Limbourg, Holland.

**C. castanea* Lang. Rügen. Also recorded by Voigt (1930 : 513, 561, pl. 30, fig. 17) from the same locality and horizon. Probably boreal Maastrichtian [as Senonian, zone of *Belemnitella mucronata*].

**C. dibleyi* (Brydone). Also recorded by Voigt (1930 : 512, 561, pl. 30, fig. 18) from Rügen. Probably boreal Maastrichtian, and ? Voigt (1930 : 512, 561, pl. 30, fig. 20) as *Castanopora* sp. from Stafversvad, Schonen [as " Unteres Mukronatensenon "].

C. faujasi (von Hagenow) (1851 : 99, pl. 10, fig. 19*a*, *b*). Also recorded by Voigt (1930 : 511, pl. 31, fig. 2) as *Rhiniopora faujasi* (von Hagenow), Maastricht, Limbourg, Holland. Maastrichtian [as " Ob. Mukronatensenon "].

**C. glandulosa* Lang. Rügen. Probably boreal Maastrichtian [as Senonian, zone of *Belemnitella mucronata*].

C. guascoi (Ubaghs) (1865 : 51, pl. 2, figs. 3*a–c*). Valkenburg (Fauquemont), east of Maastricht, Limbourg, Holland. Maastrichtian. Also recorded by Voigt (1930 : 512, 561, pl. 31, fig. 1). Maastricht. Maastrichtian [as " Ob. Mukronatensenon "].

**C.* [*Rhiniopora*] *jurassica* (Gregory). Recorded incorrectly from Ranville, Normandy, France, as of Jurassic age. Very probably from Maastricht, Limbourg, Holland. Maastrichtian.

C. [*Rhiniopora*] *labiata* (Levinsen) (1907 : 155, 156, 158, pl. opp. p. 160, fig. 1, 1*a–f*) = *Cribrilina valida* Levinsen (1925 : 376, pl. 5, fig. 58*a–d*). Møen, Aalborg, Denmark. Lower Maastrichtian, [as Skrivekridt, Senonian, zone of *Belemnitella mucronata*].

**C. magnifica* (d'Orbigny). Holotype—photograph only, either from Sainte Colombe, Manche, France, or from Royan, Charente Inférieure, France. Both localities Maastrichtian. [As Senonian, zone of *Belemnitella mucronata.*] Includes *Rhiniopora aspera*, *R. asperula*, *R. cacus* and *R. horrida*. Recorded by Voigt (1930 : 511) from Rügen and Blauen Bergen, Roslau. Probably boreal Maastrichtian [as " Ob. Mukronatensenon "]. Also recorded by Illies (1953) from Hemmoor, near Hanover, Germany. Maastrichtian.

C. multicostata Voigt (1930 : 512, 561, pl. 30, fig. 19). Rügen. Probably boreal Maastrichtian [as " Ob. Mukronatensenon "].

C. [*Rhiniopora*] *spooneri* (Butler & Cheetham) (1958 : 1154, text-figs. 1, 2). "... from a chalk bed exposed at Rayburn's dome, Bienville Parish, Lousiana ... equivalent to the Upper Cretaceous Saratoga chalk of Arkansas ..." U.S.A.

C. voigti sp. nov. (Voigt, 1930 : 511, pl. 31, fig. 6—as " *Rhiniopora hispida* Lang " which is indeterminate, see p. 238). Møen. Denmark. Lower Maastrichtian [as Oberes Mukronatensenon], and Rügen, probably boreal Maastrichtian.

HEXACANTHOPORA

**Hexacanthopora sexspinosa* Lang. Also recorded by Voigt (1930 : 495, 542, 560, pl. 26, figs. 11, 12) from " Gr. Bülten-Granulatensenon " (as *H. brightonensis*) and " Quadraten- oder Mukronatensenon " of Misburg, Germany.

H. viginticostata Voigt (1930 : 495, 560, pl. 26, fig. 13 ; 1949 : 37, 43, pl. 9, figs. 1, 2). Schwiecheldt, Hanover, Germany. " Quadraten- oder Mukronatensenon ", and Lägerdorf, Holstein, Germany, Senonian, zone of *Gonioteuthis* [*Actinocamax*] *quadrata.*

LAGYNOPORA

**Lagynopora furcifera* (Brydone). Also recorded by Voigt (1930 : 495, 542, 560, pl. 26, fig. 10 ; 1949 : 36, 43, 44, pl. 11, fig. 1). Misburg, Germany—" Quadraten- oder Mukronatensenon ", and Lägerdorf, Holstein, Germany—Senonian, zone of *Gonioteuthis* [*Actinocamax*] *quadrata.*

**L. horsleyensis* Lang. A poorly preserved specimen has been recorded by Didon (1958 : 80, 84, text-fig. 2) from the Santonian of Le Trait, near Rouen, Seine Inférieure, France.

L. lagena Lang. Also recorded by Voigt (1930 : 495, 542, 560, pl. 26, fig. 9). Balsburg. " Mammillatensenon."

?*L.* [? = *Canupora*] *prima* (Kühn) (1930 : 65, pl. 26, fig. 13). Lithothamnienkalk. Haidhof, Ernstbrunn, Austria. Danian.

LEPTOCHEILOPORA

**Leptocheilopora filliozati* (Brydone). Also recorded by Gillard (1943 : 187). "... à la côte—72 m. par le forage Font-de-Chèvres ... Royan ", France. Either Senonian, zone of *Belemnitella mucronata*, or Maastrichtian.

**L. magna* Lang. Holotype and paratype—Lüneberg, Hanover, Germany—very probably boreal Maastrichtian [as Senonian, zone of *Belemnitella mucronata*]. Other specimens—Hemmoor, Hanover, Germany—Maastrichtian, either zone of *Belemnella lanceolata* mut. *sumensis* or zone of *B. junior.* Also recorded by Voigt (1930 : 496, pl. 27, fig. 5) ? Maastricht. Maastrichtian [as " Obere Mukronatenkreide "], and from Rügen, probably boreal Maastrichtian. The species is synonymous with *Barroisina galeata* Levinsen (1925 : 385, pl. 6, fig.

67). Møen, Aalborg, Denmark. Lower Maastrichtian [as Skrivekridt, Senonian, zone of *Belemnitella mucronata*]. The species is probably synonymous also with *Leptocheilopora longuesensis* Lang (1916*a* : 396, 397 ; 1921 : 76)—horizon recorded as " Campanian, Longuesse, France ".

PELMATOPORA

**Pelmatopora brydoneis* [*sic*] Lang. Recorded by Didon (1958 : 83, 85, text-fig. 10) from the Santonian of Elbeuf, near Rouen, Seine Inférieure, France.

**P. chrysalis* (d'Orbigny). Photograph only. Villedieu, France. Senonian, Coniacian. Also recorded by Gillard (1942 : 90) from the Upper Chalk between 24 m. 50 and 26 m. 80 (Division D) from a boring at Montbazon, south of Tours, Indre-et-Loire, France.

P. ? daniensis Voigt (1925*b* : 99, 100, pl. 3, figs. 4–6 ; 1930 : 514, 561, pl. 31, fig. 11). Cöthen. Danian.

**P. fecampensis* Lang. Holotype and other specimens—Fécamp north-east of Le Havre, Seine Inférieure, France. Senonian, Coniacian. Also recorded by Didon (1958 : 82, 84, text-fig. 9) from the Santonian of Le Trait, Elbeuf and Orival, near Rouen, Seine Inférieure, France.

**P. fragilis* (d'Orbigny). Lectotype—the specimen figured by d'Orbigny (1851, pl. 717, fig. 9). Fécamp, north-east of Le Havre, Seine Inférieure, France. Senonian, Coniacian. Photograph only of a specimen from the same locality and horizon as the lectotype, and a specimen from St. Avertin, south-east of Tours, Indre-et-Loire, France. Senonian, Coniacian. Also recorded by Gillard (1942 : 90) from the Upper Chalk between 6 m. 50 and 7 m. 30 (Division A) from a boring at Montbazon, south of Tours, Indre-et-Loire, France.

**P. larva* Lang. Holotype and D.28438—La Ribochère, Loir-et-Cher, east of La Chartre-sur-le-Loire, Sarthe, France. Senonian, Coniacian.

**Pelmatopora* cf. *marsupitum* [*sic*] Lang. Recorded by Didon (1958 : 83, 85, text-fig. 11) from the Santonian of Moulineaux, near Rouen, Seine Inférieure, France.

**P. d'orbignyi* Lang. Holotype. St. Avertin, south-east of Tours, Indre-et-Loire, France. Senonian, Coniacian. Specimen also from La Ribochère, Loir-et-Cher, east of La Chartre-sur-le-Loir, Sarthe, France. Senonian, Coniacian.

**P. quadrata* Lang. Synonymous with *P. filliozati* Lang. Photograph only of the holotype of *P. filliozati* Lang, from Fécamp, north-east of Le Havre, Seine Inférieure, France. Senonian. Coniacian.

**P. striata* (d'Orbigny). Lectotype—the specimen figured by d'Orbigny (1851, pl. 686, fig. 10). Nehou, south of Valoignes, east of Sainte Colombe, Manche, France. Senonian. Photographs only of a specimen from Villedieu, France. Senonian, Coniacian. Also recorded by Gillard (1942 : 90) from the Upper Chalk between 24 m. 50 and 26 m. 80 (Division D) from a boring at Montbazon, south of Tours, Indre-et-Loire, France.

**P. suffulta* (Brydone). Canu (1911 : 252, pl. 6, figs. 7–10) has also recorded " *Cribrilina insignis* " which is possibly synonymous with *Pelmatopora suffulta*. Roca, Argentine. Rocanean.

UBAGHSIA

Ubaghsia aurita Lang. Photograph only of the holotype. Royan, south of Rochefort, Charente Inférieure, France. Maastrichtian.

**U.* [*Batrachopora*] *coaxans* (Lang). Rügen. Probably boreal Maastrichtian [as " Upper Senonian "].

**U.* [*Batrachopora*] *crassa* (Lang). Rügen. Probably boreal Maastrichtian [as Senonian, zone of *Belemnitella mucronata*].

U. [*Steginopora*] *demorgani* (Jullien) (Jullien, 1886 : 615, pl. 19, figs. 4, 5, pl. 20, fig. 1). Les Molineaux, Meudon, south-west of Paris, France. Senonian, zone of *Belemnitella mucronata.*

U. [*Steginopora*] *meudonensis* (Jullien) (Jullien, 1886 : 614, pl. 17, figs. 1–5, pl. 18, figs. 1–3). Details as for *U. demorgani.*

U. [*Steginopora*] *ocellata* (Jullien) (Jullien, 1886 : 614, pl. 19, figs. 1–3). Port Brehay, north-east of Gourbesville, north-west of Carenten, Manche, France. " Etage sénonien "—probably middle or lower part of zone of *Belemnitella mucronata.*

**U.* *ornata* (Goldfuss). Lectotype—the specimen figured by Goldfuss (1826, pl. 9, fig. 1*b*). St. Petersburg, Maastricht. Maastrichtian. Specimens from Valkenburg (Fauquemont) and Maastricht, Holland. Maastrichtian. Also recorded by Voigt (1930 : 513, pl. 31, fig. 20 only). Maastricht. Maastrichtian [as " Ob. Mukronatensenon "].

U. [*Batrachopora*] *perforata* (Marsson) (Marsson, 1887 : 98, 109, pl. 10, fig. 11). Rügen. Probably boreal Maastrichtian [as " Schreibkreide "]. Also synonymous with *Cribrilina brachiata* Levinsen (1925 : 377, pl. 8, fig. 24). Møen, Aalbourg, Denmark. Lower Maastrichtian [as Senonian, zone of *Belemnitella mucronata*].

**U.* [*Batrachopora*] *ranunculus* (Lang). Rügen. Probably boreal Maastrichtian [as Senonian, zone of *Belemnitella mucronata*]. Also recorded by Voigt (1930 : 515, 561, pl. 31, fig. 17) from the same locality and horizon.

**U.* *reticulata* (Ubaghs). Lectotype—the specimen figured by Ubaghs (1865, pl. 2*a*, fig. 7*c*). Valkenburg (Fauquemont) or Geulem, Maastricht, Holland. Maastrichtian, " Ober Bryozoenschichte " (horizon *f* of Ubaghs). Specimens from Maastricht, Holland. Maastrichtian. Also recorded by Voigt (1930 : 515, pl. 31, figs. 18, 39 only). Maastricht, Holland. Maastrichtian [as " Ob. Mukronatensenon "].

The age of some Upper Cretaceous horizons abroad, e.g. that of the Rügen Chalk, has more recently been proved different from that originally stated by some authors. In the above list such revision of horizons is indicated by giving the old opinion enclosed in square brackets.

Among the stratigraphical terms used in the above list " Coniacian " refers to beds equivalent in age to the zone of *Micraster cortestudinarium*, and " Santonian " to beds equivalent in age to the zones of *M. cor-anguinum* and *Marsupites testudinaris.* The following terms are revised or generally in use by foreign authors.

Granulatensenon. Chalk with *Gonioteuthis granulata* in Germany, approximately equivalent to the zone of *Marsupites testudinarius* in England.

Lithothamnienkalk of Haidhof. Part of the Austrian Upper Cretaceous series of Danian age.

Maastrichtian. The general use of this term as applied to English and foreign horizons has been discussed (p. 12).

Mammillatensenon. Part of the Trümmerkreide of Sweden, equivalent to the upper part of the zone of *Gonioteuthis* [*Actinocamax*] *quadrata* in Germany and England. Possibly also equivalent to part of the zone of *Belemnitella mucronata* in England (see Withers, 1935 : 71).

Møen Chalk, Aalborg, Denmark. Referred by Levinsen (1925) and others to the Senonian, zone of *Belemnitella mucronata*, but now known to be Lower Maastrichtian.

Oberes Mukronatensenon of Maastricht. Used by Voigt (1930) for the Cretaceous of the type locality of the Maastrichtian, termed subsequently by Voigt, and generally, Maastrichtian *s. str.*

Rocanean. Part of the Argentine marine Upper Cretaceous. Feruglio (1949 : 218) gave an extensive survey of previous studies of the Rocanean and concluded that " after various considerations and a critical study of the data hitherto published by previous authors it is impossible, in the present state of knowledge, to decide whether these strata belong to the Upper Cretaceous or to the lower part of the Tertiary " [translation].

Rügen Chalk. Originally (e.g. Lang, 1921 ; 1922) regarded as Senonian, zone of *Belemnitella mucronata*, but subsequently shown (e.g. Voigt, 1951 ; Hiltermann, 1952) to be probably boreal Maastrichtian.

III. NOTES ON THE COLLECTIONS STUDIED

British Museum (Natural History)

Most of the material described here is from the large collections of Chalk Polyzoa made by C. T. A. Gaster and presented by him in recent years. The total of Chalk Polyzoa in the Gaster Collections numbers many thousands, the majority of specimens coming from south-east England. Much of the material has been carefully cleaned, stained and mounted by the collector, and all specimens are accompanied by full and precise details of locality and horizon.

A number of specimens have been studied which were obtained by earlier collectors. Such specimens, from collections other than the Gaster Collection, were incorporated by Lang in his works on Cretaceous cribrimorph Polyzoa and have been re-examined here to complete the revision of some genera.

The British Museum Collections also include photographs of specimens in the F. Canu Collection, Paris.

Sedgwick Museum, Cambridge

The R. M. Brydone Collection of Chalk Polyzoa in the Sedgwick Museum, Cambridge, includes many very well preserved cribrimorph Polyzoa. I have examined the type specimens of the species established by Brydone which belong to the genera revised here.

Norwich Castle Museum

All available specimens of cribrimorph Polyzoa in the R. M. Brydone Collection of Chalk fossils from Norfolk and adjacent areas have been examined. Where appropriate, these, and specimens from other collections in the Norwich Castle Museum, are incorporated in the present study.

IV. PRESERVATION OF THE MATERIAL

Cretaceous cribrimorph Polyzoa are often well preserved. They may, however, have been worn before interment so that projecting structures, such as avicularia, oral spines and median processes of the apertural bar, are eroded and rounded. Usually it is possible to find, in a zoarial fragment of average size, an area in which the zooecia are well preserved—often through being at a slightly lower level than the general surface of the zoarium and thus escaping the general effects of wear.

In contrast, some zoarial fragments entirely lack fine skeletal details, but retain, unbroken, projecting structures such as avicularia. General fine wear of this kind was possibly produced by solution effects on the surface of the skeleton before burial, rather than by mechanical abrasion. Phleger, Parker & Peirson (1953 : 117–119, pl. 12) have described and figured a number of Recent foraminifera with calcareous tests which exhibit similar primary solution effects. As only a few zoarial fragments of Cretaceous Polyzoa, from any given locality, may show such solution effects, it seems probable—as with the Recent foraminifera—that solution was predepositional rather than secondary.

The secondary cause of loss of detail in Chalk Polyzoa is recrystallization of the calcium-carbonate of their skeletons. This seems to have occurred more commonly in specimens from earlier horizons of the English Chalk.

Most polyzoan skeletons from the Chalk are composed of finely crystalline calcium-carbonate, but some may be perfectly replaced secondarily by silica. Siliceous specimens are usually very well preserved and may occur in the hollow centres of flint nodules forming part of the infilling " flint meal ".

V. TECHNIQUES

All the material has been examined with strong reflected light at high- and low-power magnifications under a binocular microscope. Lang's method of staining specimens with indigo water-colour has been used extensively (Lang, 1921 : lxxv). Without staining, the detailed characters, such as the type and number of pelmata, are hardly visible, but the careful application of a fairly dark, temporary stain brings out such details and differentiates the polyzoan skeleton from any adherent Chalk matrix.

Many specimens encrust fragments of echinoid test or pieces of *Inoceramus* shell. With such specimens, soaking in water and gentle brushing with a soft water-colour paint brush is usually an adequate cleaning method. Unattached zoarial fragments may be freed from their Chalk matrix by repeated soaking in water and gentle brushing.

The matrix may be harder and more closely adherent to the polyzoan skeletons. In these cases, soaking for a short time (10–30 minutes) in a saturated solution of salicylic acid was found to be sufficient to loosen the matrix without damaging the polyzoan. This treatment was always followed by careful washing in running water for 24–36 hours, the material being finally dried out very slowly. Specimens treated in this way have shown no signs of deterioration after five years. The material can be stained in the usual way.

Where possible, the specimens have been mounted with gum tragacanth on mahogany microscope slides—the system used generally in the Department of Palaeontology of the British Museum (Natural History).

Accurate drawings of specimens have been made on koda-trace mounted on squared paper using a micrometer scale and squared graticule in the eyepiece of the microscope. All holotypes and some other specimens have been photographed by reflected light generally at a magnification of ×20, but occasionally at higher magnifications up to ×100 to show minute detail.

VI. MORPHOLOGY OF CRETACEOUS CRIBRIMORPH POLYZOA

Zoarial Characters

GROWTH HABIT. Many Cretaceous cribrimorph Polyzoa have encrusting zoaria firmly attached to a substrate such as an echinoid test or the shells of other organisms. Unattached zoarial fragments are also common, but many of these may have been encrusting during life on some perishable object which was destroyed during or after fossilization. Thus the distinction often made between encrusting and so-called " free " forms is very uncertain. Bassler (1953 : G7) pointed out that " although often described as free, only the upper branches actually are free, for most bryozoans are attached basally or by other parts of their surface to extraneous objects . . .".

Encrusting zoaria are usually unilaminar, consisting of a relatively widespread, single layer of zooecia attached dorsally to a substrate. Much less commonly, successive layers of zooecia may be superimposed producing a multilaminar zoarium of hemispherical or nodular form : some species of *Polycephalopora* (Lang, 1922: 105) are of this type.

Unattached zoaria may be uni-, bi- or multilaminar, and, as fossils, are almost invariably fragmentary. Unilaminar fragments may originally have encrusted a destructible substrate since lost. Their dorsal surfaces may be fairly even, with the basal wall of each zooecium smooth, slightly convex, and separated from adjacent zooecia by a slight furrow ; or they may be more irregular, the basal wall of each zooecium possessing small, scattered tubercles, with deeper furrows between adjacent zooecia, as in *Pelmatopora gregoryi* (Brydone), *P. somptingensis* Lang (Pl. 15, fig. 4) and *P. brydonei* Lang (Text-fig. 77). Some Recent Polyzoa are attached to destructible substrates by means of tubercles protruding from their dorsal surfaces, and so their presence in fossil forms may indicate an originally adherent zoarium.

Some zoaria are bilaminar, consisting of two layers of aligned zooecia growing back to back, their dorsal surfaces united. Bilaminar zoaria may be irregularly lobed expansions in one plane, or more complex, foliaceous and undulating. From very small fragments of fossil bilaminar zoaria it is usually not possible to decide what was the original form of whole colonies, though they were presumably like Recent bilaminar polyzoan colonies. Some zoaria of *Pelmatopora larva* Lang (p. 134) are bilaminar.

Unattached fragments of multilaminar zoaria occur, but these may have been broken from encrusting masses.

Castanopora voigti sp. nov. (p. 236) has an erect, cylindro-quadrate zoarium in which four adjacent rows of zooecia alternate : this zoarial form is apparently rare among cribrimorph Polyzoa. Other erect, cylindrical zoaria occur occasionally, as in *Pnictopora suffocata* Lang and *P. strangulata* Lang.

Some genera, now largely archaic, were based on zoarial growth forms alone. There is often a considerable amount of variation in the growth habit of zoaria of the same genus or species. However, Lang (1916*a* ; 1921 ; 1922) frequently used zoarial growth habit to distinguish species which are otherwise morphologically identical. In the material revised and described here, *Pelmatopora* affords several examples of species established by Lang on zoarial growth form alone. Thus he distinguished (1922 : 300) as *P. roedeanensis*, a species differing " only in colonial habit from *P. marsupitum* [*sic*]." *P. lacuum* Lang was described (1922 : 324) as " *Pelmatopora ranunculoides*, but with an erect unilaminar asty [zoarium] ", and *P. promontoriorum* Lang (1922 : 320) as having a structure which " resembles that of *Pelmatopora saltdeanensis*, but the asty is erect instead of incrusting ". Again, Lang stated (1922 : 308) " *Pelmatopora damicornis* differs from *P. gregoryi* and *P. palmata* in its colonial habit, which is bilaminar. The three species thus form a progressive series with regard to colonial habit and condition, culminating in the erect bilaminar *P. damicornis* ".

Further examples may be cited, from genera not revised here. Among these are *Sandalopora lavardinensis* Lang distinguished (1922 : 330) on zoarial growth form from *S. gallica* Lang, *S. suppulosa* Lang separated (1922 : 331) on the same basis from *S. crepidata* Lang and three species of *Rhabdopora* (*R. virgata*, *R. virgulata* and *R. tigrina*) distinguished (Lang, 1916*a* : 405 ; 1921 : 209) in the same way.

Lang was aware of the uncertainty of describing unattached unilaminar zoarial fragments as " erect ". He stated (1919*d* : 196) " It is hard to be certain . . . that an apparently erect unilaminar asty is not really incrusting, since an alga or other perishable substance may have formed its support . . . it is unusual, however, to find an habitually-incrusting form in an apparently free condition, and, unless there are other reasons against it, I should advocate keeping incrusting and free, unilaminar forms specifically distinct ". Lang preceded this statement with the assertion that " in the Pelmatoporinae, and throughout the phylum, the direction of evolution is from an incrusting to an erect asty ".

In some Recent Polyzoa it is possible to find species, such as *Steganoporella buski* Harmer, in which different parts of the same zoarium are encrusting or erect and uni- or bilaminar and, in some parts, even multilaminar. On such evidence, and in view of the uncertainty of establishing whether or not an unattached, unilaminar zoarial fragment had an encrusting mode of life, I do not agree with Lang's division of species on zoarial growth habit alone. On the same evidence, the suggested " direction of evolution " from encrusting to erect zoaria must also be questioned.

ZOARIAL BUDDING. In an encrusting zoarium the zooecia may be developed uni- or multiserially. In Cretaceous cribrimorph Polyzoa the former condition is much rarer than the latter.

Uniserial encrusting zoaria. Some genera, such as *Corymbopora* and *Andriopora* (p. 266), are always uniserial, and the zooecia may possess caudae, as in *A. major* sp. nov. (p. 268). Generally, uniserial encrusting zoaria are more extensive than multiserial, though many more zooecia may be present in a multiserial zoarium which has developed uninterruptedly on a wide substrate. The angle of divergence of lines of uniserially budded zooecia may vary considerably.

Multiserial encrusting zoaria. These are usually fan-shaped, tending ultimately to become circular as more and more zooecia are developed. The sequence of growth appears to be like that observed by Marcus (1926) [summarized by Harmer, 1931 : 122] in the Recent cheilostome polyzoan *Electra pilosa* (Linn.). Here three distal buds arise from the ancestrula producing the first generation of zooecia. Successive generations are budded off producing a young fan-shaped colony, the distal-lateral edges of which are early reflected and usually meet on the proximal side of the ancestrula. Where zooecia are crowded, usually only one bud is produced, and in later stages there is a tendency to form adjacent, but distinct, uniserial, radiating rows of zooecia (e.g., as in *Aeolopora distincta* Lang, Pl. 2, figs. 1, 2). Where sufficient space is available two or three buds may arise in *Electra pilosa*, and a new radiating row of zooecia is intercalated in the colony. This feature is commonly found in Cretaceous cribrimorph Polyzoa.

In an adult zoarium, which has attained a more or less circular form, new zooecia are budded off from the periphery. This growing edge is seldom preserved in the Cretaceous cribrimorph Polyzoa ; presumably zooecia which have only just developed are more fragile than the preceding established, probably more calcified, adult zooecia.

As a multiserial zoarium matures, particularly if its substrate is limited, the later stages of zoarial growth may obscure the ancestrular area, often by the development of secondary calcareous tissue. Frequently one part of a zoarium may develop more rapidly than another, and, especially where there is a limited substrate, two zoarial expansions may converge, producing distortion of zooecia where they meet.

Bilaminar, erect, branching or foliaceous zoaria are apparently always multiserial. They may exhibit some of the features described for encrusting zoaria, but the modifications imposed by a limited substrate are usually absent, and the zoaria develop uninterruptedly.

Zooecial Characters

Certain structural features are common to the zooecia of all Cretaceous cribrimorph Polyzoa. They all possess calcified proximal, distal and lateral walls, and a secondary front wall (the frontal shield) composed of costae. These over-arch an aperture which is divided by an apertural bar from a distal orifice with marginal oral spines. This basic arrangement may be complicated in various ways—lateral costal fusions may develop between adjacent costae; a median process, formed on the apertural bar, may fuse with the proximal pair of oral spines producing a proximal oral shield; or the oral spines may be enlarged and bifurcated. The basic morphology is often further obscured by a general accumulation of secondary calcareous material which may fill the interzooecial valleys and, in some genera, encroach on to the frontal shields, or which may accumulate round the primary orifice producing a tubular secondary orifice. Ovicells and various heterozooecia may also obscure the form of zooecia.

The following observations and remarks apply to adult zooecia, but many of the characters discussed are the same in younger stages of zoarial growth. Young zooecia and ancestrulae are discussed later.

Size of zooecia. In the Cretaceous cribrimorph Polyzoa, as in other divisions of the Cheilostomata, the size of zooecia varies greatly, but some genera, compared with others, may have consistently small zooecia. For example, the zooecia of *Aeolopora* are always much smaller than those of *Castanopora*. However, for a given species there may be considerable differences in size between zooecia of separate zoaria, and often between zooecia in the same zoarium. Zooecial size, therefore, should seldom be used as a basis for differentiating species in the absence of more definite characters. But Lang has described (e.g. 1921 : 224–227) as separate species, forms such as *Dishelopora binoculata* and *D. claviceps*, which differ only very slightly in their zooecial length.

Shape of zooecia. It is strictly correct to say that in the great majority of cribrimorph Polyzoa the zooecia are polygonal and not broadly oval or elliptical. This polygonal form is most easily seen on the dorsal surfaces of unattached unilaminar zoaria (as in *Pelmatopora somptingensis* Lang, Pl. 15, fig. 4), or in encrusting zoaria in which the frontal shields have been destroyed, revealing the vertical walls of the zooecia. On the front of the zoarium interzooecial secondary tissue may obscure the true limits of each zooecium, or the frontal shields of adjacent zooecia may appear almost contiguous as in the specimens of *Castanopora* shown in Pl. 17. The upper surfaces of zooecia may vary in shape from almost circular to almost quadrate, though they are most commonly longitudinally oval.

In a given zoarium, most zooecia have approximately the same shape, but a few usually differ, possibly because of growth restrictions imposed by a limited substrate. Apparently, if only a limited amount of space is available for development, a zooecium tends to fill that space—its shape being to some extent controlled by it.

Zooecial Walls other than the Frontal Wall

Basal walls. A distinction may be made between the zooecia of some unilaminar encrusting zoaria and those which are now unattached. In the former, where the frontal shield is removed by wear, or artificially, there is often no indication of a calcified basal wall. This is shown in broken zooecia of *Lagynopora* and *Hexacantho-*

pora (Pl. 5, figs. 1, 4). In contrast, zooecia of unilaminar encrusting zoaria of *Pelmatopora* usually have a substantial calcified basal wall.

The zooecia of unattached uni- and bilaminar, or erect cylindrical zoaria always possess calcified basal walls, which may be smooth on their lower surface or have a number of scattered tubercles, as in *Pelmatopora somptingensis*.

Lateral, proximal and distal walls. These vary in height, and may slope as in *Lagynopora* and *Leptocheilopora.* Large distal and lateral communication pores may be visible in the distal and lateral walls respectively. These are often seen in *Lagynopora, Hexacanthopora* and *Leptocheilopora* and, less frequently, in other genera. The distal communication pores are usually large, single, circular or oval, and may be regarded as septula possibly covered originally by multiporous septular discs. Ordinary septula are usually smaller. The lateral communication pores are smaller than the distal, but more numerous. Often three or four occur in line along the middle of each lateral wall, or they may be scattered. Distal and lateral communication pores are visible in the specimen of *Leptocheilopora magna* shown in Pl. 6, fig. 5, and in the single zooecium of *Castanopora magnifica* shown in Text-fig. 112. In the latter their structure is unusual : small, circular pores occur in the centres of oval depressions surrounded by low rims. Communication pores have not been seen in *Pelmatopora* or other genera : septula are probably present but very minute. Dietellae occur in *Lagynopora, Hexacanthopora* and *Leptocheilopora*—they are visible, for example, in some zooecia of *Hexacanthopora sexspinosa* Lang (Pl. 5, fig. 3).

FRONTAL WALL. Lang (1921 : xxxvi) described the frontal wall of membranimorphs, from which the cribrimorphs are to be derived as a " composite structure. Laterally and proximally it is calcareous [the gymnocyst], and slopes outwards, gradually becoming continuous with the lateral walls ; but all the middle and distal parts [the aperture] are occupied by a flat and chitinous roof ".

In the Cretaceous cribrimorph Polyzoa a secondary frontal wall of costae (the frontal shield) is formed over the aperture, the only part of the primary frontal wall to be retained being the calcareous gymnocyst surrounding the aperture. The frontal membrane, or chitinous part of the primary frontal wall originally covering the aperture, is not preserved.

PRIMARY FRONTAL WALL—GYMNOCYST. The gymnocyst is apparently always smooth. It is variable in width, usually wider proximally to the frontal shield and narrower along its sides. Occasionally a narrow band of gymnocyst may extend round the distal and lateral margins of the orifice. The amount of visible gymnocyst is commonly diminished by close growth of the zooecia, and, in many genera of Cretaceous cribrimorphs, it is totally concealed by plentiful interzooecial secondary tissue. As in some species of *Pelmatopora,* the secondary tissue may contain lacunae through which the gymnocyst is locally exposed.

SECONDARY FRONTAL WALL—FRONTAL SHIELD. The general and detailed features of the frontal shield, and its relationship to other zooecial characters, provide one of the most useful means of differentiating genera and species in the Cretaceous cribrimorph Polyzoa. Essentially, the frontal shield is composed of a series of costae which develop from the edge of the gymnocyst and overarch the aperture, usually fusing along the mid-line of the zooecium. This basic arrangement is complicated in various ways. There is often considerable morphological similarity between the frontal shields of fossil cribrimorphs and those of some of the Recent Cribrilinidae. The development of some features in Recent species (observed by Harmer and others) shows that some analogous characters in the fossils may have had an origin different from that suggested by Lang.

Development and structure of the costae. Hincks has described the costae of Recent Cribrilinidae, such as those of *Membraniporella nitida* (Johnston) as " probably the equivalent of the marginal spines on many of the Membraniporae ". Harmer (1902 : 292) agreed with Hincks's opinion and again mentioned *M. nitida*, suggesting that the modified, overarched, marginal spines developed by closure of embayments in the edge of the gymnocyst (Text-fig. 2*A–C*). Harmer also figured (1902, pl. 15, fig. 7) the ancestrula and some young zooecia of a specimen of *Cribrilina radiata* (Moll), in which he suggested the costae are analogous to horizontal calcareous lobes occurring on the inner sides of the marginal spines of the Membraniporid ancestrula. Norman (1903 : 91) described a more complex development of costae and has shown that lumen pores and lateral costal fusions may be formed as the costae develop. Costae of four of the species described by Norman are shown in Text-fig. 2*D–G*, p. 27.

Other modes of formation of costae were recorded, somewhat obscurely, by Waters (1923 : 545–573). Apparently, in the Recent species *Lepralia otto-mulleriana* (Moll), a thin calcareous lamina may gradually spread from the proximal and lateral margins of the aperture leaving uncalcified pores. Subsequently, irregular ridges may separate costa-like areas, each with one pore (Text-fig. 3*C*). In other specimens, irregular " broad spines " with small " nodulations " may spread secondarily from the proximal and lateral margins of the aperture and so produce a costate appearance (Text-fig. 3*A*, *B*, p. 28).

Thus the costae of different and the same species of Recent Cheilostomata may differ structurally and have developed in different ways. In those forms in which the costae are produced by apparent closure of embayments in the edge of the gymnocyst the costae are continuous tubular structures. The costae of all the Cretaceous cribrimorph Polyzoa described here appear to be of this type. It is possible that a costa with the structure described by Norman (1903) is not a simple tubular outgrowth of the type described by Harmer. Lumen lines may have been present in the costae of some Cretaceous species, but fine detail of this kind is easily destroyed during fossilization.

Harmer has pointed out that Lang's studies confirmed palaeontologically the supposed Membraniporid origin of Recent forms with costate frontal shields. Lang (1921 : 6, text-fig. 5) compared Membraniporid and cribrimorph Polyzoan structures —his text-figure is reproduced here, somewhat modified, in Text-fig. 4*A*, *B*, p. 29.

Number of costae. In the Recent Cribrilinidae and in the Cretaceous cribrimorph Polyzoa the number of costae in any single zooecium of a given zoarium may vary considerably. Illies (1953) even suggested that zooecial dimensions provide a better basis for specific diagnosis in species of *Rhiniopora* [*Castanopora*]. The number of costae is often greater in longer zooecia and fewer in shorter ones. Similarly, adult zooecia usually possess more costae than young zooecia in the same zoarium, though in some species the reverse may be true. Taken alone, the number of costae is thus of little diagnostic value, but, provided a sufficient number of specimens is available, the general range in number of the costae may be used, in conjunction with all other available characters, to differentiate species. Lang's statement (1919*d* : 197) that the number of costae " tends to increase during development [in the Pelmatoporinae] but may be very much reduced catagenically " indicates the variability of this character.



Fig. 2. *A–C.* Diagram showing the development of continuous tubular costae from closure of bays in the edge of the gymnocyst.

D " *Cribrilina annulata* Fabricus "—based on Norman's figure—1903, pl. 8, fig. 10.

E. " *Cribrilina balzaci* (Audouin) "—based on Norman's figure—1903, pl. 9, fig. 6.

F. " *Gephyrotes nitido-punctata* Smitt "—based on Norman's figure—1903, pl. 8, fig. 13.

G. " *Cribrilina innominata* Couch "—based on Norman's figure—1903, pl. 9, fig. 3.

g, edge of gymnocyst ; *is*, intercostal space or " lacuna " ; *l.c.f*, lateral costal fusion ; *ll*, lumen line ; *l.p*, lumen pores or pelmata ; *m*, median line of frontal shield ; *p*, non-calcareous papillae.

Lumen pores : pelmata : pelmatidia. The lumen pores which occur along the mid-line of each costa in some of the Recent Cribrilinidae are closely analogous to the pores which occur in the same position on the costae of many Cretaceous cribrimorph Polyzoa. However, Lang (1921 : xli) termed the latter *pelmata* and *pelmatidia*, and described them as " hollow upturned prolongations, presumably continued in the living organism as free spines, but appearing in the fossil as a row of broken ends ". In some species, such as the more complex forms of *Pelmatopora*, he distinguished " primary, secondary, tertiary and quaternary " pelmata and pelmatidia corresponding in position to equivalent lateral costal fusions. The terms " primary, secondary " etc. were used with reference to the supposed order in which the pelmata developed and shifted outwards from the median area of the frontal shield. In agreement with Harmer (1926 : 471) I regard the pelmata and pelmatidia as lumen pores, which, as in the Recent Cribrilinidae, have developed during the formation of the costae bearing them. In Recent specimens, as a costa is formed, the lumen pores are apparently left as uncalcified, membranous areas on its upper surface. There is no indication that the pores are formed first at the inner end of the costa and subsequently migrate towards its outer end, an hypothesis with which Lang attempted to account for the position of pelmata and pelmatidia in most of the



FIG. 3. Development of costa-like structures in the Recent species *Lepralia ottomulleriana* Moll. *A, B, C,* based on parts of Water's figures, 1923, pl. 17, figs. 1, 2 and 5 respectively.

c.l, thin calcareous lamina ; *irn,* irregular ridges with " nodulations " ; *isn,* irregular broad " spines " with " nodulations " ; *n,* " nodulations" ; *p,* pores.

Cretaceous forms. Thus Lang stated (1922 : 6) " It is convenient to consider that the primary [first formed] pelma has shifted proximally on the costa, carrying with it a part of the median line of fusion, and that the secondary pelma has been added at the distal [inner] end of the costa. . . . This process is repeated in more ad-



FIG. 4. *A, B*. Diagram showing the development of a cribrimorph polyzoan (*B*) from a Membraniporid (*A*). Modified from Lang (1921, text-fig. 5). Chitinous parts stippled.

C–F. Types of costae, pelmata, pelmatidia and lateral costal fusions in Cretaceous cribrimorph Polyzoa.

C. Simple Pelmatoporid costae with long, single intercostal spaces and each with a large, oval pelma at the inner end. The cross-section shows the slight rim round the pelmata.

D. Costae of *Castanopora* with numerous equally small pelmatidia and a corresponding number of lateral costal fusions, each curved parallel to the edge of the gymnocyst producing rows of somewhat oblong intercostal spaces.

E. Costae of *Ubaghsia* with two very large pelmata, short lateral costal fusions and irregular intercostal spaces. The cross-section shows that the costae are wider than in other genera.

F. Costae of *Aeolopora* each with a single boss-like pelmatidium and no lateral costal fusions. The cross-section shows the extended rim of the pelmatidia.

vanced species". He summarized the supposed process in a series of diagrams (1919*d* : 198, text-figs. 2–5 ; 1922 : 6, text-fig. 2).

Pelmata and pelmatidia, or lumen pores, were recorded by Lang (1916*a* : 83, 84 ; 1922 : 1) in all the genera which he assigned to the Pelmatoporidae. In the material studied here they have also been found in *Lagynopora*, *Hexacanthopora*, *Leptocheilopora* and *Aeolopora*. There is no reason to suppose, as Lang has done (1922 : 8), that the pelmata in *Diacanthopora* are the broken ends of branched spines. In agreement with Harmer (1926 : 471), I regard them also as lumen pores. Lang (1921) has recorded minute pores in the costae of *Pliophloea* (p. 172), *Dishelopora* (p. 217) and *Histricopora* (p. 227). Although he did not call these minute pores "pelmata" or lumen pores, I regard them as such. It is very probable that minute pelmata, discernible only under high-power magnification, are, or were originally, present in most of the Cretaceous cribrimorph Polyzoa but have been lost during fossilization.

Although Lang stated (1922 : 5), "It is not possible to impose a very rigid line of demarcation between a pelma and a pelmatidium", he has used the supposed difference between the two to separate some subfamilies. Very often, as in the more complex species of *Pelmatopora*, a graded series occurs with the largest lumen pore (pelma) at the outer end, successive pores becoming gradually smaller towards the inner end of each costa. In the simpler species of *Pelmatopora* and in *Tricephalopora*, there is often a single, large, oval pelma near the inner end of each costa. In *Castanopora* a series of up to seven or eight equally small pores may occur, or, as in *Aeolopora*, a single, minute, boss-like structure with a central pore may be present near the inner end of each costa. The pelmata may have a distinct rim produced by local thickening of the costal wall round the pore. This feature is also found in Recent forms ; Waters (1923) has recorded such thickened pore edges produced during the development of the secondary frontal wall in *Lepralia otto-mulleriana* (Text-fig. 3*C*). In other genera of Cretaceous cribrimorph Polyzoa lumen pores may not occur in a continuous series and may lack distinct rims, as in *Leptocheilopora magna* Lang (Pl. 6, fig. 1). The different types of pelmata and pelmatidia which occur in the Cretaceous material described here are shown in Text-fig. 4*C–F*.

Lang's terms "pelmata" and "pelmatidia" are retained in the systematic descriptions, but only for convenience of reference to the size of the pores : pelmata are interpreted as large lumen pores, and pelmatidia as small lumen pores. I do not accept in any way Lang's hypothesis of their origin.

Lateral costal fusions. Lang's concept of a "proximal shift" of pelmata and lateral costal fusions presupposes that both features are secondary modifications of the costae. In some Recent forms, however, lateral costal fusions, like the pelmata or lumen pores, are developed during the primary growth of the costae (Text-fig. 2*D–G*) ; in others they may develop subsequently between established costae and divide an originally long, single intercostal slit into a series of small intercostal spaces. It is difficult to decide whether the lateral costal fusions which occur in some Cretaceous cribrimorph Polyzoa are formed, like the pelmata, during the initial growth of the costae or subsequently. Pairs of lateral costal fusions usually coincide with the pelmata on the costae. Norman (1903) suggested that in some

Recent forms adjacent costae produce lateral processes which meet half-way between the two costae and fuse (Text-fig. 2*E*). In other cases he suggested that adjacent, irregular costae may touch at various points along their length producing apparent lateral fusions, e.g., in *Membraniporella nitida* (Text-fig. 2*D*). In the Cretaceous material most of the costae are regular and the lateral fusions between them are distinct tubular structures, often curved parallel with the edge of the gymnocyst (Text-fig. 4*D*). In some species, such as *Ubaghsia ornata*, in which the pelmata may be very large, the costae expand at the levels of the pelmata and the lateral fusions are correspondingly short (Text-fig. 4*E*). Lateral costal fusions are not always present where pelmata are developed—*Lagynopora*, *Hexacanthopora* and *Leptocheilopora* have pelmata but no lateral costal fusions. Although the number of lateral costal fusions on each costa of the same frontal shield is usually constant, the larger costae may possess extra fusions.

Intercostal spaces. As in the Recent Cribrilinidae, the intercostal spaces of Cretaceous cribrimorph Polyzoa vary widely in shape and size. In many genera the spaces between adjacent costae are elongate, single openings (Text-fig. 4*C*). In others a series of lateral costal fusions subdivides the space between adjacent costae into a number of smaller intercostal spaces termed, by workers on Recent forms, " lacunae ". The term " lacunae " is used here, however, to describe openings in the interzooecial tissue. In some species with lateral costal fusions the outermost intercostal spaces may be fairly constant in shape, but little diagnostic value can be placed upon this feature. In general, the intercostal spaces of the earliest, simpler, Cretaceous cribrimorph Polyzoa are wider than the costae they separate, but in the majority of later forms the reverse is true. The intercostal spaces may be filled with secondary calcareous tissue which spreads outwards from the median area of the frontal shield : this is well shown by some species of *Lagynopora*.

Median area of fusion. The size and shape of this area may vary considerably in a given species between zooecia of different zoaria and even between zooecia of the same zoarium. Many median areas of fusion are very narrow, involving only the inner tips of the costae and almost confined to the mid-line of the frontal shield ; others are wider and fusiform producing a flattened top to the arched frontal shield. In some species of *Lagynopora* and of other genera, the costae may be firmly united for about two-thirds of their length by an outward extension of the median area of fusion.

Lang's concept of a proximal shift along the costae of the pelmata and lateral costal fusions led him, in his descriptions of some Pelmatoporid species, to use the term " median area of fusion " to include all the frontal shield inside the outermost lateral costal fusions, which he regarded as parts of the original median area of fusion. Thus he described (1922 : 301) *Pelmatopora somptingensis* as having " two paired rows of perforations [intercostal spaces] in the original median area of fusion (now comprising the greater part of the intraterminal front-wall [frontal shield]) ". The median area of fusion of *P. somptingensis* is, in fact, very narrow (Text-fig. 96). The same remarks apply to other species with lateral costal fusions : these structures are not part of the median area of fusion. Different types of median areas of fusion, as defined here, are shown in Text-fig. 5*a–d*.

Apertural bars. Lang (1921 : 20) has shown that the apertural bar of the less complex, earlier genera of Cretaceous cribrimorph Polyzoa is asymmetrical but that in later more complex forms it is symmetrical and composed of two costae firmly fused in the mid-line. There is often considerable variation in its form between different genera and species, and occasionally between specimens of the same species. The apertural bar may be straight and relatively featureless, as in *Andriopora minor* sp. nov., or distinctly " V " shaped, as in *Leptocheilopora filliozati*. In *Hexacanthopora* *sexspinosa* it is curved with a prominent median process. In *Lagynopora* the median process of the apertural bar is more pronounced and usually bifurcated, producing two hollow spine-like processes which curve outwards from the mid-line of the orifice and fuse with the proximal-lateral oral spines forming a " Y "-shaped proximal oral shield.



FIG. 5. Types of median area of fusion in Cretaceous cribrimorph Polyzoa.
a, Narrow median area of fusion found in less complex species of *Pelmatopora* ; *b*, narrow median area of fusion typical of frontal shields with numerous lateral costal fusions—found in the more complex species of *Pelmatopora* ; *c*, *d*, Langynoporid frontal shields, (*c*) narrow median area of fusion, (*d*) very wide median area of fusion. Median area of fusion shaded with horizontal lines.

The costae composing the apertural bar may also possess pelmata or pelmatidia if these are present in the costae of the frontal shield. Secondary calcareous tissue may accumulate on the apertural bar, as in some genera of the Tricephaloporinae, spreading proximally as a tongue of tissue covering the median area of the frontal shield. The whole apertural bar is easily modified by wear, median processes being particularly easily destroyed. Diagrams of different types of apertural bars developed in the material studied here are shown in Text-fig. 6*a*–*q*.

TERTIARY FRONTAL WALL. Continued upgrowth of interzooecial secondary tissue (p. 40), combined with lengthening and expansion of the tissue of tubular secondary orifices (p. 35), and parts of other secondary structures, produces a prominent, irregular tertiary frontal wall above the general level of the frontal shields. This composite type of tertiary frontal wall is well developed in some species of *Ubaghsia* (Pls. 21–23).

A less complex tertiary frontal wall is developed in some species of *Tricephalopora*. In some a tongue of secondary tissue, spreading from the apertural bar, forms part



FIG. 6. Types of apertural bars found in Cretaceous cribrimorph Polyzoa.

a–d, Apertural bars of *Pelmatopora*. (*a*) *P. solearis*, (*b*) *P. d'orbignyi*, (*c*) *P. somptingensis*, (*d*) *P. palmata*. *e–g*, Apertural bars of *Castanopora*. (*e*) *C. magnifica*, (f) *C. castanea*, (*g*) *C. jurassica*. *h*, Apertural bar of *Ubaghsia ornata* with four large pelmata. *i*, *j*, Apertural bars of (*i*) *Andriopora major*, (*j*) *A. minor*. *k*, *l*, Apertural bars of (*k*) *Aeolopora distincta*, (*l*) *A. nebulosa*. *m*, *n*, Apertural bars of (*m*) *Leptocheilopora filliozati* showing the median ridge which is continued along the mid-line of the frontal shield and (*n*) *L. vulnerata* in which there is a prominent median facet on the distal side of the "V"-shaped apertural bar. *o*, Apertural bar of *Hexacanthopora sexspinosa* with prominent median process. *p*, *q*, Apertural bars of two species of *Langynopora* showing the bifurcated median processes and their fusion with the proximal-lateral oral spines, (*p*) *L. pustulosa*, (*q*) *L. ampulla*. For convenient comparison each orifice and apertural bar is drawn at approximately the same scale, irrespective of actual size.

of a tertiary frontal wall, in others this structure is combined with a general upgrowth of tubular secondary orifices and interzooecial secondary tissue to form a more complete tertiary frontal wall, which is less raised above the general level of the frontal shields than in *Ubaghsia*. Lang (1921 : xlix) has observed that " *Hippiopora* appears to be the only described Cretaceous Cribrimorph outside the Pelmatoporidae which possesses a tertiary front-wall ".

ORIFICES. The primary orifices of young zooecia vary in shape but are sharply defined, and, apart from being larger, may remain still sharply defined and otherwise unmodified in adult zooecia. Accumulation of secondary calcareous tissue round a primary orifice, or other modications, produces a secondary orifice.



FIG. 7. Evolution and terminology of orifices suggested by Lang. Diagram based on Lang (1921, text-fig. 10).

PRIMARY ORIFICES. Lang (1921 : xlvi) suggested that the primary orifice in Cretaceous cribrimorph Polyzoa, although varying in shape, tended to pass from a " primitive " so-called " sub-semicircular " or " semicircular " form to more complex shapes. This evolutionary sequence is summarized in Text-fig. 7 ; the terms used are those devised by Lang to describe ten different shapes which he recognized. He was well aware of the uncertainty of suggesting such a sequence, for he stated, " This evolution must not, however, be strained. It is easy to see how a normal aperture [orifice] may become distinctly pliophloean without passing through a cribriline stage ".

The present work shows how, in most genera, and often within a given species, the shape of the orifice can be very variable. This feature is discussed and illustrated in some detail under the remarks on *Leptocheilopora filliozati* (p. 115), of which many specimens are available. Similar remarks apply to other species of *Leptocheilopora* and to species of other genera. *Pelmatopora* often has a comparable range of variation in shape of the primary orifice. Where proximal-lateral constrictions occur in an orifice they may be more or less pronounced, and the shape of the proximal margin may vary according to the form of the apertural bar. Conversely, but in only a few species, the orifice may have a fairly constant shape, as in *Leptocheilopora magna* Lang.

Thus the shape of the primary orifice is not usually of diagnostic value. It may vary in shape naturally or its form may be changed by wear ; probably the median proximal notch found in some orifices of *Leptocheilopora tenuilabrosa* (p. 107) is due to wear. No general evolutionary sequence in the shape of the primary orifice could be distinguished in the material studied here.

In contrast to the variable shapes which may occur, the length and breadth of the orifice in a given species are often constant within a very small range. Lang (1919*d* : 198) only briefly considered the size of the orifice in relation to that of the frontal shield, and suggested " This tends, as a rule, to increase ; but, in some cases . . . becomes smaller catagenetically ". In the genera studied here it is difficult to find evidence to support this view. It is apparent, however, that in a given species the length (*hr*) and the width (*lr*) remain fairly constant although the shape may vary. This feature is again well shown by *Leptocheilopora filliozati* (p. 115) in which the dimensions of the orifice, irrespective of its shape, are fairly constant, even in abnormal zooecia.

Lang has suggested (1919*d* : 197) that the shape of the orifice " tends to alter from being rather longer than wide to rather wider than long ", or, in other words, that *lr* generally exceeds *hr* in later genera and species. In those genera revised here in which the form of the primary orifice is not secondarily modified in the adult stages, I can find no indication of this tendency. It is apparent that species with orifices which are wider than high predominate throughout.

SECONDARY ORIFICES. In many genera the primary orifice is modified in adult zooecia by the marginal accretion of secondary calcareous tissue. A slightly thickened oral rim may be developed as in *Pelmatopora*, or increased accumulation may produce a tubular secondary orifice, as in some species of *Tricephalopora*. Other structures, such as oral spines and avicularia, may be incorporated in the general upgrowth of calcium-carbonate round the orifice, as in *Ubaghsia*, or distinct distal and proximal oral shields may be produced, as in *Lagynopora*.

Where a slightly thickened oral rim is developed the secondary tissue is usually accumulated fairly evenly round the distal and lateral margins. The proximal margin of the orifice, formed by the apertural bar, is seldom affected. Thus, although the dimensions of the orifice are slightly decreased they usually remain the same in proportion to one another and may be used, in conjunction with all other available characters, in specific diagnosis.

In genera with a tubular upgrowth of calcareous tissue round the primary orifice the secondary orifice is often very different in shape and size, and is separated from the primary orifice by a peristome of variable length. The formation of distal and proximal oral shields, incorporating other structures, is considered later.

ORAL SPINES. In some Recent cheilostome Polyzoa the shape and size of the oral spines varies widely even in a single species. Most oral spines are hollow in the centre, but they may be long, unbranched and rod-like, or bifurcated, or variously branched.

Frequently only the proximal-lateral pair of oral spines is branched, the remainder being unmodified. The oral spines are branched, for example, in some species of *Chaperia*, *Scrupocellaria* and *Amphiblestrum* (Text-fig. 9). The spines may be jointed near the base so that the upper part is easily removed by wear, leaving a small, hollow spine-base (Text-figs. 8*c*; 9*k*–*n*).

In most fossil specimens, including many of the Cretaceous cribrimorphs, only the bases of the oral spines are preserved. They may have been jointed near the base and have broken off at this point, or, as is more probable, they may have been unjointed and worn down to the base. Thus the apparent simplicity and similarity of the oral spines in many genera of Cretaceous cribrimorphs may be largely due to wear.



FIG. 8. Oral spines in Recent species.

a, *b*, *Cribrilina alcicornis* Jullien (1883, pl. 14, figs. 24, 25), (*a*) expanded, lobed oral spines preserved, (*b*) a worn orifice, the oral spines reduced to hollow spine-bases, and two worn avicularia. *c*, *d*, *Chaperia* "*quadrispinosa*" [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 549*c*], (*c*) distal end of worn zooecium with four hollow oral spine-bases and distal-median avicularium, (*d*) distal end of unworn zooecium with four complete oral spines and distal-median avicularium.

However, bifurcated and branched oral spines are preserved in some genera: *Ubaghsia* may possess branched spines, and some species of *Pelmatopora* have distinctive bifurcated oral spines. Jullien (1886, pls. 17–19) figured branched spines in *Ubaghsia* [*Steginopora*] *meudonensis*, *U.* [*S.*] *ocellata*, *U.* [*S.*] *demorgani* and in *U. reticulata* (see Pl. 22). All these species possess a substantial tertiary frontal wall which evidently protected the oral spines from wear. The variously modified oral spines which occur in some species of *Pelmatopora* were interpreted by Lang (1916*a*: 102; 1919*d*: 200; 1921: xxxiii–xxxiv; 1922: 247) as secondary avicularia which have replaced the oral spines. I have examined a very large number of specimens of *Pelmatopora*, including those seen by Lang, and have found nothing to suggest



FIG. 9. Types of oral spines in three genera of Recent cheilostomatous Polyzoa. *a–d*, "*Scrupocellaria frondis*" [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 31.12.19.7B]. Branched oral spines from four zooecia of the same zoarium. *e–g*, "*Scrupocellaria securifera*" [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 28.3.6.170]. Small oral spines tending to bifurcate. *h*, "*Amphiblestrum cervicorne*" [B.M. (N.H.) Zool Dept. Polyzoa Coll. slide 87.12.9.328]. Front view of paired, branched lateral oral spines and two unbranched distal oral spines. *i–j*, "*Chaperia* sp." [B.M. (N.H.) Zool Dept. Polyzoa Coll. slide TN 863A]. Front and side views of paired, bifurcated lateral oral spines and four unbranched distal oral spines. *k–m*, "*Chaperia cervicornis*". (*k*) [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 99.5.1.548]—stout, bifurcated oral spine, jointed at the base and with the central hollow visible; (*l*) [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 444D]—branched oral spine, jointed at the base and with large central hollow; (*m*) [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 99.7.1.1202]—bifurcated oral spine, jointed at the base. *n*, "*Chaperia galeata*" [B.M. (N.H.) Zool. Dept. Polyzoa Coll. slide 53B]. Unbranched oral spine, jointed at the base and with central hollow.

that the structures are avicularian. Their bifurcated form, as in *Pelmatopora gregoryi*, is a condition which is unknown in avicularia, and they lack true avicularian characters. They occur consistently as part of the symmetrically arranged series of spines round the orifice, and they are most closely comparable with the expanded and bifurcated oral spines which occur in some Recent cheilostome Polyzoa.

Lang (1916*a* ; 1919) has referred to the structures as " avicularia " and later (1921) as a " kind of (supposed) aviculoecia [avicularia] ". In his fullest description of *Pelmatopora* (1922) he was evidently undecided as to their nature. In a discussion of " the elaboration of the aperture [orifice] " he has stated (p. 244) " . . . what appears to be a new structure [in *Pelmatopora*] occurred on the distal edge of the secondary apertural [oral] rim, corresponding in position to the obliterated distal pair of apertural [oral] spines, though probably lying just outside them ; it is even possible that the apparently new structure was a new development of the apertural [oral] spines, as the evidence of specimen D.23397 of *P. brydonei* is not altogether conclusive. . . . In this specimen the right distal apertural [oral] spine and the new structure on the same side are both present and seem to be separate. On the other hand, the astogenetic evidence of *Pelmatopora saltdeanensis* [= *P. gregoryi*] points to this supposed new structure being but a development of the distal pair of apertural [oral] spines. . . . It is probable that, if not developments of apertural [oral] spines, these projections are aviculoecia [avicularia], and that the pointed depressions on their distal surfaces are the rostra. During evolution they enlarge, lengthen, and become bifid . . . ". In his section on the systematics of *Pelmatopora* Lang referred to the structures as " secondary aviculoecia ".

The structure in D.23397 is discussed in detail under the remarks on *P. brydonei* (p. 166) and is figured (Text-fig. 78*a*, *b*). It is an enlargement of the unmodified oral spine, and regeneration of the whole orifice (see Lang, 1922 : 294) has not occurred. As Lang suggested in his reference to the "astogenetic evidence" of *P. saltdeanensis* [= *P. gregoryi*] and in his remarks on that species (1922 : 319), a comparison of oral spines of young and adult zooecia in a given zoarium points to his "secondary aviculoecia" as only a development of the unmodified distal oral spines. In the species of *Pelmatopora* in which the adult zooecia possess enlarged oral spines,[1] the young zooecia have unmodified spines which gradually become expanded or bifurcated as the zooecia mature.

In a given species of *Pelmatopora*, as in Recent cheilostome Polyzoa, the size and shape of the enlarged oral spines may vary considerably. This variation, which is produced largely by different degrees of wear, is shown particularly in Text-fig. 81*a–d* (*P. marsupitorum*), Text-fig. 88*a–f* (*P. gregoryi*), Text-fig. 97*a–g* (*P. somptingensis*), and in Text-fig. 101*a*, *b* (*P. palmata*). It follows that variations in shape of the enlarged oral spines are not always suitable characters on which to differentiate species.

The oral spines are always arranged symmetrically around the orifice, whatever their number. They are often equally spaced, but occasionally occur in paired groups of two or three. In the material studied, the adult zooecia in some genera have fewer oral spines than the young zooecia (e.g., some species of *Pelmatopora*). In other cases the number of oral spines is the same in adult and young zooecia. There is no indication that the later species tend to have fewer oral spines than the earlier ones. In the following Table, 15 species are listed in which it has been possible to compare the numbers of oral spines in young and adult zooecia. It can be seen that, in the four genera represented, the number of oral spines is more variable in the young stages, and that reduction in number, if any, is commonly to four oral spines in the adult zooecia.

---

[1] *Pelmatopora coryli*, *P. quadrivolucris*, *P. brydonei*, *P. marsupitorum*, *P. gregoryi*, *P. somptingensis* and *P. palmata*.

		Number of oral spines	
Horizon	Species	Young zooecia	Adult zooecia
M. cortestudinarium and *M. cor-anguinum* Zones	*Lagynopora lagena*	4, 5 or 6	4
M. cor-anguinum Zone	*L. birlingensis*	4	4
	L. praecursor	6	4
Marsupites Zone	*L. ollula*	6 or 7	4 or 5
O. pilula Zone	*L. urceolus*	5 or 6	4
	L. vasculum	4, 5 or 6	4
O. pilula and *B. mucronata* Zones	*L. amphora*	6	4
G. [*A.*] *quadrata* Zone	*L. ampulla*	5	4
	L. furcifera	5 or 6	4
Marsupites and *G.* [*A.*]*quadrata* Zones	*Hexacanthopora sexspinosa*	6 or 7	4, 5 or 6
O. pilula and *G.* [*A.*] *quadrata* Zones	*Leptocheilopora filliozati*	4, 5 or 6	4
B. mucronata Zone and *O. lunata* Chalk	*L. vulnerata*	7	7
B. mucronata Zone and Maastrichtian	*L. magna*	4	4
B. mucronata Zone and *O. lunata* Chalk	*Castanopora castanea*	4	4
	C. dibleyi	4 or 5	4 or 5

Lang (1921 : xliv) has stated that the oral spines " are primarily six in number " and that " In most families of Cretaceous Cribrimorphs the primary six apertural [oral] spines have been reduced to four ". He also mentioned a number of genera and species as not showing this tendency. From the exceptions mentioned by Lang, and from the evidence of the present work, it becomes difficult to recognize a general reduction in the number of the oral spines. The most that may be said is that some species of some genera show a variable reduction in the number of oral spines, others show no reduction, but none shows an increase.

ORAL SHIELDS. Accumulation of secondary calcareous tissue round the orifice may conceal the oral spines or they may form a visible part of a distal or proximal oral shield. Occasionally, as in the species of *Pelmatopora* with enlarged distal oral spines, the space between the spines may be filled with secondary tissue forming a small distal oral shield. More frequently, as in *Lagynopora*, the proximal-lateral pair of oral spines fuse with the bifurcated median process of the apertural bar thus forming a prominent " Y "-shaped proximal oral shield (Text-fig. 6*p*, *q*). In *Ubaghsia* the proximal-lateral oral spines may be enlarged, but otherwise unmodified, and fused on to the distal side of large lateral oral avicularia. Lang (1921 : xlv) mentioned other kinds of proximal and distal oral shields in which the oral spines may be involved, but these are not found in the genera described here. In genera with prominent hyperstomial ovicells the distal pair of oral spines often flank the entrance to the ovicell; in some cases they appear to be displaced by it (e.g., Text-fig. 67—*Pelmatopora fecampensis*). A projecting distal oral shield may be formed in species with endozooecial ovicells. In these the space between the distal oral spines may be filled with secondary calcareous tissue (e.g., Text-fig. 89—*P. gregoryi*).

INTERZOOECIAL SECONDARY TISSUE. Lang has pointed out (1922 : 4) that the absence or extreme scarcity of interzooecial secondary tissue may be a useful diagnostic character. Many of the genera he described (1921) are devoid of it, and in the present material, it is absent in *Lagynopora*, *Hexacanthopora*, *Leptocheilopora* and *Aeolopora*.

When interzooecial tissue is present it may be very variable in amount. In the Pelmatoporidae, for example, a few genera, such as *Castanopora*, lack it, but in others, such as *Pnictopora* (Lang, 1922 : 145) the characters of the zooecia are almost totally concealed by it. *Pelmatopora* provides a good example of the extreme variation in amounts of secondary tissue which may be present in different species of the same genus and in different specimens of the same species. Thus in *P. calceata* there is no interzooecial secondary tissue; in *P. crepidaria* it is only very slightly developed near the orifices; and in *P. solearis* it may occur in contour-like ridges between the zooecia or it may be abundant, filling the interzooecial valleys. Other species of *Pelmatopora* may have extremely variable amounts of interzooecial secondary tissue, but it is generally abundant, either filling the interzooecial valleys or standing up as irregular ridges above the general level of the frontal shields.

The amounts of interzooecial tissue vary so widely that it is not a character which is useful for specific differentiation. Thus, although Lang (1922 : 319) separated *P. cellium* [= *P. gregoryi*] from *P. saltdeanensis* [= *P. gregoryi*] solely on its supposedly greater accumulation in the former, the two species are alike in other characters and are here regarded as synonymous.

Apparently there is a tendency, seen in some young zoaria, for interzooecial tissue to be concentrated near the orifices in the early stages of its accumulation. In later stages, if it is at all abundant, it is fairly evenly distributed in the interzooecial valleys. Its continued accumulation, spreading out from the adult parts of a zoarium, may almost obscure the young zooecia. The ancestrula is often thus covered.

In the early stages of zoarial growth, before deposition of interzooecial tissue is complete, the gymnocysts of the zooecia may remain locally exposed. As accumulation continues, irregular openings or lacunae are maintained. These expose tissue which was deposited earlier or, occasionally, small areas of the gymnocyst may remain visible.

Lang (1921, text-figs. 7–9) has summarized three stages in the accumulation of interzooecial secondary tissue. It is difficult to be certain whether the tissue is first deposited in a narrow band round the gymnocyst as Lang has shown (text-fig. 7). In the material described here, the interzooecial tissue is apparently not deposited in a narrow ridge near the edge of the gymnocyst, but in a thin layer spreading from isolated points in the lowest parts of the interzooecial furrows. Subsequently, as shown by Lang (text-figs. 8, 9), lacunae remain in the accumulating tissue, but are gradually eliminated as it is built up into irregular interzooecial ridges.

General upgrowth of interzooecial tissue may contribute to the formation of a tertiary frontal wall (p. 33).

AVICULARIA. Workers on Recent and Tertiary cheilostome Polyzoa usually distinguish two main types of avicularia: (*a*) " adventitious " forms which occupy various positions on the outside walls of normal zooecia, and (*b*) " vicarious " or " interzooecial " forms which occur in series with normal zooecia. The terms " interzooecial " and " vicarious " are usually regarded as interchangeable.

In the Cretaceous cribrimorph Polyzoa, however, I distinguish between interzooecial and vicarious avicularia. The former are usually much smaller than the zooecia which they accompany and they do not occupy positions which would usually contain normal zooecia. The latter are often as large as, or larger than, the zooecia with which they occur, and occupy spaces which would otherwise be taken up by normal zooecia (Text-fig. 10*a*).

Adventitious avicularia are rare in the Cretaceous cribrimorphs. Lang (1921 : 166) has recorded distal adventitious avicularia in *Monoceratopora.* In the material described here, larger, apparently adventitious, distal avicularia occur in *Castanopora armata* sp. nov. (Text-fig. 103).

Vicarious avicularia are also infrequent in the Cretaceous cribrimorph Polyzoa. Some zoaria of *Leptocheilopora tenuilabrosa* (Pl. 7 ; Text-figs. 47, 48) possess very distinctive, large, spatulate, costate vicarious avicularia with distal apertural spines. Levinsen (1907 : 155, pl. opp. p. 160, figs. 1, 1*b*, 1*c*) has recorded large, spatulate, vicarious avicularia of a more usual type in *Castanopora* [*Cribrilina*] *labiata* (Text-fig. 10*b*).

Interzooecial avicularia occur very commonly in many genera of the Cretaceous cribrimorphs. They vary considerably in size, shape, number and position, and they may be directed distally, proximally, or variously. *Pelmatopora* affords a good example of the variation in detail which may occur in the interzooecial avicularia of one genus (Text-fig. 10*c–g*) and within a given species of one genus (Text-fig. 98*a–f—P. somptingensis*).

The shape of interzooecial avicularia is easily modified by wear. Sharply pointed rostra, which occur in various species of *Pelmatopora* and *Castanopora,* are often removed by wear leaving an apparently oval avicularian aperture. It may thus be misleading to record as oval the avicularia of species of which only one or very few specimens are available. This may be so in the case of *Pelmatopora calceata.* Similarly the transverse bar which may divide the rostrum of an avicularian aperture from the proximal portion is often broken, and it is difficult to decide whether the remaining lateral constrictions are the broken ends of such a transverse bar or true condyles on which the mandible was hinged (Text-fig. 10*h*). Interzooecial avicularia are often raised on fairly stout stalks, as in *Pelmatopora somptingensis* (Text-fig. 98*a, b*). In avicularia of this type, very little wear can remove the rostrum completely and reduce the visible structure to an oval cross-section of the stalk.

Where the avicularia are paired, one on either side of an orifice, they are often secondarily thickened and raised on very stout stalks which may fuse above the orifice, forming part of the proximal oral shield. This structure is well developed in *Ubaghsia ranunculus* (Text-fig. 117) and in other species of the same genus. Occasionally, as in *Haplocephalopora* (Lang, 1922 : 94), the avicularia may be very small and completely raised on the rim of a secondary tubular orifice.



FIG. 10. Types of avicularia in Cretaceous cribrimorph Polyzoa.
a, Diagram showing the occurrence of the three main types. *b*, Vicarious avicularium of *Castanopora labiata* (Levinsen). *c–g*, Interzooecial avicularia of *Pelmatopora*, (*c*) *P. gregoryi*, (*d*) *P. somptingensis*, (*e*) *P. calceata*, (*f*, *g*) *P. marsupitorum*. *h*, General terminology of avicularia.

In species with abundant interzooecial tissue the avicularia may be almost immersed, and it becomes difficult to decide whether they are interzooecial or adventitious. Judging from the avicularia of species which lack interzooecial tissue the great majority appear to be truly interzooecial rather than adventitious. Waters (1923) has observed that Lang's text-figures of Cretaceous cribrimorph Polyzoa (1921 ; 1922) give the impression that most avicularia grow out of parts of gymnocysts of the zooecia, but that in his plates Lang has not shown this relationship. There is no indication that the majority of avicularia are adventitious.

Waters (1923) has also pointed out that there is great variation in the avicularia of some Recent cheilostomatous Polyzoa, and that there are genera, and even species, in which they are sometimes present and sometimes not. It has already been mentioned that in the Cretaceous cribrimorphs there is considerable variation in the interzooecial avicularia, but provided only well-preserved material is used and that avicularian characters are considered in relation to all other available features, the avicularia are certainly of value for differentiating species.

In some species the avicularia are always distally pointed and directed and paired near the orifices, while in others they may have a noticeably sporadic distribution and be variously directed. Avicularia of two or more distinct sizes may be present, as in *Hexacanthopora sexspinosa*. In the material described here the general characters of the avicularia of a given genus are fairly constant although the detailed characters may vary. Thus, in *Leptocheilopora*, apart from the large vicarious avicularia of *L. tenuilabrosa*, avicularia are absent. Those of *Lagynopora* are commonly very large and oval with lateral constrictions. In *Castanopora* they are usually long and pointed with somewhat spatulate rostra and often with transverse bars; in *Pelmatopora* the unworn avicularia generally have pointed rostra and often lack transverse bars.

The " secondary aviculoecia [avicularia] ", which Lang described as occurring in the more complex species of *Pelmatopora*, are modified oral spines. The structures have been discussed already (p. 38).

In the interzooecial avicularia of the material described here no example has been found in which minute spines are present on the distal margin of the avicularian aperture.

Heterozooecia other than avicularia. These are apparently rare in the Cretaceous cribrimorph Polyzoa. In some species of *Pliophloea*, Lang (1921 : 174) has recorded interzooecial " swollen and hollow " tissue with small round pores and has suggested that the structures are " cenoecia " [kenozooecia]. In the present work, one specimen of *Aelopora distincta*, D.40658, was found to have clusters of very small, convex, smooth heterozooecia with large oval apertures on their upper surfaces (Pl. 1, fig. 5). These structures may be kenozooecia. Heterozooecia other than avicularia have not been found in the other genera and species studied.

Ovicells. I agree entirely with the remarks made by Brown (1952 : 36) in a general discussion on the ovicells of the Cheilostomata. The present study shows that his remarks are completely applicable to the ovicells of Cretaceous cribrimorph Polyzoa.

Brown stated " In the majority of the Cheilostomata the ovicells appear as globular distal chambers sometimes resting lightly on and sometimes completely immersed in and covered by the frontal wall of the distal zooecium. Gradations between these two forms can be traced and no distinct boundary can be defined between the first or *hyperstomial* type (Text-fig. 11*a*), and the second or *entozooecial* [endozooecial] type (Text-fig. 11*b*). In fact, Silén (1944) has shown that in the Anasca there is essentially no difference between these two forms so far as their structure is concerned, and I have no doubt that the same remarks will be found to apply to those ovicells of the Ascophora which are of the same type. . . . Thus it appears impossible to sustain the view of some workers that the difference between " hyperstomial " and " entozooecial " ovicells can be used to subdivide not only genera but also families. . . . However, the terms are undoubtedly useful for descriptive purposes . . . ".

The examples given by Brown of the occurrence of both types of ovicell in species of the same genus and of the use of the two forms for subdivision of families can be paralleled by examples from the Cretaceous cribimorph Polyzoa. In *Pelmatopora* the globular ovicells may be external, as in *P. fecampensis* (Text-figs. 66–68), or partly concealed by the wall of the distal zooecium, as in *P. quadrata* (Text-fig. 64), or completely concealed, producing externally only a distal transverse shelf over the orifice, as in *P. marsupitorum* (Text-fig. 82), *P. somptingensis* (Text-fig. 99*a*, *b*) and *P. gregoryi* (Text-fig. 89). Lang (1921 : 4–6) has used the two types of ovicells to separate into two major groups the eleven families which he recognized in the Cretaceous cribrimorphs. One group with four families, including his Pelmatoporidae, he diagnosed as having endozooecial ovicells; the remaining families he grouped together as having hyperstomial ovicells. Lang's statement (1921 : xxxv) that " the ovicell is of the same type throughout any given family of Cretaceous Cribrimorphs " is incorrect.



FIG. 11. Two types of ovicell in Cretaceous cribrimorph Polyzoa.
a, Hyperstomial. *b*, Endozooecial. *dz*, distal zooecium; *ec*, ectooecium; *ech*, brood chamber; *en*, endooecium; *mv*, median vesicle; *op*, operculum. (After Harmer, 1926, text-fig. 1, and Brown, 1952, text-fig. 5.)

Following general usage (e.g. Brown, 1952 : 37) the outer and inner layers which compose the wall of the ovicell are termed " ectooecium " and " endooecium " respectively—" endooecium " is used in the present work as an alternative to " entooecium ".

In the hyperstomial ovicells in the present material the endooecium is apparently always smooth and complete. The ectooecium may be complete, or marginal with a median-proximal fenestra which exposes part of the endooecium. The ectooecium is often smooth, but diagonal and median ridges may be present, or it may be costate as in *Leptocheilopora tenuilabrosa* (Text-figs. 46, 47), or corrugated as in some specimens of *Aeolopora distincta* (Text-fig. 16). The upper margin of the opening of the ovicell may have a distinct labellum bounded by lateral slits, as in *Aeolopora*, or it may be entire as in the majority of genera and species. The opening of the ovicell is frequently flanked by a pair of oral spines, but equally often these may be obscured by the growth of the ovicell. Occasionally the growth of hyperstomial ovicells may displace the oral spines—this has apparently happened in some specimens of *Pelmatopora fecampensis*.

The presence of endozooecial ovicells is often completely concealed by interzooecial secondary tissue. Endozooecial ovicells may, however, produce a distal oral shield by the development of a distal transverse shelf filling the space between the distal pair of oral spines.

Other types of ovicell have been distinguished by Levinsen (1909), Canu & Bassler (1920), Brown (1952) and by other authors. In the material studied here I have seen only hyperstomial and endozooecial types.

Young zooecia and ancestrulae. Only two generalizations seem to be true of the young zooecia of all the Cretaceous cribrimorph Polyzoa. They are always smaller than the adult zooecia, and in any given zoarium the young zooecia never possess fewer oral spines than the adult zooecia.

The general characters of young zooecia, compared with the equivalent characters of the adult zooecia in any given zoarium, may be summarized briefly as follows :

(1) Size and shape of zooecia	Always smaller, but equally variable in shape.
(2) Extent of gymnocyst	Often more widely exposed, due partly to the general absence of interzooecial secondary tissue.
(3) Frontal shield	Often apparently more arched, again partly due to the absence of interzooecial secondary tissue.
(4) Number of costae	Very commonly less, but occasionally the same or more.
(5) Number of pelmata and lateral costal fusions	Very commonly less, but may be the same.
(6) Median area of fusion	Often narrower, but may be relatively as wide.
(7) Apertural bar	Usually simpler, lacking secondary modifications such as median processes or extra tissue.
(8) Primary orifice	Always smaller and usually the same shape, but liable to exhibit the same degree of variation in relative *hr* : *lr* values.
(9) Number of oral spines	Often greater or the same, but never less.
(10) Interzooecial secondary tissue	Commonly absent or scanty.
(11) Avicularia	Usually smaller and sporadic, but otherwise similar.

The majority of these remarks could be applied to the ancestrula of any given zoarium when it is compared with the young or adult zooecia.

Measurements

In each species studied, measurements are given where possible for the ancestrula, young and adult non-ovicelled zooecia and avicularia of parts of zoaria where normal growth appears to have taken place. It has been pointed out already (p. 24) that zooecial size may vary so greatly, often within a single zoarium, that in the absence of more definite characters it should seldom be used as a basis for differentiating species. On the other hand, the length and breadth of the orifice in a given species are often constant within a very small range and when considered in relation to all other available characters are often of value for specific differentiation.

The margins of the primary orifice are well defined and it is possible to make accurate measurements. In contrast, measurement of the zooecia is often made difficult by the fact that the distal and proximal walls may slope, and by the accumulation of interzooecial secondary tissue which obscures the true zooecial boundaries.

The following abbreviations are used in the present work :

Lz = length of zooecium.
lz = width of zooecium.
hr = length of orifice.
lr = width of orifice.
Lc = length of caudae.
lc = width of caudae.

The first two abbreviations are also used for the length and width of avicularia. The last two are used in the description of *Andriopora major* sp. nov. (p. 268).

Terminology

Brown (1952 : 32) has already commented on the large number of new terms which Lang (1913–19 ; 1921 : xxvi–li) introduced in his studies on the Cretaceous cribrimorphs to describe the " skeleton as distinguished from the tissue which secretes it ". I agree that there is no need in practice for such a complete duplicate terminology, and accordingly follow the general usage of modern zoologists, although I retain a few of Lang's terms such as "apertural bar". See also Lagaaij (1952 : 12).

ADULT ZOOECIA. See Zooecia.

ANCESTRULA. The first zooecium of a zoarium, formed by the metamorphosis of the larva.

APERTURAL BAR. Transverse proximal margin of the orifice formed by the fusion of the most distal pair of costae—symmetrical in all but the earliest Cretaceous cribrimorph Polyzoa.

APERTURE. The opening occupied in life by the frontal membrane. Used loosely by earlier workers and by Lang in the sense of the term " orifice " as defined here.

AVICULARIA. Heterozooecia provided in life with a powerful musculature producing movement in a mandible. They may be (a) *adventitious*—occupying various positions on the external walls of normal zooecia ; (b) *vicarious*—as large as, or larger than, the zooecia with which they occur, and occurring in positions usually occupied by normal zooecia ; or (c) *interzooecial*—usually much smaller than the zooecia which they accompany, not occupying positions which would usually contain normal zooecia.

CAUDAE. Elongate, narrow, tubular, proximal part of the gymnocyst in some uniserial genera.

COMMUNICATION PORES. Pores perforating the walls separating two zooecia (see also *dietellae*). [Syn., *septulum* (pl., *septula*)—a single small pore. Some of the large, single communication pores which occur in some genera of Cretaceous cribrimorph Polyzoa may have contained *multiporous septular discs* or thin disc-like structures with numerous small pores.]

CONDYLES. The tooth-like pivots of the operculum or the mandible.

COSTAE. [Syn., *costulae*.] Modified marginal spines, overarching the frontal membrane and forming the frontal shield in cribrimorph Polyzoa.

CRYPTOCYST. A more or less horizontal calcareous lamina on the basal side of the frontal membrane developed from the edge of the gymnocyst, but not completely subdividing the body-cavity.

DIETELLAE. Small enclosed spaces at the base of the distal and distal-lateral walls of the zooecia of certain Cheilostomata, and connected with the zooecia by communication pores.

DISTAL. The portions of a structure on the side away from the origin of growth or from the ancestrula.

DORSAL. The side of a zooecium opposite that composed of the frontal shield and orifice, or the side of a zoarium opposite that bearing the apertures [syn., *back*, *reverse*].

ECTOOECIUM. Outer layer of the ovicell wall.

ENDOOECIUM. Inner layer of the ovicell wall [syn., *Entooecium*].

ENDOZOOECIAL. Type of ovicell which may be completely immersed in and covered by the frontal wall of the distally adjacent zooecium, while opening into the vestibular arch [syn., *Entozooecial*].

FENESTRA. Uncalcified area in the ectooecium of an ovicell through which the endooecium is exposed.

FRONTAL MEMBRANE. Uncalcified part of the primary frontal wall—a chitinous membrane covering the aperture in life.

FRONTAL SHIELD. The secondary frontal wall in cribrimorph Polyzoa, formed above the aperture by the over-arching and fused costae.

FRONTAL WALL. Chitinous or calcareous cover of the ventral surface of the zooecia. In the cribrimorph Polyzoa it may be (a) *primary*—consisting of the calcareous gymnocyst, and in life of the chitinous frontal membrane, (b) *secondary*—formed by the frontal shield, or (c) *tertiary*—formed by an upgrowth and spread of secondary calcareous tissue above the primary and secondary frontal walls.

GYMNOCYST. The part of the frontal wall which lies between the aperture and the upper edges of the vertical zooecial walls—the calcified part of the primary frontal wall : it is usually most developed on the proximal side of the aperture.

HETEROZOOECIA. Modified zooecia either (*a*) lacking a polypide, or with only a vestige of one, but provided with a musculature, as in avicularia, or (*b*) without polypide or a musculature, as in kenozooecia.

HYPERSTOMIAL. Type of ovicell which rests on or indents the frontal wall of the distally adjacent zooecium.

INTERCOSTAL SPACES. Openings between the costae—they may be (*a*) single, long, slit-like spaces, or (*b*) subdivided into a series of smaller openings by lateral costal fusions. [Syn., *lacunae* of workers on Recent Polyzoa.]

INTERZOOECIAL. Term applied to any structure developed between zooecia of adjacent series, hence " interzooecial secondary tissue " is calcareous tissue accumulated between zooecia in the interzooecial valleys.

KENOZOOECIA. See Heterozooecia.

LABELLUM. The free tongue of the descending lamina of the ovicell.

LACUNAE. Irregular openings in the interzooecial secondary tissue.

LAMINA PERISTOMICA. Term introduced by Jullien and used by Lang, and here as synonymous with " tertiary frontal wall "—see Frontal Wall.

LATERAL COSTAL FUSIONS. Hollow cylindrical, lateral fusions between adjacent costae, usually arising at the levels of lumen pores. [Lang referred to the outermost lateral costal fusions as " primary ", and to the successive inner ones as " secondary, tertiary ", etc.]

LATERAL SLITS. The pair of slits at the sides of the labellum in some ovicells.

LUMEN LINE. Clear, membranous, median line seen along the upper surface of the costae in some Recent Cribrilinidae.

LUMEN PORES. Uncalcified, small, round, membranous areas on the upper surface of costae in some Recent Cribrilinidae. Lang's *pelmata* and *pelmatidia* of Cretaceous cribrimorph Polyzoa are lumen pores.

MANDIBLE. The triangular or rounded chitinous or calcareous structure which closes the orifice in the avicularium. Not preserved in the material studied here.

MARGINAL SPINES. Spines, other than oral spines, which surround the aperture. Modified as costae in the Cretaceous cribrimorph Polyzoa.

MEASUREMENTS. See p. 45.

MEDIAN AREA OF FUSION. The imperforate area formed by the fusion of the inner ends of the costae, which extends laterally from the mid-line of the frontal shield and which is bounded by the innermost intercostal spaces.

MEDIAN PROCESS. Median distal prolongation of the apertural bar. It may be a single, more or less pointed process, or it may bifurcate and fuse with the proximal pair of oral spines.

MULTIPOROUS SEPTULAR DISCS. See Communication Pores.

OPERCULUM. Chitinous or calcareous lamina, generally semicircular in shape, closing the orifice of the zooecium.

OPESIA. The opening into the body cavity which remains after development of the cryptocyst. In the Cretaceous cribrimorph Polyzoa it usually coincides with the aperture.

ORAL AVICULARIA. Avicularia which are associated with the orifice. In the Cretaceous cribrimorph Polyzoa a pair of avicularia, termed " lateral oral avicularia ", may occur one on either side of the orifice.

ORAL SHELF. The flattened rim round the distal and lateral margins of the orifice on which the operculum or mandible rests.

ORAL SHIELDS. Structures, other than a peristome, which develop round the primary orifice. They may form a *distal oral shield*, usually a marginal upgrowth produced by infilling of the space between distal oral spines with secondary tissue ; or a *proximal oral shield*, often formed by the fusion of the median process of the apertural bar with the proximal pair of oral spines, or by the thickening, upgrowth and fusion of lateral oral avicularia.

ORIFICE. The opening of the zooecium, covered by the operculum, for the extrusion of the polypide. It may be either (a) *primary* (the opening of the tentacular sheath covered by the operculum) leaving in the fossil state a well defined orifice with sharp limits and no marginal accumulation of secondary tissue, or (*b*) *secondary* when the primary orifice becomes surrounded by a peristome with a secondary orifice at its upper end. Lang introduced a series of terms such as "cribriline, normal, pliophloean" to describe various shapes of primary orifice (Text-fig. 7).

OVICELL. The distal, globular, structure in the fertile cheilostomatous zooecia, which serves as a brood-chamber to the developing larvae.

PELMA (pl., *pelmata*) and PELMATIDIUM (pl., *pelmatidia*). Large and small lumen pores respectively in the costae of Cretaceous cribrimorph Polyzoa. [Lang referred to the outermost pelmata and pelmatidia as "primary", and successively to the inner ones as "secondary, tertiary" etc.] I have retained these two terms merely for convenience of reference but with no implication of acceptance of Lang's hypothesis of their origin (see p. 28).

PERISTOME. Tubular upgrowth of the rim of the primary orifice with the secondary orifice at its upper end.

POLYPIDE. The polyzoan animal with its tentacles placed in the interior of the zooecium.

PROXIMAL. The parts of a structure on the side near to the origin of growth or ancestrula.

ROSTRUM. That part of an avicularium which is distal to the transverse bar or condyles and on which the mandible rests.

SEPTULUM. See Communication Pores.

TRANSVERSE BAR. Narrow calcareous bar in some avicularia, on which the mandible hinges.

YOUNG ZOOECIA. See Zooecia.

ZOARIUM. The entire polyzoan colony.

ZOOECIUM. An individual member of the polyzoan colony. There is a difficulty in applying the terms "young" and "adult" to zooecia. They are here used as follows :

Young zooecia are those formed in the early stages of growth of the zoarium, after the ancestrula.

Adult zooecia are those formed in the late stages of growth of the zoarium.

I reject the following terms used by Lang, and indicate for each, after the colon, the term used by me :

Ancestroecium : ancestrula.

Apertural spines : oral spines. [Also used correctly by Lang for the spines which may occur on the distal edge of the aperture in some avicularia.]

Asty : zoarium.

Aviculoecium : avicularium.

Cenoecia : kenozooecia.

Costal spines : marginal spines.

Cribriline : a term used by Lang to describe a particular shape of primary orifice (see Orifice).

Epheboecium : adult zooecium.

Extraterminal front-wall : gymnocyst.

Fenestra : Lang used this term for openings in the calcareous tertiary frontal wall developed in some Cretaceous cribrimorph Polyzoa.

Gerontoecium : adult zooecium.

Heteroecia : heterozooecia.

Interoecial : interzooecial.

Intraterminal front-wall : frontal shield.

Lamina : cryptocyst.

Median lacunae : lacunae.

Neanoecia : young zooecia.

Normal : a term used by Lang to describe a particular shape of primary orifice (see Orifice).

Oecium : zooecium.

Orthoecium : zooecium.

Pliophloean : a term used by Lang to describe a particular shape or primary orifice (see Orifice).

Primary aperture : primary orifice.

Termen : a term used by Lang for the edge of the gymnocyst surrounding the aperture.

Note on the General Arrangement of the Systematics

The scheme adopted in the systematic sections of the present work is as follows.

Families and subfamilies. These are redefined where necessary. Synonymies precede the diagnoses and any changes suggested are discussed under remarks on the families or subfamilies involved.

Genera. The synonymy of each genus is given first, then the type species is cited and the genus diagnosed. Any changes are discussed under the remarks, and the stratigraphical range of the genus is given.

Species. After the synonymy the type of each species is cited in detail and its status given. When no type has been previously selected lectotypes are chosen in order to establish the species firmly. The diagnosis which follows is based on structures preserved in fossilization only, and the succeeding description is based on the characters of all material available to me. Where species are known from the literature only, or from photographs only, this is stated, and when the systematic position of a species is uncertain on this account, the species is listed as such and is not included in the general systematic assessment. Measurements are given where possible, and then general remarks on each species.

The stratigraphical distribution is stated and all the specimens which have been examined are listed with full details of locality and horizon. Under the heading " Specimens " is listed all the appropriate material which is in the collections of the

Department of Palaeontology of the British Museum (Natural History), each specimen with its registration number prefaced by the letter " D ". Material in the collections of other institutions is listed under " Other Material ". This includes specimens from the R. M. Brydone Collection in the Sedgwick Museum, Cambridge, the registration numbers of which are prefaced by the letters " S.M., B.". Specimens from the collections in the Norwich Castle Museum are denoted by the letters " N.C.M.".

Unless otherwise stated all the specimens are from C. T. A. Gaster Collection.

VII. SYSTEMATIC DESCRIPTIONS

Phylum *POLYZOA*
Class ECTOPROCTA
Tribe GYMNOLAEMATA
Order CHEILOSTOMATA Busk, 1852
Sub-Order ANASCA Levinsen, 1909

Division CRIBRIMORPHA Harmer, 1926

Family Pelmatoporidae Lang, 1916*a*

1916*a*	Pelmatoporidae Lang, p. 83.	
1921	Pelmatoporidae Lang :	Lang, pp. 1, 6.
1922	Pelmatoporidae Lang :	Lang, pp. 1–394.
1923	Pelmatoporidae Lang :	Waters, pp. 547, 553.
1925*b*	Pelmatoporidae Lang :	Voigt, pp. 97, 98, 99.
1927	Pelmatoporidae Lang :	Canu & Bassler, p. 18.
1930	Pelmatoporidae Lang :	Voigt, pp. 537, 538.
1935	Pelmatoporidae Lang :	Bassler, p. 31.
1953	Pelmatoporidae Lang :	Bassler, pp. G23, G190.
1958	Pelmatoporidae Lang :	Didon, p. 82.

Diagnosis. Multiserial Cribrimorpha with one or more pelmata or pelmatidia on each costa.

Remarks. Lang (1921 : 1) has stated that " If ovicells are present in the Pelmatoporidae, they are endozooecial ". In his key to the families of Cretaceous cribrimorph Polyzoa (1921 : 4), he divided into two groups the eleven families he recognized : group A was diagnosed as having hyperstomial ovicells and no pelmata or pelmatidia, group B as having endozooecial ovicells with pelmata or pelmatidia generally present.

It is shown below that some genera of Lang's group A possess pelmata or pelmatidia, and that both hyperstomial and endozooecial ovicells occur in the Pelmatoporidae. Brown (1952 : 43), in a brief discussion of the Membraniporidae, has stated that " although other workers (Canu & Bassler, 1920 ; Osburn, 1940) have used the difference between hyperstomial and entozooecial [endozooecial] ovicells to distinguish families, this is a very unreliable criterion, especially in the light of Silén's researches (1944) into the nature of the reproductive processes of the Polyzoa " (see also p. 43). It is apparent that the same remarks apply to the Cretaceous cribrimorph Polyzoa. Ovicells are not used here to distinguish families.

STRATIGRAPHICAL DISTRIBUTION. Turonian to Danian.

Subfamily PELMATOPORINAE Lang, 1916*a*

1916*a* Castanoporinae Lang, pp. 83, 93.
1916*a* Pelmatoporinae Lang, pp. 84, 101.
1916*a* Andrioporinae Lang, p. 382 [*partim*—*Aeolopora* only].
1916*a* Lagynoporidae Lang, p. 393.
1916*a* Lagynoporinae Lang, p. 393.
1916*a* Leptocheiloporinae Lang, pp. 394, 396.
1919*d* Pelmatoporinae Lang: Lang, pp. 191–228.
1921 Castanoporinae Lang: Lang, pp. xxxv, xli, xliv, xlv, xlvii, l, li, and p. 1.
1921 Pelmatoporinae Lang: Lang, pp. xli, l, li.
1921 Lagynoporidae Lang: Lang, pp. 5, 42.
1921 Lagynoporinae Lang: Lang, pp. 5, 44.
1921 Leptocheiloporinae Lang: Lang, pp. 5, 68.
1921 Andrioporinae Lang: Lang, p. 86 [*partim*—*Aeolopora* only].
1922 Castanoporinae Lang: Lang, pp. 5, 7, 10, 15, 154.
1922 Pelmatoporinae Lang: Lang, p. 235.
1927 Andrioporidae Lang: Canu & Bassler, p. 17 [*partim*—*Aeolopora* only].
1927 Lagynoporidae Lang: Canu & Bassler, p. 17.
1930 Lagynoporidae Lang: Voigt, p. 495.
1935 Andrioporidae Lang: Bassler, p. 30 [*partim*—*Aeolopora* only].
1935 Lagynoporidae Lang: Bassler, p. 30.
1943 Lagynoporidae Lang: Gillard, p. 186.
1953 Lagynoporidae Lang: Bassler, p. G188.
1953 Andrioporidae Lang: Bassler, p. G189 [*partim*—*Aeolopora* only].
1953 Pelmatoporinae Lang: Bassler, p. G192.
1953 Castanoporinae Lang: Bassler, p. G192.

DIAGNOSIS. Pelmatoporidae with small or large zooecia; with unmodified, or, enlarged or branched oral spines; with or without oral shields; apertural bar with or without a median process; pelmata or pelmatidia present, either with one boss-like pelmatidium, or with one or more pelmata or pelmatidia on each costa; with or without lateral costal fusions; with or without a tertiary front wall; with or without avicularia (if present these may be either sporadic, small and rounded, or paired, large and distally pointed); ovicells endozooecial or hyperstomial.

REMARKS. The genera of Cretaceous cribrimorph Polyzoa which are revised here were placed by Lang (1916*a*; 1921; 1922) in five subfamilies. This arrangement may be summarized as follows:

Revised genera	Subfamilies of Lang
Castanopora *Rhiniopora* (= *Castanopora*) *Ubaghsia*	Placed in the Castanoporinae [Lang, 1922 : 154].
Pelmatopora *Batrachopora* (= *Ubaghsia*)	Placed in the Pelmatoporinae [Lang, 1922 : 235].
Lagynopora *Prodromopora* (= *Lagynopora*) *Hexacanthopora*	Placed in the Lagynoporinae[1] [Lang, 1921 : 44].
Leptocheilopora . . .	Placed in the Leptocheiloporinae[1] [Lang, 1921 : 68].
Aeolopora	Placed in the Andrioporinae [Lang, 1921 : 89].

---

[1] Lang (1921 : 5, 44) placed these two subfamilies in the Lagynoporidae.

No distinction can be made between the Castanoporinae and the Pelmatoporinae. In his key to the subfamilies of the Pelmatoporidae Lang (1922 : 17) diagnosed them as follows : (*a*) Castanoporinae are Pelmatoporidae in which " costae bear *pelmatidia*[1] of several orders with lateral costal fusions corresponding in number and position to the pelmatidia ", (*b*) Pelmatoporinae are Pelmatoporidae in which " costae bear each one or more *pelmata*,[2] and lateral costal fusions correspond in number and position to the pelmata ".

As it is not possible to distinguish clearly between pelmata and pelmatidia (see p. 30) the two subfamilies are not separable on the basis given by Lang, nor are there any other characters which clearly help to maintain the division.

All genera previously assigned by Lang to the Lagynoporinae, and *Leptocheilopora*, the only genus which he placed in the Leptocheiloporinae, have distinct pelmata or pelmatidia and thus belong to the family Pelmatoporidae, and, according to Lang's definitions of its subfamilies, can be placed only in the Pelmatoporinae. Thus the subfamilies Lagynoporinae and Leptocheiloporinae, and the family Lagynoporidae which comprises them, are all synonymous with the subfamily Pelmatoporinae.

Aeolopora possesses boss-like pelmatidia and consequently belongs to the family Pelmatoporidae and is excluded from the Andrioporinae which Lang (1921 : 86) diagnosed as lacking, on the costae, either " a median row of pores " or " a median slit ". Again, according to Lang's definitions of the subfamilies of the Pelmatoporidae, *Aeolopora* can be assigned only to the subfamily Pelmatoporinae.

Thus all the genera listed in the above table are here regarded as belonging to the Pelmatoporinae.

STRATIGRAPHICAL DISTRIBUTION. Turonian to Danian.

KEY TO THE GENERA OF THE PELMATOPORINAE REVISED HERE

(I) Pelmatoporinae lacking a tertiary front wall ; secondary orifice absent or only slightly developed.
  (A) Zooecia small ; a single boss-like pelmatidium at the inner end of each costa 1. ***AEOLOPORA*** Lang
  (B) Zooecia large ; one or more normal pelmata or pelmatidia on each costa.
    (1) Proximal oral shield well developed—formed by the bifurcation of the median process of the apertural bar and its fusion with the proximal pair of oral spines . . . . . . 2. ***LAGYNOPORA*** Lang
    (2) Proximal oral shield absent.
      (*a*) Median process of the apertural bar present and not bifurcated 3. ***HEXACANTHOPORA*** Lang
      (*b*) Median process of the apertural bar absent. 4. ***LEPTOCHEILOPORA*** Lang
        (i) Costae less numerous with fewer pelmata or pelmatidia and lateral costal fusions ; avicularia usually of one kind, either small, rounded and sporadic, where no distal oral shield is developed, or large, elongate, distally pointed and directed, paired where a slight distal oral shield is formed by the enlargement of the distal pair of oral spines . . . . 5. ***PELMATOPORA*** Lang
        (ii) Costae numerous with numerous pelmata or pelmatidia and lateral costal fusions ; avicularia large, variously directed, of one, two or three kinds . . . . . 6. ***CASTANOPORA*** Lang
(II) Pelmatoporinae with a well-developed secondary orifice and a tertiary front wall 7. ***UBAGHSIA*** (Jullien)

---

[1] My italics.

Twelve other genera were placed by Lang (1916*a*; 1922) in his subfamilies Castanoporinae and Pelmatoporinae.

In the Castanoporinae he included seven genera other than those revised here. Provisionally I use Lang's definitions of these seven genera to distinguish them from those revised. Thus, according to Lang (1922 : 158), *Carydiopora, Anornithopora, Hesperopora* and *Stichocados* are distinguished from the other genera here placed in the revised subfamily of the Pelmatoporinae by their small size and fewer costae; *Phrynopora* by its unusual type of "secondary aperture" [proximal oral shield]; and *Steginopora* and *Disteginopora* by their types of tertiary front walls.

Similarly, Lang (1922 : 240) included in his subfamily Pelmatoporinae *Decurtaria, Murinopsia, Sandalopora, Ichnopora* and *Pachydera.* Various characters of these five genera, particularly the types of oral shields which may be developed, distinguish them from other genera placed in the revised subfamily of the Pelmatoporinae.

Genus ***AEOLOPORA*** Lang

1916*a* *Aeolopora* Lang, pp. 383, 390.
1921 *Aeolopora* Lang: Lang, p. 158.
1924 *Aeolopora* Lang: Gaster, second of three tables opp. p. 110.
1927 *Aeolopora* Lang: Canu & Bassler, p. 23.
1930 *Aeolopora* Lang: Gaster, table opp. p. 340.
1930 *Aeolopora* Lang: Voigt, p. 498.
1935 *Aeolopora* Lang: Bassler, p. 43.
1953 *Aeolopora* Lang: Bassler, p. G189.

TYPE SPECIES (by original designation). *Aeolopora distincta* Lang, 1916*a* : 390. Senonian, zone of *M. cor-anguinum.* Upper Basildon, Berkshire.

EMENDED DIAGNOSIS. Small Pelmatoporinae with a single boss-like pelmatidium near the distal end of each costa; no lateral costal fusions; ovicells hyperstomial, prominent, with labellum and lateral slits.

REMARKS. As is mentioned below, the boss-like structure near the distal end of each costa, which is characteristic of the genus, is interpreted as an unusual form of pelmatidium.

Lang (1916*a* : 382, 383) originally diagnosed *Aeolopora* as a multiserial, encrusting, Andrioporine genus lacking a "secondary aperture" [orifice], but with the "median area of fusion surrounded by a ring of solid spines; [and with] apertural [oral] spines thickened". This diagnosis remained almost unchanged in his subsequent key to the Andrioporine genera (1921 : 88, 89).

Lang (1921 : 160), in his remarks on *A. distincta,* stated that "at the margin of the median area of fusion each costa bears a bead, which, to judge from comparable instances, is the original constricted distal end of the costa, which has retreated from the mid-line." It has been possible, using a magnification of ×140, to see that the so-called "beads" are not solid but possess a minute central pore which apparently connects with the central hollow of the costa. The structure is therefore regarded as a pelmatidium with an unusually high rim.

Remarking on the genus *Aeolopora,* Lang (1921 : 158, 159) also stated that "A tendency to this arrangement [the development of 'solid spines or beads' round the median area of fusion] has already been noticed in *Andriopora gasteri* ... *Eucheilopora* ... *Pancheilopora*, and again in the species *Andriopora gallica* and *A. frequens*." These species of *Andriopora* and all available specimens belonging to the other two genera have been examined for the feature mentioned. At the most, there is a tendency for irregular median ridges to form, but circlets of boss-like pelmatidia, or of " solid spines or beads ", are definitely not present.

The presence of pelmatidia in *Aeolopora* removes the genus from both the subfamily Andrioporinae Lang, 1916*a*, and the family Andrioporidae Lang, 1916*a*, and places it in the subfamily Pelmatoporinae. The boss-like pelmatidia particularly distinguish the genus, which is always small and lacks lateral costal fusions, but possesses frequent, prominent, hyperstomial ovicells.

In the key to the families of Cretaceous cribrimorph Polyzoa Lang (1921 : 4–6) made a basic division of all families into (*a*) those with hyperstomial ovicells but without pelmata or pelmatidia and (*b*) those with endozooecial ovicells and pelmata or pelmatidia. *Aeolopora* is clearly a genus which has distinct hyperstomial ovicells and pelmatidia, though the latter are of an unusual form. The wider classification of the families and subfamilies of some of the Cretaceous cribrimorph Polyzoa is discussed elsewhere (p. 52).

STRATIGRAPHICAL DISTRIBUTION. Senonian, lower part of zone of *M. cor-anguinum* to zone of *B. mucronata.*

Of the two species *A. distincta* and *A. nebulosa*, here regarded as composing the genus, the former is apparently more abundant in the Senonian of southern England, ranging from the lower part of the zone of *M. cor-anguinum* into the zone of G. [*A.*] *quadrata*, while the latter has been recorded only from the zone of G. [*A.*] *quadrata* (or its equivalent, see Voigt, 1930 : 498) and from that of *B. mucronata* in the Isle of Wight.

A. nebulosa is somewhat larger and more robust and may be regarded as a development from *A. distincta*. Apparently *A. nebulosa* occurs much less frequently.

KEY TO SPECIES OF *AEOLOPORA*

(A) *Aeolopora* with 10–16, often 11–13, costae ; apertural bar of variable thickness, with or without two distinct boss-like pelmatidia near the mid-line ; avicularia occasionally present ; ovicells hyperstomial with labella and lateral slits 1. ***A. distincta*** Lang

(B) *Aeolopora* with larger zooecia ; usually 9–12 costae ; apertural bar constantly thickened and prominent with a small median process above two indistinct boss-like pelmatidia ; avicularia apparently absent ; ovicells hyperstomial, with small labella and lateral slits . . . . . . 2. ***A. nebulosa*** Lang

1. ***Aeolopora distincta*** Lang

(Pl. 1 ; Pl. 2, figs. 1–7 ; Text-figs. 12–17)

1916*a* *Aeolopora distincta* Lang, p. 390.
1916*a* *Aeolopora stellata* Lang, p. 390.
1921 *Aeolopora distincta* Lang : Lang, p. 159, pl. 5, fig. 4 ; text-fig. 75.
1921 *Aeolopora stellata* Lang : Lang, p. 161, text-fig. 76.
1924 *Aeolopora stellata* Lang : Gaster, second of three tables opp. p. 110.

1930 *Aeolopora stellata* Lang : Gaster, table opp. p. 340.
1935 *Aeolopora distincta* Lang : Bassler, p. 43.
1953 *Aeolopora distincta* Lang : Bassler, p. G189, text-fig. 143, 2.

HOLOTYPE. D.28062. Zoarial fragment encrusting a piece of echinoid test. Senonian, high in the zone of *M. cor-anguinum.* Upper Basildon, north-west of Pangbourne, Berkshire. L. Treacher Collection.

EMENDED DIAGNOSIS. *Aeolopora* with orifice of equal height and width, or hr exceeds lr very slightly ; oral spines 6, costae 10–16, but very often 11–13 only ; apertural bar of variable thickness with or without two distinct boss-like pelmatidia near the mid-line ; heterozooecia (? kenozooecia) occasionally present ; avicularia present ; ovicells abundant, hyperstomial, with labella and lateral slits.







FIG. 12. *Aeolopora distincta* Lang, D.40599. Orifice and ovicell with complete ectooecium and well-formed labellum and lateral slits.

FIG. 13. *Aeolopora distincta* Lang, D.28062. Holotype. Ovicelled zooecium.

FIG. 14. *Aeolopora distincta* Lang, D.28911. Holotype of *A. stellata* Lang. Ovicelled zooecium.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* closely placed, oval. *Orifice* small, distally rounded, the sides straight, or tending to diverge proximally, the proximal-lateral corners may be somewhat rounded, proximal margin straight to widely " V "-shaped. *Oral spines* 6, apparently secondarily thickened, but small, arranged on the distal-lateral margins and sides of the orifice, usually paired symmetrically. *Frontal shield* well arched. *Costae* variable from 10–16, but very often 11–13, fairly closely placed, widest at their proximal (outer) ends and tapered distally. Each costa bears near its distal (inner) end a small, prominent, boss-like *pelmatidium.*[1] There are no *lateral costal fusions.* The costae slope upwards to the pelmatidia and then flatten to join the *median area of fusion*, which is flat, of variable width, and may be occasionally absent. *Apertural bar* variable in thickness, often narrow, fairly straight or widely " V "-shaped, usually with, but sometimes apparently lacking, a pair of distinct boss-like pelmatidia near the mid-line. The apertural bar is never so consistently thickened as in *A. nebulosa*. *Gymnocyst* only slightly exposed. There is no *interzooecial secondary tissue.*

---

[1] The " bead or solid spine " of Lang (1921 : 161).





FIG. 15. *Aeolopora distincta* Lang, D.40578. Orifice and ovicell, the latter with a lateral growth of ectooecium.

FIG. 16. *Aeolopora distincta* Lang. D.40559. Two ovicelled zooecia and interzooecial avicularium. Both ovicells are damaged but that on the left shows the corrugated ectooecium.

Avicularia interzooecial, occurring infrequently in any zoarium, fairly large, oval, distally directed and with slight lateral constrictions (Text-fig. 16). In most zoaria avicularia are absent.

Heterozooecia (? kenozooecia) occasionally present, very small (one-fifth to one-quarter the size of the normal zooecia), convex, smooth, each with a large oval aperture on the upper surface, occurring singly or grouped in irregular clusters between the zooecia (seen only in D.40658).

Ovicells abundant, hyperstomial, prominent, globular, smooth, with a distinct *labellum* bounded by short *lateral slits.* The endooecium of the primary ovicell may be covered by a slightly corrugated ectooecium (Text-figs. 12, 15, 16). The ovicelled zooecia sometimes occur in linear series.

MEASUREMENTS. *Zooecia* Lz = 0·27 to 0·33 mm.
lz = 0·19 to 0·25 mm.
hr = 0·05 to 0·07 mm.
lr = 0·05 to 0·06 mm.
Avicularia Lz = 0·05 mm.
lz = 0·03 mm.

REMARKS. Lang (1916*a* : 390) originally referred two species, *A. distincta* Lang and *A. stellata* Lang, to *Aeolopora*, but the present revision shows that they are synonymous ; his later (1921) species, *A. nebulosa*, is clearly definable, however.



FIG. 17. *Aetopora distincta* Lang, D.40571. Zooecium, with very worn ovicell, showing arrangement of oral spine-bases.

A. distincta and *A. stellata* were separated primarily on the number of " beads surrounding the median area of fusion " and on the form of the apertural bar. All the available specimens of *Aeolopora*, labelled by Lang as belonging to these species, have been re-examined, and the results may be tabulated briefly as follows :

	Specimens assigned by Lang to *A. distincta*	Specimens assigned by Lang to *A. stellata*
Lz .	0·30 to 0·33 mm. .	0·27 to 0·32 mm.
lz .	0·20 to 0·25 mm. .	0·19 to 0·23 mm.
hr .	0·05 to 0·07 mm. .	0·06 mm.
lr .	0·05 to 0·06 mm. .	0·05 to 0·06 mm.
Number . of costae	10–16 (often 12 or 13)	11–15 (often 11 or 12)

As Lang indicated (1921 : 159, 161), the dimensions of the two species are virtually the same.

The apertural bar of *A. distincta*, according to Lang (1921 : 160), " is somewhat flattened in a vertical plane, and each of the costae of which it is composed retains the bead formed by its constricted distal end, and this takes its place in the ring of beads ". In contrast, he stated (p. 161), the apertural bar of *A. stellata* " is considerably flattened in a vertical plane, and, unlike the apertural bar of *A. distincta*, does not normally bear a median pair of beads ".

I have not found it possible to distinguish between the two by the form of the apertural bar. The structure is easily modified by wear, and is in itself variable both in the zooecia of a single zoarium and in those of different zoaria of each species recognized by Lang. The apertural bars of adjacent zooecia may be of varying thickness, and the boss-like pelmatidia, which are present near the mid-line, may or may not be distinct (cf. Text-figs. 13–16) : they are often visible in the specimens labelled by Lang *stellata* and are not, as he suggested, " not normally present ".

The number of boss-like pelmatidia necessarily varies with the number of costae, and, as is apparent from the above table, the range in number of costae of *A. stellata* is contained within that of *A. distincta*.

Thus the two species are regarded as synonymous, the name *distincta* being retained for them.

Many new specimens, ranging from the zone of *M. cor-anguinum* to that of *G.* [*A.*] *quadrata*, have been assigned to the species, and although much of this material has been measured it has not proved necessary to extend the ranges established in the re-examination of Lang's specimens. Many specimens show the details of the ovicell, particularly D.28062, D.28911, D.40578, D.40559 and D.40653. Heterozooecia are present in D.40658, (Pl. 1, fig. 5).

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum* to zone of *G.* [*A.*] *quadrata*.

SPECIMENS. D.28062. Holotype—see above.

D.28911. Holotype of *A. stellata* Lang, encrusting a fragment of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 2 of Gaster, by reservoir, near Hill Barn, North Lancing, east of Worthing, Sussex.

D.21212–13. Two paratypes of *A. distincta* Lang. Horizon, locality and collection as for D.28056 and D.28052 respectively.

D.23326–28. Three paratypes of *A. distincta* Lang, D.23326–27 encrusting *Inoceramus* fragments, D.23328 encrusting a piece of echinoid test. Senonian, top of zone of *M. cor-anguinum*. Medical College Pit, Epsom, Surrey. F. Möckler Collection.

D.23950. Paratype of *A. distincta* Lang, six well-preserved zooecia encrusting a very small shell fragment. Horizon, locality and collection as for D.23326.

D.28052–55. Four paratypes of *A. distincta* Lang. Four or five incomplete zoaria encrusting one fragment of echinoid test [the fifth zoarial fragment numbered D.21213]. Senonian, zone of *M. cor-anguinum*. Upper Basildon, north-west of Pangbourne, Berkshire. L. Treacher Collection.

D.28056–61 and D.28063–66. Ten paratypes of *A. distincta* Lang. Ten of twelve incomplete zoaria encrusting one fragment of echinoid test [the other

	two zoarial fragments numbered D.21212 and D.28062—the latter the holotype]. Horizon, locality and collection as for the holotype.
D.28246.	Paratype of *A. distincta* Lang, encrusting a fragment of *Inoceramus*. Senonian, zone of *M. cor-anguinum*, about 40 ft. from base of zone (20 ft. below the " strong *cor-anguinum* tabular " [flint band] of Whitaker and Rowe). Cliffs between Beltout and Birling Gap, west of Beachy Head, Sussex.
D.3248.	Paratype of *A. stellata* Lang, encrusting a fragment of *Inoceramus*. Senonian, zone of *G.* [*A.*] *quadrata*. Romsey, south-west of Winchester, Hampshire. H. P. Blackmore Collection.
D.28918–21.	Four paratypes of *A. stellata* Lang, encrusting fragments of echinoid tests. Horizon, locality and collection as for D.28911.
D.28965–66.	Two paratypes of *A. stellata* Lang, encrusting one fragment of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 7 of Gaster, in Lambleys Lane, Sompting, south of Lambleys Barn, north of Upton Cottages, north-west of Sompting Church (just above the 200 ft. contour).
D.28986.	Paratype of *A. stellata* Lang, encrusting a fragment of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 10 of Gaster, Western Pit, Charman Dean, north of Broadwater, north of Worthing, Sussex.
D.29885.	Incomplete zoarium labelled " ?*A. stellata* Lang ", encrusting a fragment of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs between last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.
D.29888.	Specimen labelled "*A. distincta* Lang ", encrusting a fragment of echinoid test. Senonian, zone of *M. cor-anguinum*. Coombs Pit, West Horsley, north-east of Guildford, Surrey.
D.29889–90.	Two specimens labelled " *A. stellata* Lang ", the former encrusting a fragment of *Ostrea*, the latter a fragment of *Inoceramus*. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 2 of Gaster—see D.28911.
D.29891.	Specimen labelled " *A. stellata* Lang ", encrusting a fragment of *Inoceramus*. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit 4 of Gaster. Western Pit, below Lancing Ring, west side of Boundstone Lane, north-east of Worthing, Sussex.
D.29892.	Specimen labelled " *A. stellata* Lang ", encrusting a fragment of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Locality and collection as for D.29885.
D.29893.	Specimen labelled " *A. stellata* Lang ", encrusting a fragment of echinoid test. Senonian, zone of *Marsupites testudinarius*, subzone of *Uintacrinus westphalicus*. Medical College Pit, Epsom, Surrey.
D.40555.	Zoarial fragment encrusting part of a *Pelmatopora* zoarium. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 3 of Gaster, east pit at top of Boundstone Lane, North Lancing, near Worthing, Sussex.

D.40556. One of two incomplete zoaria encrusting a fragment of echinoid test [ancestrula present]. Horizon as for D.40555, Pit 4 of Gaster, western pit at top of Boundstone Lane, North Lancing, near Worthing, Sussex.

D.40559–61. Three incomplete zoaria—D.40559–60 encrusting the same *Inoceramus* fragment, D.40561 encrusting a piece of echinoid test. Horizon as for D.40555, Pit 2 of Gaster—see D.28911.

D.40562. Incomplete zoarium encrusting a small, whole valve of *Ostrea*. Horizon, locality and collection as for D.40555.

D.40563–80. Eighteen specimens, all encrusting fragments of echinoid tests. Two specimens—Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*, Pit 2 of Gaster (see D.28911—above), the rest from the subzone of *E. scutata* var. *depressula* of Section 113 of Gaster—cliffs between the last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.

D.40581–94. Fourteen specimens, all encrusting fragments of echinoid tests, except D.40581–83, D.40587–88 and D.40590–91 which encrust fragments of *Inoceramus*. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pits 2 and 4 of Gaster—see D.28911 and D.29891 respectively.

D.40595–98. Four specimens encrusting fragments of *Inoceramus*. Senonian, zone of *M. cor-anguinum* (less than 10 ft. below strong *cor-anguinum* tabular). Cliffs, Beltout, near Eastbourne, Sussex.

D.40599. Specimen encrusting a fragment of *Inoceramus*. Senonian, zone of *M. cor-anguinum*. Southern England—precise locality and collection unknown.

D.40600. Specimen encrusting a fragment of *Inoceramus*. Senonian, base of zone of *M. cor-anguinum* (about 20 ft. below the strong *cor-anguinum* tabular). Cliffs between Beltout and Birling Gap, west of Beachy Head, Sussex.

D.40601. Incomplete zoarium encrusting the same echinoid fragment as D.40556.

D.40602–05. Four specimens, D.40602–03 encrusting fragments of echinoid tests, D.40604–05 encrusting fragments of *Inoceramus*. Senonian, zone of *M. cor-anguinum* (*Trochiliopora* Bed). Cliffs, Seaford district, Sussex.

D.40606–18. Thirteen specimens, all encrusting echinoid fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 10 of Gaster, Charman Dean, near Worthing, Sussex.

D.40619. Specimen encrusting a fragment of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 24 of Gaster, Warningcamp Hill, Warningcamp, north-east of Arundel, Sussex.

D.40620–23. Four specimens encrusting fragments of echinoid tests. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 26 of Gaster, South Woodleighs Pit, east of gravel pit, north-east of Woodleighs Lodge, about half a mile west of Pit 24, Warningcamp, north-east of Arundel, Sussex.

D.40624–26. Three specimens encrusting fragments of echinoid tests. Senonian, zone of *G.* [*A.*] *quadrata*. Locality, horizon and collection as for D.40606.

D.40627. Specimen encrusting a fragment of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 2 of Gaster. Horizon, locality and collection as for D.28911.

D.41064. Zoarial fragment encrusting a piece of *Ostrea* shell. Senonian, zone *O. pilula*, subzone of *E. scutata* var. *depressula*. Section 113 of Gaster, cliffs Rottingdean, Sussex.

D.40628. Specimen encrusting a fragment of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Locality and collection as for D.40606.

D.40629–31. Three specimens, encrusting fragments of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 12 of Gaster, large pit north of Pumping Station, top of Waterworks Lane, north-east of "The Warren", north of Broadwater and Worthing, Sussex.

D.40632–33. Two specimens encrusting fragments of echinoid tests. Senonian, either lowest part of zone of *G.* [*A.*] *quadrata* or topmost part of zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 3 of Gaster, eastern pit at top of Boundstone Lane, south of Lancing Ring, north of Fir Croft, North Lancing, Sussex.

D.40634–36. Three specimens, encrusting fragments of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 15 of Gaster, large pit about one-third of a mile south-south-east of Salvington Windmill, and north of Hill Cottages, Salvington, Sussex.

D.40637–44. Eight specimens, all encrusting inner surfaces of fragments of echinoid tests. Senonian, zone of *G.* [*A.*] *quadrata*. South side of Pit 1143 of Brydone. Redhampton Limeworks Pit, Portsdown, Hampshire.

D.40645. Specimen encrusting a fragment of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 16 of Gaster. Pit by main road, east of Swandean Isolation Hospital and half a mile south of Salvington Windmill, Durrington, Sussex.

D.40646–54. Nine specimens, all encrusting fragments of echinoid tests. Horizon, locality and collection as for D.40606.

D.40655–57. Three specimens, encrusting one fragment of echinoid test. Horizon, locality and collection as for D.40634.

D.40658. Incomplete zoarium with heterozooecia, encrusting a fragment of echinoid test. Horizon, locality and collection as for D.28965.

D.38852. Zoarial fragment encrusting an *Inoceramus*. Upper Chalk. Grove Park Road, Bury St. Edmonds, Suffolk. J. L. Gilbert Collection.

2. ***Aeolopora nebulosa*** Lang

(Pl. 2, figs. 8, 9; Text-figs. 18, 19)

1921 *Aeolopora nebulosa* Lang, p. 163, pl. 5, fig. 5; text-fig. 77.
1924 *Aeolopora nebulosa* Lang: Gaster, second of three tables opp. p. 110.
1930 *Aeolopora nebulosa* Lang: Voigt, pp. 498, 542, 560, pl. 27, fig. 1.

Holotype. D.29061. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 7 of Gaster, in Lambleys Lane, Sompting, south of Lambleys Barn, north of Upton Cottages, north-west of Sompting Church, just above the 200 ft. contour.

Emended diagnosis. *Aeolopora* with somewhat larger, more robust zooecia than those of *A. distincta*; orifice of equal height and width, or hr exceeds lr very slightly or vice versa; oral spines 6, considerably thickened; costae usually 9–12 (occasional adult zooecia may have 8 or 13); apertural bar prominent, considerably thickened, wide, its distal face nearly vertical, noticeably produced upwards in the mid-line, with or without two distinct boss-like pelmatidia; avicularia apparently absent; ovicells prominent, hyperstomial, with small labellae and lateral slits.



Fig. 18. *Aeolopora nebulosa* Lang, D.29061. Holotype. Zooecium with very thick apertural bar.

Fig. 19. *Aeolopora nebulosa* Lang, D.29061. Holotype. Zooecium with narrow apertural bar and well-developed oral spine bases.

Description. *Zoarium* encrusting, unilaminar. *Zooecia* closely placed, oval, somewhat larger than those of *A. distincta* and more robust. *Orifice* small, always distally rounded, the sides straight, proximal margin slightly curved. *Oral spines* 6, prominent, considerably thickened but otherwise unmodified, usually paired and evenly spaced. *Frontal shield* well arched. *Costae* thicker and more robust than in *A. distincta*, usually 9–12, but occasionally adult zooecia may possess only 8, and less frequently may have 13. The costae narrow inwards, and each bears a small, prominent, boss-like *pelmatidium* fairly close to its distal (inner) end. There are no *lateral costal fusions*. The outline of the distal extremity of each costa may often be traced in the *median area of fusion*, which is relatively flat, and usually narrower than in *A. distincta*. Outwards from the boss-like pelmatidia the costae slope down to meet the gymnocyst less steeply than in *A. distincta*. *Apertural bar* prominent, considerably thickened, wide or very wide, its distal face vertical, produced upwards in the mid-line to form a small *median process*. Beneath the median process and proximal to it, a pair of boss-like pelmatidia may or may not be distinct. The apertural bar is generally straighter than that of *A. distincta*. *Gymnocyst* only slightly exposed. There is no *interzooecial secondary tissue*.

Avicularia apparently absent. *Heterozooecia* not seen. *Ovicells* prominent, hyperstomial, globular, smooth with a small *labellum* and *lateral slits*. If an ectooecium is present it is not corrugated—as may be the case with *A. distincta*.

MEASUREMENTS. *Zooecia* Lz = 0·30 to 0·40 mm.
lz = 0·20 to 0·30 mm.
hr = 0·06 to 0·08 mm.
lr = 0·06 to 0·07 mm.

REMARKS. The dimensions given by Lang for *A. nebulosa* (Lz = about 0·40 mm. and lz = about 0·30 mm.) were maximum measurements. It is, however, apparent that *A. nebulosa* may be distinguished from *A. distincta* by its rather larger size, its more robust appearance, its definitely thicker oral spines, its thicker, and generally fewer costae, and its more constantly thickened apertural bar which may be very prominent (Text-figs. 18, 19). The labellum tends to be smaller than in *A. distincta.* Avicularia have not been seen in any of the specimens examined and their absence may again be of diagnostic value.

Lang (1921 : 163) described the apertural bar of *A. nebulosa* as " thick, flattened in a vertical plane, and produced medianly in a spine-like projection ". This median projection may not always be present and is never very marked ; it is particularly prone to wear. The apertural bar of *A. distincta* also tends to be thickened medianly and this may occasionally produce a median projection. It is not true to describe the form of the orifice as " cribriline "[1] (Lang, 1921 : 163) ; the proximal-lateral constrictions characteristic of this type of orifice are not present.

Voigt (1930 : 498, pl. 27, fig. 1) very briefly recorded a specimen which he assigned to *A. nebulosa* Lang from the Mammillatensenon of Ifö, Sweden.

Specimens D.40660–65, from the zone of *B. mucronata* of the Isle of Wight extend the upward range of the species. The holotype, D.29061, is from the zone of *G.* [*A.*] *quadrata* of Sussex and the paratype specimens from the same horizon in Wiltshire. No specimens of *A. nebulosa* have been discovered as yet below this horizon. Much new material belonging to the genus, has been examined, but apart from the specimens from the zone of *B. mucronata* all have proved to belong to *A. distincta* Lang.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *G.* [*A.*] *quadrata* to zone of *B. mucronata.*

SPECIMENS. D.29061. Holotype—see above.

D.3251. Paratype encrusting a fragment of ?echinoid test. [D.29052–53 are probably parts of the same specimen.] Senonian, zone of *G.* [*A.*] *quadrata*. East Harnham, south of Salisbury, Wiltshire. H. P. Blackmore Collection.

---

[1] See Lang, 1921 : xlvii.

D.29052–53.	Two paratypes [poorly preserved], encrusting. Horizon, locality and collection as for D.3251.
D.40660.	Zoarial fragment, encrusting a piece of echinoid test. Senonian, zone of *B. mucronata*. Pit 9 of Rowe, south of the Briary, between Alum Bay and Freshwater, Isle of Wight.
D.40661–65.	Five incomplete zoaria, encrusting fragments of echinoid tests. Senonian, zone of *B. mucronata*. Pit 6 of Rowe, south of Newbarn, Afton Down, Isle of Wight.

Genus ***LAGYNOPORA*** Lang

?1874	*Lepralia* : Reuss, p. 129 [*partim*].
1877	*non Lepralia* : Novák, pp. 82, 93 [*partim*].
1889	*non Lepralia* : Fric, p. 90.
1893	*Cribrilina* : Vine, pp. 316, 336 [*partim*].
?1900*b*	*Cribrilina* : Canu, p. 445 [*partim*].
1904	*Cribrilina* : Jukes-Browne, p. 490 [*partim*].
1910	*Cribrilina* : Brydone, p. 391.
1916*a*	*Prodromopora* Lang, p. 394.
1916*a*	*Lagynopora* Lang, p. 394.
1921	*Lagynopora* : Lang, pp. lxi, xci–v, 52.
1921	*Prodromopora* : Lang, pp. lxiii, xciv, 50, 51.
1927	*Lagynopora* : Canu & Bassler, p. 17.
1927	*Prodromopora* : Canu & Bassler, p. 36.
1930	*Prodromopora* : Kühn, p. 65.
?1930	*Canupora* Kühn, p. 65.
1930	*Lagynopora* : Gaster, table opp. p. 340.
1930	*Lagynopora* : Voigt, pp. 495, 542, 560.
?1935	*Canupora* : Bassler, pp. 30, 64.
1935	*Prodromopora* : Bassler, pp. 30, 176.
1935	*Lagynopora* : Bassler, pp. 30, 135.
1949	*Lagynopora* : Voigt, pp. 36, 43, 44.
?1953	*Canupora* : Bassler, p. G188.
1953	*Prodromopora* : Bassler, p. G188.
1953	*Lagynopora* : Bassler, p. G188.
1958	*Lagynopora* : Didon, pp. 80, 84.

TYPE SPECIES. *Lagynopora lagena* Lang, 1916*a* : 395. Senonian, zone of *M. cortestudinarium* or zone of *M. cor-anguinum*. Chatham, Kent. W. Gamble Collection.

EMENDED DIAGNOSIS. Pelmatoporinae with a median process on the apertural bar which bifurcates and fuses with the proximal pair of oral spines producing a prominent oral shield ; avicularia usually present.

REMARKS. *Lagynopora* may be clearly distinguished from other genera of the Pelmatoporinae by its distinctive proximal oral shield. In specimens which have been worn the proximal shield may often be destroyed. Usually, however, a few of these structures survive. Also it is possible to infer their presence originally in worn zooecia by the form of the apertural bar, and, in contrast to those of *Hexacanthopora*, by the lengthened and enlarged proximal oral spines, originally fused to the median process.

Prodromopora Lang is synonymous with *Lagynopora*. Lang (1916*a* : 394; 1921 : 50) regarded the few, narrow, widely spaced costae and the slight median area of fusion in *Prodromopora* as generically significant. Other characters were recognized as similar to those of *Lagynopora*. The characters of the frontal shield are, however, of specific value only, and *Prodromopora praecursor* Lang is placed in *Lagynopora*.

Canupora prima Kühn (1930 : 65) is here provisionally referred to *Lagynopora* and is listed and described as a species of uncertain systematic position (p. 96). I exclude from this genus *Lepralia pediculus* Reuss, which Lang doubtfully referred to *Lagynopora* (see p. 95).

Excluding *Lagynopora*? *prima* (Kühn), 12 species are recognized, one of which, *L. pustulosa*, is new.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cortestudinarium* to zone of *B. mucronata* in England, zone of *G.* [*A.*] *quadrata* in Germany, and ?Danian in Austria.

KEY TO THE SPECIES OF *Lagynopora* LANG

(I) *Lagynopora* with boss-like pelmata . . . . . . 1. ***L. pustulosa*** sp. nov.
(II) *Lagynopora* in which the pelmata are minute pores.
  (A) Distinct lateral costal fusions present . . . . 2. ***L. horsleyensis*** Lang
  (B) No lateral costal fusions.
    (1) Intercostal spaces wide, as long as the costae; median area of fusion very slight; costae narrow (11–14) . . . . . 3. ***L. praecursor*** (Lang)
    (2) Intercostal spaces narrower, but still distinct and nearly as long as the costae.
      (a) Zooecia small, Lz = 0·43 to 0·60 mm., lz = 0·25 to 0·37 mm.
        (i) Costae 11–14; avicularia occasional, fairly large, oval 4. ***L. saltdeanensis*** Lang
        (ii) Costae 17–18; avicularia abundant, small, oval 5. ***L. birlingensis*** Lang
      (*b*) Zooecia large, Lz = 0·64 to 0·75 mm., lz = 0·37 to 0·50 mm.; costae 13–17; frontal shield often with a median ridge . 6. ***L. amphora*** Lang
    (3) Intercostal spaces limited to the outer margins of the frontal shield.
      (*a*) Costae 12–14. . . . . 7. ***L. ampulla*** Lang
      (*b*) Costae 20–23. . . . . 8. ***L. furcifera*** (Brydone)
    (4) Intercostal spaces absent; costae juxtaposed, outlined by furrows only.
      (*a*) Zooecia small, Lz = 0·48 to 0·68 mm., lz = 0·35 to 0·51 mm.
        (i) Costae 9–11; oral spines 4 or 5; avicularia of one kind, large, prominent, oval . . . . 9. ***L. ollula*** Lang
        (ii) Costae 13–20; oral spines always 4; avicularia of two sizes 10. ***L. lagena*** Lang
      (*b*) Zooecia large, Lz = 0·65 to 0·80 mm., lz = 0·35 to 0·50 mm.
        (i) Median area of fusion extends only along the inner third of each costa; costae 12–18; avicularia very large, broadly oval, prominent, fairly frequent, directed just proximally and outwards from the zooecia they accompany . . 11. ***L. urceolus*** Lang
        (ii) Median area of fusion extends along the inner two-thirds of each costa; costae 15–17; avicularia rare or absent . 12. ***L. vasculum*** Lang

Lepralia pediculus Reuss[1] and *Lagynopora ?prima* (Kühn)[2] are of uncertain systematic position and have therefore been omitted from the above key.

1. *Lagynopora pustulosa*[3] sp. nov.

(Pl. 3, fig. 1 ; Text-figs. 20, 21)

HOLOTYPE. D.39419. Small zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *B. mucronata.* Felpham Chalk, Bognor, Sussex.

PARATYPES. D.39420–22. Three zoarial fragments encrusting one piece of echinoid test. Senonian, zone of *B. mucronata.* Pit 8 of Rowe, near Nodewell, Freshwater, Isle of Wight.



FIGS. 20, 21. *Lagynopora pustulosa* sp. nov., D.39419. Holotype. Front and side view of adult zooecium.

a.b, apertural bar ; *d.o.s*, distal oral spine-bases ; *f*, fusion between median process of apertural bar and proximal oral spines ; *l.w*, lateral wall ; *m.p*, median process of apertural bar ; *o*, orifice.

DIAGNOSIS. *Lagynopora* with zooecia of small size (Lz = 0·50 to 0·53 mm., lz = 0·27 to 0·32 mm.) ; proximal oral shield heavy, large, prominent, very obtusely " Y "-shaped, with a marked median furrow ; frontal shield well arched ; costae 13–16 ; each costa with a distinct small, boss-like pelma at its inner end ; ovicells hyperstomial, globular, small.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* elliptical, closely placed, each with a small, round, *distal communication pore. Secondary orifice* distally and laterally rounded, proximal margin straight. *Oral spines* 4, thickened, arranged symmetrically on the distal margin of the orifice. *Proximal oral shield* very prominent, large, very obtusely " Y "-shaped, with a straight distal margin and a marked median furrow, formed by the fusion of the bifurcated median process of the apertural bar with the proximal pair of oral spines. *Frontal shield* well arched. *Costae* 13–16, fairly narrow, tapering inwards and each with a small, boss-like *pelma* very close to its inner end. The pelmata produce two rows of small bosses, one on either side of the mid-line. Distinct, narrow *intercostal spaces* separate the costae. No *lateral costal fusions*. *Median area of fusion* slight. *Apertural bar* wider than the most distal costae, straight, the inner ends of its two component costae curve distally at the mid-line to form the prominent, wide *median process* which bifurcates to form part of the proximal oral shield. *Gymnocyst* concealed by the close placing of the zooecia.

---

[1] Lang (1916*a* : 395 ; 1921 : 61) very tentatively placed this species in *Lagynopora.*

[2] *Canupora prima* (Kühn) (1930 : 65) may be a *Lagynopora,* but the available details are insufficient to place it with any certainty.

[3] With reference to the form of the pelmata.

Avicularia apparently absent. *Ovicells* prominent but small, smooth, hyperstomial, globular, of about equal height and width.

MEASUREMENTS. *Zooecia* Lz = 0·50 to 0·53 mm.
lz = 0·27 to 0·32 mm.
hr = 0·09 mm.
lr = 0·15 mm.

[The majority of zooecia in the available specimens are ovicelled or slightly damaged, thus making accurate measurements based on non-ovicelled zooecia difficult.]

REMARKS. This species is clearly distinguished by its boss-like pelmata and its large, prominent and distinctive proximal oral shield.

In the holotype, D.39419, young zooecia are present but not sufficiently well preserved to show whether 4 or 6 oral spines are typically developed at this stage of zoarial growth. The young zooecia apparently do not possess proximal oral shields. The same remarks apply to the young zooecia of the paratypes. In D.39422 some ovicelled zooecia possess prominent, long, proximal oral spines—these seem to be the remains of broken proximal oral shields.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *B. mucronata*.

SPECIMENS. D.39419. Holotype—see above.
D.39420–22. Paratypes—see above.

2. *Lagynopora horsleyensis* Lang

(Pl. 3, figs. 2, 3 ; Text-fig. 22)

1916*a* *Lagynopora horsleyensis* Lang, pp. 395, 396.
1921 *Lagynopora horsleyensis* Lang : Lang, pp. xciii, 57, pl. 2, fig. 3 ; text-fig. 28.
?1958 *Lagynopora horsleyensis* Lang : Didon, pp. 80, 84, text-fig. 2.

HOLOTYPE. D.28908. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Pit 264 of Young, Coombs Pit, West Horsley, north-east of Guildford, Surrey.

EMENDED DIAGNOSIS. *Lagynopora* with 14–17 costae, each with 3, occasionally only 2, distinct lateral costal fusions ; avicularia occasional, large, oval, interzooecial, directed distally and obliquely outwards from the zooecia they accompany.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* broadly oval, fairly closely placed, with large, oval *distal communication pores*. *Secondary orifice* wider than high, rounded distally with straight sides and a straight proximal margin. *Proximal oral shield* probably formed by the fusion of the lateral oral spines with the median process of the apertural bar ; this structure is usually destroyed by wear. *Oral spines* 4, somewhat thickened, but otherwise unmodified. *Frontal shield* flatly arched. *Costae* 14–17, tapering inwards, each with 3, occasionally only 2, distinct *lateral costal fusions*. Outermost *intercostal spaces* long and slot-like, inner intercostal spaces small and regular. *Median area of fusion* narrow, outwards from the mid-line the costae are almost horizontal but turn down fairly steeply beyond the outermost lateral costal fusions to meet the gymnocyst. Minute *pelmatidia* are present on the costae at the levels of the lateral costal fusions, and at least one other pelmatidium apparently occurs near the outer end of each costa. *Apertural bar* a little wider than the most distal costae, distally bent at its outer ends and with a *median*



FIG. 22. *Lagynopora horsleyensis* Lang, N.C.M.247.956 (4/2). Adult zooecium with damaged apertural bar, the proximal oral shield worn away. A large interzooecial avicularium with proximal-lateral constrictions is shown to the left of the zooecium.

process. This probably fused with the lateral oral spines to produce a proximal oral shield. *Gymnocyst* apparently smooth, exposed proximally to some frontal shields, but generally obscured by the close placing of the zooecia. *Interzooecial secondary tissue* apparently absent.

Avicularia interzooecial, of one kind, large, oval but somewhat parallel-sided, unpaired, with proximal-lateral constrictions separating a smaller proximal portion from a larger, rounded *rostrum* ; apparently directed distally and obliquely outwards from the zooecia they accompany. *Ovicells* prominent, smooth, hyperstomial, usually longer than wide, tending to be pointed distally.

MEASUREMENTS. *Zooecia* Lz = 0·47 to 0·57 mm.
lz = 0·30 to 0·40 mm.
hr = 0·09 to 0·10 mm.
lr = 0·12 to 0·15 mm.
Avicularia Lz = 0·15 to 0·22 mm.
lz = 0·10 to 0·17 mm.

REMARKS. *Lagynopora horsleyensis* is easily distinguished from other species of the genus by its distinct lateral costal fusions. It is unfortunate that in the available specimens all the proximal oral shields have been destroyed by wear, only the worn median process of the apertural bar and the somewhat lengthened and thickened form of the lateral oral spines suggesting the original presence of this structure. D.40942 extends the upward range of the species into the zone of *G.* [*A.*] *quadrata.*

Lang (1921 : 57) slightly overestimated the dimensions of *L. horsleyensis* and its number of costae. His text-figure 28 and pl. 2, fig. 3, do not indicate the very distinct form of the lateral costal fusions.

Didon (1958) has assigned to this species a poorly preserved specimen from the Santonian of Le Trait, near Rouen, France.

STRATIGRAPHICAL DISTRIBUTION. Senonian, upper part of zone of *M. cor-anguinum* to zone of *G.* [*A.*] *quadrata* and ? Santonian of Le Trait, near Rouen, France.

SPECIMENS. D.28908. Holotype—see above.

D.29899. Incomplete zoarium encrusting a shell fragment. Senonian, zone of *M. cor-anguinum.* Pit in Houndean Bottom, west of Lewes, Sussex.

D.40942. Incomplete zoarium encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 12 of Gaster, Broadwater Pit, north of waterworks pumping station, top of Waterworks Lane, north of Broadwater, near Worthing, Sussex.

D.41069. Small zoarial fragment of six young zooecia encrusting a piece of *Inoceramus* shell. Senonian, high in the zone of *M. cor-anguinum.* Locality and collection as for D.29899.

OTHER MATERIAL

N.C.M.3951. Zoarial fragment with numerous ovicells encrusting the base of a whole test of *Conulus albogalerus.* Senonian, zone of *M. cor-anguinum.* Northfleet, Kent. A. W. Rowe Collection.

3. ***Lagynopora praecursor*** (Lang) nov. comb

(Pl. 3, fig. 5 ; Text-figs. 23, 24)

1916*a* *Prodromopora praecursor* Lang, p. 394.
1921 *Prodromopora praecursor* Lang : Lang, pp. lxiii, xciv, 51, pl. 2, fig. 1 ; text-fig. 25.
1935 *Prodromopora praecursor* Lang : Bassler, p. 176.
1953 *Prodromopora praecursor* Lang : Bassler, p. G188, text-fig. 142, 8.

HOLOTYPE. D.8351. Zoarial fragment with many damaged ovicelled zooecia and a broken ancestrular area, encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Gillingham, north-east of Chatham, Kent. W. Gamble Collection.



FIGS. 23, 24. *Lagynopora praecursor* (Lang), D. 8351. Holotype.

(23) Ovicelled adult zooecium, proximal oral shield worn away ; *d.o.s*, distal oral spines ; *o*, outline of ovicell of preceding zooecium.

(24) Interzooecial avicularium, of the same zoarium, showing lateral constrictions. The avicularium is figured in the normal orientation to the zooecium—directed distally and obliquely outwards.

EMENDED DIAGNOSIS. *Lagynopora* with small zooecia (Lz = about 0·60 mm., lz = 0·40 to 0·42 mm.) ; frontal shield well arched ; costae 11–13, narrow, separated by wide intercostal spaces; median area of fusion narrow; avicularia frequent, apparently of one kind only, large (Lz = 0·23 mm., lz = 0·15 to 0·20 mm.), oval or almost parallel-sided with proximal-lateral constrictions, directed distally and often obliquely outwards from the zooecia they accompany; ovicells hyperstomial, often with a slight median ridge and small lateral slits.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* oval, fairly closely placed, each with a large, round *distal communication pore*. *Secondary orifice* wider than high, rounded distally and laterally with a straight proximal margin. *Oral spines* 4, arranged symmetrically; the distal pair are smaller than the proximal-lateral pair and may flank the entrance to the ovicell when this is present. The proximal-lateral pair are enlarged; they very probably fused with the median process of the apertural bar to form a *proximal oral shield*. *Frontal shield* well arched, sloping down steeply at the margins to meet the gymnocyst. *Costae* 11–13, narrow, fused only near the mid-line and separated by wide *intercostal spaces*. Some costae retain minute *pelmatidia* along their mid-line. *Apertural bar* well formed, only slightly wider than the most distal costae, with a *median process* which probably fused with the proximal-lateral oral spines to form the proximal oral shield. *Gymnocyst* exposed proximally to most frontal shields and narrowly along their sides. No *interzooecial secondary tissue*.

Avicularia of one kind, very large, frequent, directed distally and often obliquely outwards from the zooecia they accompany. The avicularia are oval or almost parallel-sided with well-marked proximal-lateral constrictions which may be the remains of worn transverse bars. Sometimes the constrictions may be completely removed by wear. *Ovicells* prominent, hyperstomial, globular, smooth, of medium size, often with a slight median ridge and with small *lateral slits*. *Young zooecia* smaller than the adult zooecia and with 6 oral spines. *Ancestrula* apparently with 6 oral spines.

MEASUREMENTS. *Adult zooecia* Lz = about 0·60 mm.
lz = 0·40 to 0·42 mm.
hr = 0·10 to 0·12 mm.
lr = 0·14 to 0·15 mm.
Avicularia Lz = 0·23 mm.
lz = 0·15 to 0·20 mm.
Young zooecia Lz = 0·37 to 0·42 mm.
lz = 0·26 to 0·30 mm.
hr
lr } Not measurable.
Ancestrula Lz = 0·29 mm.
lz = 0·21 mm.
hr
lr } Not measurable.

REMARKS. *Prodromopora praecursor* Lang is placed in *Lagynopora*, as it apparently has a proximal oral shield formed by the fusion of the proximal-lateral oral spines with the median process of the apertural bar. Further, a prominent distal communication pore is present, there are 4 oral spines in the adult zooecia, and large, distally directed oval avicularia are present. All these characters are typical of *Lagynopora.* Unlike most other species of *Lagynopora,* there are only a few widely spaced costae which fuse only along the mid-line of the frontal shield: this distinctive frontal shield is only specifically diagnostic.

Only one specimen, D.8351, is available. Apart from three adult zooecia the zoarium is badly damaged. It is, however, possible, using high-power magnification, to distinguish minute pelmatidia on the costae of these zooecia.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum.*

SPECIMEN. D.8351. Holotype—see above.

4. ***Lagynopora saltdeanensis*** Lang

(Pl. 3, fig. 4; Text-figs. 25, 26)

1916*a* *Lagynopora saltdeanensis* Lang, p. 395.
1921 *Lagynopora saldeanensis* Lang: Lang, pp. xcii, 56, text-fig. 27.
1930 *Lagynopora saltdeanensis* Lang: Gaster, table opp. p. 340.

HOLOTYPE. D.28896. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula,* subzone of *E. scutata* var. *depressula.* Cliffs between last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.



FIGS. 25, 26. *Lagynopora saltdeanensis* Lang, D.28896. Holotype.
(25) Adult ovicelled zooecium in which the proximal oral shield has been worn away.
(26) Distal half of another adult ovicelled zooecium with two worn interzooecial avicularia.

Emended diagnosis. *Lagynopora* with fairly small zooecia (Lz = 0·43 to 0·45 mm., lz = 0·25 to 0·27 mm.); secondary orifice fairly small (hr = 0·08 mm., lr = 0·10 mm.); costae, 11–14, with distinct intercostal spaces, but no lateral costal fusions; avicularia occasional, fairly large, oval; ovicells hyperstomial.

Description. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, closely placed. *Secondary orifice* small, wider than high, rounded distally with straight sides and a straight proximal margin. *Oral spines* apparently 4, not much thickened. *Proximal oral shield* formed by the fusion of the lateral oral spines with the median process of the apertural bar. Nearly all zooecia are worn, retaining only the abraded median process on the apertural bar. *Frontal shield* well arched. *Costae* 11–14, narrow, tapering inwards, united along the mid-line in a narrow *median area of fusion*. Outwards from the median area the costae slope down fairly steeply to meet the gymnocyst. Two or three small *pelmatidia* are visible on the upper surfaces of most costae, but they may be obscured by wear. *Apertural bar* straight with a *median process* which is fused with the proximal-lateral oral spines to produce the proximal oral shield. *Gymnocyst* smooth, fairly widely exposed proximally to each frontal shield, but it may be concealed by the close placing of the zooecia or by the small amounts of *interzooecial secondary tissue* which may occur.

Avicularia occasional, unpaired, fairly large, oval, and of uncertain orientation. *Ovicells* large, smooth, prominent, hyperstomial, globular, of about equal height and width and apparently without *lateral slits*. *Young zooecia* similar to the adult zooecia, but smaller and with fewer costae.

Measurements. *Adult zooecia* Lz = 0·43 to 0·45 mm.
lz = 0·25 to 0·37 mm.
hr = 0·08 mm.
lr = 0·10 mm.
Young zooecia Lz = 0·30 to 0·33 mm.
lz = 0·20 to 0·25 mm.
hr = 0·06 mm.
lr = 0·09 mm.
Avicularia Not measurable.

Remarks. *L. saltdeanensis* is smaller than *L. birlingensis* and possesses fewer costae, and has, apparently, fewer, larger, sporadic avicularia. The only available specimen of *L. saltdeanensis* is the holotype—D.28896. It possesses only a few well-preserved zooecia, at least two of which are representative of a young stage of zoarial growth. It might thus be argued that D.28896 is merely an early growth stage of *L. birlingensis*. The problem can only be settled when new material becomes available.

The form of the proximal oral shield in *L. saltdeanensis* is unknown, but this feature has generic rather than specific diagnostic value and its original presence can be determined with certainty from the form of the worn orifice.

Stratigraphical distribution. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*.

Specimen. D. 28896. Holotype—see above.

5. *Lagynopora birlingensis* Lang

(Pl. 3, fig. 6; Text-figs. 27, 28)

1916*a* *Lagynopora birlingensis* Lang, p. 395.
1921 *Lagynopora birlingensis* Lang: Lang, pp. xciv, 55, pl. 2, fig. 2; text-fig. 26.
1930 *Lagynopora birlingensis* Lang: Gaster, table opp. p. 340.

HOLOTYPE. D.28254. Incomplete zoarium encrusting a fragment of *Inoceramus* shell. Senonian, zone of *M. cor-anguinum.* Cliffs between Beltout and Birling Gap, west of Beachy Head, Sussex.



FIGS. 27, 28. *Lagynopora birlingensis* Lang, D.28254. Holotype.
(27) Adult zooecium with worn orifice, the proximal oral shield worn away, and worn, oval avicularium.
(28) Ovicell and orifice, partly obscured by matrix, of another adult zooecium—the proximal oral shield is partly preserved; *m.p*, median process of apertural bar; *p.o.s*, proximal oral spine.

EMENDED DIAGNOSIS. *Lagynopora* with fairly large zooecia (Lz = 0·50 to 0·60 mm., lz = 0·30 to 0·37 mm.); secondary orifice of normal size (hr = 0·10 mm., lr = 0·12 to 0·16 mm.); costae numerous (17–18) with distinct intercostal spaces; no lateral costal fusions; avicularia interzooecial, oval, fairly abundant; ovicells hyperstomial, longer than wide.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, fairly closely placed. *Secondary orifice* wider than high, rounded distally and laterally, proximal margin fairly straight. *Oral spines* 4, somewhat thickened but otherwise unmodified, arranged in two pairs on the distal-lateral margins of the orifice. *Proximal oral shield* very obtusely " Y "-shaped, formed by the fusion of the proximal-lateral pair of oral spines with the median process of the apertural bar. *Frontal shield* fairly well arched. *Costae* 17–18, narrow, tapering inwards and separated by distinct, narrow *intercostal spaces* : no *lateral costal fusions* present. The costae are united about the mid-line into a narrow *median area of fusion* in which may be traced the outlines of the inner ends of individual costae. Minute *pelmatidia* are present, possibly three or four on each costa. *Apertural bar* prominent with a *median process* forming part of the proximal oral shield—it is often destroyed by wear leaving a straight apertural bar with a worn median process. *Gymnocyst* exposed proximally to some frontal shields but generally obscured by the small amount of irregular but widespread *interzooecial secondary tissue.*

Avicularia interzooecial, unpaired, numerous, small, rounded to oval, of uncertain orientation. *Ovicells* hyperstomial, smooth, prominent, globular, somewhat longer than wide and without *lateral slits*. *Young zooecia* smaller than the adult zooecia, with fewer (11 or 12) costae, but otherwise similar.

MEASUREMENTS. *Adult zooecia* Lz = 0·50 to 0·60 mm.
lz = 0·30 to 0·37 mm.
hr = 0·10 mm.
lr = 0·12 to 0·16 mm.
Young zooecium Lz = 0·35 mm.
lz = 0·22 mm.
hr
lr } Not measurable.
Avicularia Lz = 0·08 to 0·12 mm.
lz = 0·07 to 0·11 mm.

REMARKS. *Lagynopora birlingensis* may be distinguished by its dimensions, the shape of the orifice which is wider than high, by the form of its proximal oral shield and by its numerous costae and avicularia.

In his description of the species Lang (1921 : 55) described the orifices as normal to cribriline ; however, the marked proximal-lateral constrictions typical of cribriline orifices are not present. He also commented " aviculoecia [avicularia] dimorphic, both kinds fairly numerous, sporadically distributed, with more or less circular apertures, which, even in the larger kind, are decidedly smaller than those [i.e., the orifices] of the orthoecia [zooecia] and, at least in the larger kind, bear spines on the distal rim ". I have been unable to distinguish two kinds of avicularia and have not seen the spines which Lang mentioned. In his figure of the species (1921 : 12, text-fig. 2) Lang showed, at the lower left-hand side, a large round pore and a smaller one proximal to it. These might be avicularia, but as they are very worn this remains uncertain. The only complete proximal oral shield in the specimen is shown much improved in Lang's figure. Although it is more obscured by matrix than is indicated, it is possible to determine that it is much more obtusely " Y"-shaped than Lang has shown. He regarded *L. birlingensis* as a " very primitive form, approaching *Prodromopora* [= *Lagynopora*] in the thinness and distant spacing of the costae." See also remarks on *L. saltdeanensis* (p. 74).

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum.*

SPECIMENS. D.28254. Holotype—see above.

D.41068 Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum* (above the " strong *cor-anguinum* tabular flint band"). Cliffs between Birling Gap and Beltout, west of Eastbourne, Sussex.

6. *Lagynopora amphora* Lang

(Pl. 3, fig. 9; Text-fig. 29)

1916*a* *Lagynopora amphora* Lang, pp. 395, 396.
1921 *Lagynopora amphora* Lang : Lang, pp. xcii, 59, text-fig. 29.
1930 *Lagynopora amphora* Lang : Gaster, table opp. p. 340.

HOLOTYPE. D.28255. Incomplete zoarium encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Cliffs east of Telscombe Staircase, west of Newhaven, Sussex.



FIG. 29. *Lagynopora amphora* Lang, D.28255. Holotype. Orifice and broken ovicell of adult zooecium. The enlarged proximal oral spines fused originally with the prominent, broken median process of apertural bar.

EMENDED DIAGNOSIS. *Lagynopora* with large zooecia (Lz = 0·64 to 0·75 mm., lz = 0·37 to 0·50 mm.); frontal shield often with a distinct median ridge; costae 13–17 with distinct intercostal spaces; avicularia interzooecial, small, oval, occasional; ovicell hyperstomial, slightly pointed distally, slightly longer than wide.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* very closely placed, broadly oval, each with a large, oval distal, and a smaller, oval *lateral communication pore. Secondary orifice* wider than high, distally and laterally rounded with a straight proximal margin. *Oral spines* 4, slightly enlarged, arranged symmetrically on the distal and lateral margins of the orifice. *Proximal oral shield* developed, but form unknown, all the oral shields in the available specimens having been destroyed by wear. *Frontal shield* evenly arched, often with a distinct median ridge. *Costae* 13–17, tapering slightly inwards, separated by narrow but distinct *intercostal spaces*. No *lateral costal fusions*. Each costa possesses a series of minute *pelmatidia* on its upper surface. *Median area of fusion* slight, often consisting of the median ridge only, or extending only slightly on either side of it. The outline of the inner extremity of each costa is visible in the median area of fusion. *Apertural bar* wider than the most distal costae, somewhat triangular, with distally curved ends, and with a prominent, fairly wide *median process* which almost certainly fuses, when complete, with the proximal-lateral pair of oral spines to form a proximal oral shield. *Gymnocyst* not exposed, owing to the very close placing of the zooecia. No *interzooecial secondary tissue*.

Avicularia interzooecial, occasional, small, with oval apertures. *Ovicells* hyperstomial, prominent, smooth, globular, longer than wide, sometimes tending to be slightly pointed distally. *Young zooecia* generally resemble the adult zooecia, but are smaller and have 6 oral spines, and slightly fewer (11–14) costae.

Measurements. *Adult zooecia* Lz = 0·64 to 0·75 mm.
lz = 0·37 to 0·50 mm.
hr = 0·12 to 0·13 mm.
lr = 0·15 to 0·16 mm.
Young zooecia Lz = 0·40 to 0·47 mm.
lz = 0·30 to 0·35 mm.
hr = 0·11 to 0·12 mm.
lr = 0·13 mm.
Avicularia Not measurable.

Remarks. *Lagynopora amphora* is distinguished from other species of the genus particularly by the characters of its frontal shield, which, in contrast to that of *L. ampulla*, is evenly arched and unconsolidated. The median ridge is especially distinctive.

Stratigraphical distribution. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*, and lower part of zone of *B. mucronata* (not yet recorded from the zone of *G.* [*A.*] *quadrata*).

Specimens. D.28255. Holotype—see above.

D.40943. Incomplete zoarial fragment with young and adult zooecia, some ovicelled, encrusting a piece of echinoid test. Horizon, locality and collection as for the holotype.

Other material

N.C.M.76.937(32), 76.937(33). Two zoarial fragments encrusting one piece of echinoid test. Senonian, lower part of zone of *B. mucronata*, Wells, Norfolk. R. M. Brydone Collection.

7. *Lagynopora ampulla* Lang

(Pl. 3, figs. 7, 8 ; Text-fig. 30)

1916*a*	*Lagynopora ampulla* Lang, pp. 395, 396.
1921	*Lagynopora ampulla* Lang : Lang, pp. xcii, 60, text-fig. 31.
1930	*Lagynopora ampulla* Lang : Gaster, table opp. p. 340.

HOLOTYPE. D.28256. Specimen in five fragments. Incomplete zoarium encrusting a cheilostome polyzoan. Senonian, zone of *G.* [*A.*] *quadrata*. East side of Old Nore Point, west of Newhaven, Sussex.



FIG. 30. *Lagynopora ampulla* Lang, D.28256. Holotype. Adult ovicelled zooecium with small, round, worn avicularium (*a*). The ovicell is slightly damaged proximally ; the proximal oral shield is well preserved showing the slender fusions with lengthened proximal oral spines and wide median process of apertural bar. *o*, outline of the ovicell of the preceding zooecium.

EMENDED DIAGNOSIS. *Lagynopora* with large zooecia (Lz = 0·62 to 0·75 mm., lz = 0·35 to 0·50 mm.); proximal oral shield obtusely " Y "-shaped, prominent, flat and wide in front with a distinct median furrow, and slender fusions with the proximal-lateral oral spines; costae 12–14, closely placed, no lateral costal fusions.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, closely placed, with large, oval *distal communication pores* and smaller *lateral communication pores*. *Secondary orifice* wider than high, distally rounded with straight sides and a straight proximal margin, the sides of the orifice may converge slightly proximally. The form of the secondary orifice is often obscured by the well-developed proximal oral shield or by ovicells. *Oral spines* 4, slightly lengthened, somewhat thickened, arranged symmetrically on the distal rim of the orifice. *Proximal oral shield* very prominent, obtusely " Y "-shaped, flat and wide in front, composed of the bifurcated, enlarged median process of the apertural bar which fuses, on either side, with the proximal-lateral oral spines. *Frontal shield* very flatly arched, well consolidated. *Costae* 12–14 (often 13). Each costa is fairly flat and tapered inwards. Adjacent costae are in contact and horizontal for most of their length. At their outer ends they bend down steeply to meet the gymnocyst and are separated by short, narrow *intercostal spaces*. Although the costae are in contact with one another, their outlines remain distinct, even in the median area. *Median area of fusion* extremely wide. Minute *pelmatidia* are visible on the upper surfaces of some costae. *Apertural bar* wider than the most distal costae, straight, or slightly curved distally at either end, with a strong *median process* which bifurcates and fuses with the proximal-lateral oral spines to produce the proximal oral shield. *Gymnocyst* apparently smooth, but little exposed owing to the close placing of the zooecia. No *interzooecial secondary tissue*.

Avicularia very occasional, small, circular, interzooecial. *Ovicells* hyperstomial, prominent, smooth, globular, of about equal height and width. The entrance to the ovicell may be prominently flanked by a pair of oral spines. *Young zooecia* similar to the adult zooecia, but smaller and with 5 oral spines, and generally fewer (10–12) costae.

MEASUREMENTS. *Adult zooecia* Lz = 0·62 to 0·75 mm.
lz = 0·35 to 0·50 mm.
hr = 0·14 to 0·15 mm.
lr = 0·15 to 0·17 mm.
Young zooecia Lz = 0·40 to 0·47 mm.
lz = 0·25 to 0·32 mm.
hr = 0·08 to 0·10 mm.
lr = 0·09 to 0·14 mm.
Avicularia Lz = 0·04 mm.
lz = 0·04 mm.

REMARKS. *L. ampulla* is distinguished from other species of *Lagynopora* by its well consolidated frontal shield, short intercostal spaces, and particularly by its proximal oral shield. Lang (1921 : 60) overestimated the zooecial dimensions of *L. ampulla*.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *G.* [*A.*] *quadrata.*
SPECIMEN. D.28256. Holotype—see above.

8. *Lagynopora furcifera* (Brydone)

(Pl. 4, figs. 1, 2; Text-figs. 31, 32)

1910 *Cribrilina furcifera* Brydone, pp. 391, 392, 482 [*partim*—pl. 30, fig. 7 only, *non* fig. 6 (= *Lagynopora vasculum*), *non* fig. 8 (= *L. ollula*)].
1916*a* *Lagynopora* [*Cribrilina*] *furcifera* (Brydone): Lang, p. 395 [*partim*—that referring to Brydone, 1910, pl. 30, fig. 7 only].
1917 *Cribrilina furcifera* Brydone: Brydone, p. 492.
1921 *Lagynopora furcifera* (Brydone) Lang, pp. xci, 67.
?1930 *Lagynopora furcifera* (Brydone): Voigt, pp. 495, 542, 560, pl. 26, fig. 10.
?1949 *Lagynopora* aff. *furcifera* (Brydone): Voigt, pp. 36, 43, 44, pl. 11, fig. 1.

LECTOTYPE. (Chosen by Lang, 1921 : 67.) S.M., B.36165. An extensive, well-preserved zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata.* Bramford, near Ipswich, Suffolk. R. M. Brydone Collection.



FIGS. 31, 32. *Lagynopora furcifera* (Brydone), S.M., B.36165. Lectotype.
(31) Adult zooecium and worn avicularium, the proximal oral shield destroyed.
(32) Ovicell and orifice of another adult zooecium with complete proximal oral shield and part of an avicularium.

EMENDED DIAGNOSIS. *Lagynopora* with very large zooecia (Lz = 0·75 to 0·85 mm., lz = 0·50 to 0·56 mm.); proximal oral shield very prominent, very widely " Y "-shaped, composed of a wide median process which rises from the apertural bar, bifurcates, and fuses with the proximal-lateral pair of oral spines; frontal shield well arched; costae very numerous, 20–23, distinct, separated by intercostal spaces for at least half their length; median area of fusion wide; avicularia interzooecial, fairly frequent, oval.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, closely placed, each with a large, oval *distal communication pore* and one or two, somewhat smaller, oval *lateral communication pores*. *Secondary orifice* wider than high, rounded distally and laterally with a straight proximal margin. *Oral spines* 4, slightly enlarged, symmetrically arranged round the orifice. *Proximal oral shield* prominent, formed by the fusion of the proximal-lateral pair of oral spines with the wide, flat median process of the apertural bar. *Frontal shield* slightly flattened in the median area but otherwise well arched. *Costae* 20–23, narrow, well defined, separated at their outer ends, and for about half their length, by distinct *intercostal spaces*; the outlines of their inner ends may be traced across the wide *median area of fusion*. No *lateral costal fusions*. Minute *pelmatidia* are visible on the upper surfaces of unworn costae. *Apertural bar* wide, somewhat swollen, triangular, the apex of the triangle pointing proximally. The outer ends of the apertural bar are slightly bent proximally to the frontal shield, but generally obscured by the close placing of the zooecia, or by avicularia. *Interzooecial secondary tissue* absent.

Avicularia interzooecial, frequent, fairly large, oval. [In the lectotype all the avicularia are worn.] *Ovicells* hyperstomial, frequent, smooth, prominent, globular, some may be slightly pointed distally with a slight median ridge. The proximal-lateral margins of the ovicells normally incorporate the distal pair of oral spines. *Young zooecia* smaller than the adult zooecia but otherwise similar, except that some of the very young zooecia may possess 5 or 6 oral spines, and the orifice tends to be of equal height and width.

MEASUREMENTS. *Adult zooecia* Lz = 0·75 to 0·85 mm.
lz = 0·50 to 0·56 mm.
hr = 0·15 to 0·16 mm.
lr = 0·18 to 0·20 mm.
Young zooecia Lz = 0·57 to 0·63 mm.
lz = 0·40 to 0·52 mm.
hr = 0·15 mm.
lr = 0·15 mm.
Avicularia Lz = About 0·20 mm.
lz = About 0·13 mm.

REMARKS. Brydone figured three specimens with his original description of *Cribrilina furcifera*, 1910, pl. 30, figs. 6–8). Lang (1916*a* : 395) placed these three specimens in a single species of *Lagynopora*. Subsequently (1921) he suggested that the material figured by Brydone as *Cribrilina furcifera* belonged to three separate species of *Lagynopora* (that shown in fig. 6 being *L. vasculum*, that in fig. 7—*L. furcifera*, and that in fig. 8—*L. ollula*).

Having examined Brydone's figured specimens, I agree that they are three separate species of *Lagynopora*. I have accordingly extended the brief diagnoses given by Lang [these were based solely on Brydone's somewhat inadequate figures] and have added full descriptions and measurements.

Lagynopora furcifera (Brydone) possesses more costae than *L. vasculum* or *L. ollula*, and is larger. The degree of consolidation of the arched frontal shield and the avicularia also further distinguish *Lagynopora furcifera* from other species of the genus.

Voigt (1930 : 495, pl. 26, fig. 10) assigned to the species a specimen from the " Quadraten-oder Mukronatensenon " of Misburg. Both the description, in which he stated only that there are about 20 costae and that Lz = 0·70 mm., and the figure, which is quite indistinct, are inadequate to place the material with any certainty. More recently (Voigt, 1949 : 36, pl. 11, fig. 1) has described and figured, as " *Lagynopora* aff. *furcifera* (Brydone) ", another specimen from the " Quadratenkreide " of Lägerdorf, Holstein. It is slightly larger than the lectotype, with slightly fewer costae and with 7 oral spines in the youngest zooecia.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *G.* [*A.*] *quadrata* of England and Germany.

MATERIAL. S.M., B.36165. Lectotype—see above.

9. *Lagynopora ollula* Lang

(Pl. 4, fig. 3 ; Text-figs. 33–35)

1910 *Cribrilina furcifera* Brydone, p. 391 [*partim*—pl. 30, fig. 8 only].
1916*a* *Lagynopora furcifera* (Brydone) : Lang, p. 395 [*partim*—that referring to Brydone, 1910, pl. 30, fig. 8 only].
1921 *Lagynopora ollula* Lang, pp. xciv, 60.
1930 *non Lagynopora furcifera* (Brydone) : Voigt, pp. 495, 542, 560, pl. 26, fig. 10.
1949 *non Lagynopora* aff. *furcifera* (Brydone) : Voigt, pp. 36, 43, 44, pl. 11, fig. 1.

LECTOTYPE. (Chosen by Lang, 1921 : 60.) S.M., B.36166. The specimen figured by Brydone (1910, pl. 30, fig. 8). Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites*, subzone of *Uintacrinus*. Kingsgate, Kent. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Lagynopora* with large zooecia (Lz = 0·58 to 0·67 mm., lz = 0·40 to 0·51 mm.) ; orifice of constant size (hr = 0·12 mm., lr = 0·15 mm.) ; proximal oral shield very prominent, obtusely " Y "-shaped, the flat, wide median process standing out almost at right angles to the plane of the apertural bar ; frontal shield well consolidated ; costae 9–11, broad, closely placed, fused to neighbouring costae throughout most of their length ; avicularia interzooecial, unpaired, large, prominent, oval, with proximal-lateral constrictions.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* closely placed, very broadly oval, each with a very large, oval *distal communication pore* and several smaller, oval *lateral communication pores*. *Secondary orifice* wider than high, of constant size, distally rounded with straight sides and a straight proximal margin. The form of the orifice may be obscured by the proximal oral shield or by ovicells. *Oral spines* 4 or 5, arranged symmetrically round the orifice. The proximal-lateral pair of oral spines is often larger than the distal pair, all are slightly secondarily thickened. *Proximal oral shield* very prominent, obtusely " Y "-shaped, formed by the flat, wide median process of the apertural bar which bifurcates and fuses with the lateral pair of oral spines. *Frontal shield* wide, flat, except at the margins where the costae descend steeply to meet the gymnocyst. *Costae* 9–11, distinct, wide, rounded, tapering inwards. Each costa is completely fused to its neighbouring costae for most of its length, but the outline of each costa is clearly visible up to the mid-line of the frontal shield. Only at their outer ends are the costae clearly separate, and the frontal shield is well consolidated and the *median area of fusion* extremely wide. Minute *pelmatidia* are visible on the upper surfaces of unworn costae. *Apertural bar* straight, except for a distal curve at either end, swollen, wider than the most distal costae. A strong, flat, wide *median process* rises almost



Figs. 33–35. *Lagynopora ollula* Lang, S.M., B.36166. Lectotype.

(33) Adult ovicelled zooecium; the proximal oral shield is broken on one side.

(34) Orifice of another adult zooecium with a fifth distal-median oral spine-base; the proximal oral shield broken.

(35) Avicularium with proximal-lateral constrictions.

perpendicularly from the centre of the apertural bar and forms part of the proximal oral shield. *Gymnocyst* smooth, but seldom exposed owing to the close placing of the zooecia; usually visible round the interzooecial avicularia. No *interzooecial secondary tissue.*

Avicularia interzooecial, unpaired, large, round or oval, probably with a *transverse bar,* now broken, and represented by proximal-lateral constrictions, originally dividing a smaller semicircular proximal portion from a larger, transversely oval *rostrum.* Apparently the avicularia are variously directed; some are definitely distally directed obliquely away from the zooecia they accompany, others may be directed proximally and obliquely towards the zooecia they accompany.

Ovicells hyperstomial, very prominent, smooth, globular, as wide as high. The proximal-lateral margin of the ovicell may incorporate or obscure the distal-lateral pair of oral spines. Some ovicells tend to be slightly pointed distally.

Young zooecia smaller than the adult zooecia. Orifice of less constant size than in the adult stages. Oral spines 6 or 7. Proximal oral shield probably developed. Frontal shield more arched than in the adult. Costae more numerous, 10–12, but, as in the adult stages, fused for most of their length. Apertural bar wide, straight.

Ancestrula [of lectotype]. Only the orifice is visible; it is wider than high, and there are 8 closely spaced, symmetrically arranged oral spines.

MEASUREMENTS. *Adult zooecia* Lz = 0·58 to 0·67 mm.
lz = 0·40 to 0·51 mm.
hr = 0·12 mm.
lr = 0·15 mm.
Avicularia Lz = 0·12 to 0·14 mm.
lz = 0·19 to 0·15 mm.
Young zooecia Lz = 0·40 to 0·43 mm.
lz = 0·23 to 0·26 mm.
hr = 0·07 to 0·10 mm.
lr = 0·10 to 0·11 mm.
Ancestrula $\left.\begin{matrix}\text{Lz}\\ \text{lz}\end{matrix}\right\}$ Not measurable.
hr = 0·07 mm.
lr = 0·10 mm.

REMARKS. *Lagynopora ollula* (Brydone) is clearly distinguished from other species of the genus by its much consolidated, wide frontal shield, its relatively few costae, the constant size of its orifice, and particularly by the characters of its avicularia.

In the lectotype, S.M., B.36166, where ovicells have been preserved they apparently afforded some protection for the proximal oral shield, for this is most complete in the ovicelled zooecia. In non-ovicelled zooecia, although the proximal oral shield is very prominent, it is frequently destroyed by wear so that the apertural bar appears as a straight, featureless structure with no trace of the very strong median process present in complete individuals. Very slight wear of the surfaces of the costae in this well-preserved specimen has been sufficient to obscure most of the minute pelmatidia which may now be seen only occasionally on the upper surfaces of some costae.

The lectotype zoarium, although small, is almost complete, for it possesses the ancestrula and its surrounding young zooecia, ovicelled adult zooecia, and, in one place, a small zooecium possibly close to part of the growing edge of the zoarium.

Fewer oral spines are developed in the adult than in the young stages, and the ancestrula possesses the largest number of oral spines, 8. This reduction in the number of oral spines during growth is apparently accompanied by a slight reduction in the number of costae.

Lang (1921 : 60) has suggested that *L. ollula*, apart from its well-consolidated frontal shield, shows " primitive characters ".

Mr. A. G. Brighton has kindly informed me that Brydone (*in litt.* 23rd June, 1938) stated that the lectotype came from the " *Uintacrinus* Band " of Kingsgate, Kent. See also remarks under *L. furcifera* (Brydone), p. 82.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *Marsupites*, subzone of *Uintacrinus*.

MATERIAL. S.M., B.36166. Lectotype—see above.

10. ***Lagynopora lagena*** Lang

(Pl. 4, figs. 4–6 ; Text-figs. 36, 37)

1893	*Cribrilina nitidiformis* Vine, pp. 316, 336 [*nomen nudum—vide* Lang, 1921 : 62].
1904	*Cribrilina nitidiformis* Vine : Jukes-Browne, p. 490 [*vide* Lang, 1921 : 62].
1916*a*	*Cribrilina nitidiformis* Vine : Lang, p. 410 [*vide* Lang, 1921 : 62].
1916*a*	*Lagynopora lagena* Lang, p. 395.
1921	*Lagynopora lagena* Lang : Lang, pp. lxi, xciv, 62, pl. 2, fig. 4 ; text-fig. 30.
?1930	*Lagynopora* aff. *lagena* Lang : Voigt, pp. 495, 542, 560, pl. 26, fig. 9.
1935	*Lagynopora lagena* Lang : Bassler, p. 135.
1953	*Lagynopora lagena* Lang : Bassler, p. G188, text-fig. 142, 9.

HOLOTYPE. D.4042. Zoarial fragment encrusting a piece of echinoid test. Senonian, either zone of *M. cortestudinarium* or zone of *M. cor-anguinum*. Chatham, Kent. W. Gamble Collection.

EMENDED DIAGNOSIS. *Lagynopora* with zooecia of medium size (Lz = 0·48 to 0·68 mm., lz = 0·35 to 0·46 mm.) ; proximal oral shield more acutely " Y "-shaped than in other species of *Lagynopora*, formed by a fairly narrow median process of the apertural bar which fuses with large, robust proximal-lateral oral spines ; frontal shield evenly arched ; costae 13–20 (often 15–19), juxtaposed but delimited for their whole length by distinct furrows and united only in the narrow median area of fusion which may form a slight ridge ; avicularia frequent, apparently of two sizes, interzooecial, oval with proximal-lateral constrictions ; ovicells hyperstomial, often with a slight median ridge, occasionally with small lateral slits.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, fairly closely placed, each with an oval *distal communication pore*. *Secondary orifice* wider than high, rounded distally and laterally with a straight proximal margin. *Oral spines* 4, symmetrically arranged round the orifice, the proximal-lateral pair slightly larger than the distal pair. *Proximal oral shield* prominent, more acutely " Y "-shaped than in other species of *Lagynopora,* formed by the narrow median process of the apertural bar which fuses with the proximal pair of robust oral spines. *Frontal shield* well arched, with a slight, often ridge-like, narrow *median area of fusion.* *Costae* 13–20, variable in number but often 15–19, narrow, juxtaposed but defined along their whole length by distinct furrows and properly fused only along the mid-line of the frontal shield. No *lateral costal fusions.* Minute *pelmatidia* are visible on the upper surfaces of some well-preserved costae. *Apertural bar* wider than the most distal costae, straight, with a slight distal bend at either end, and with



FIGS. 36, 37. *Lagynopora lagena* Lang.

(36) D.4042. Holotype. Adult ovicelled zooecium with complete proximal oral shield and small oral, worn avicularium of type *b*. The distal oral spines are unusually long.

(37) D.2633. Paratype. Large interzooecial avicularium of type *a*, shown in relation to Fig. 36 in the usual orientation for the species—directed distally and obliquely outwards from the zooecium.

a narrow *median process* which forms part of the proximal oral shield. *Gymnocyst* smooth, only slightly exposed proximally to some frontal shields or more extensively round some avicularia, but generally concealed by the close placing of the zooecia. No *interzooecial secondary tissue.*

Avicularia of one kind, but, in some zoaria, of two sizes, interzooecial, oval with proximal-lateral constrictions which are, apparently, the broken ends of straight *transverse bars.* Where avicularia of both sizes are developed, the larger type (*a*) occurs near the orifice and is directed distally and obliquely outwards from the zooecium it accompanies. The small type (*b*) occurs less frequently and always distal to the orifice or distal to the ovicell when this is present, and is distally directed. In zoaria with both types the avicularia occur frequently, but they are less numerous in zoaria with only the larger form. Where only the larger form of avicularium occurs it may also be placed distally to the orifice. Occasionally avicularia are rare or even absent in a zoarium. No example has been seen in which only the smaller type of avicularium occurs. *Ovicells* hyperstomial, small but prominent, smooth, globular, wider than high and often with a median ridge. Occasional ovicells may have small *lateral slits. Young zooecia* resemble the adult zooecia, but are smaller and may have from 4 to 6 oral spines and fewer costae. Often the youngest zooecia have the most oral spines. *Ancestrulae* lack the proximal oral shield which occurs throughout young and adult zooecia. Some ancestrulae possess 6 oral spines, that of D.11280, however, has only 4.

MEASUREMENTS. *Adult zooecia* Lz = 0·48 to 0·68 mm.
lz = 0·35 to 0·46 mm.
hr = 0·10 to 0·13 mm.
lr = 0·13 to 0·18 mm.
Avicularia :
Type *a* Lz = 0·18 to 0·20 mm.
lz = 0·15 to 0·17 mm.
Type *b* Lz = 0·10 to 0·11 mm.
lz = 0·07 to 0·08 mm.
Young zooecia Lz = 0·40 to 0·45 mm.
lz = 0·26 to 0·33 mm.
hr = 0·07 mm.
lr = 0·11 mm.
Ancestrula (of D.11280) Lz = 0.30 mm.
lz = 0·23 mm.
hr = 0·07 mm.
lr = 0·10 mm.

REMARKS. *Lagynopora lagena*, the type species, was regarded by Lang (1921 : 63) as " the lowest term in a series " which, he suggested, exhibits an increasing consolidation of the frontal shield and a gradual loss of avicularia, first of the smaller kind [type (*b*) in *L. lagena*] and finally of the larger kind also. Certainly *L. lagena* is the only species of *Lagynopora* which possesses more than one size of avicularium, and it also exhibits, though not necessarily in a single zoarium, the widest range in number of costae. Like other species, it undergoes, in the early stages of zoarial growth, a rapid reduction in the number of oral spines.

L. lagena is distinguished by its large number of costae, its well-arched frontal shield and narrow, ridge-like median area of fusion, and by its avicularia and small ovicells.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *M. cortestudinarium* and *M. cor-anguinum* in England. ?Upper part of zone of *G.* [*A.*] *quadrata* in Germany.

SPECIMENS. D.4042. Holotype—see above.

D.4027–28. Two paratype zoarial fragments, the former encrusting an erect cheilostome polyzoan, the latter with only one complete zooecium and now unattached. Horizon, locality and collection as for the holotype.

D.4046. Paratype. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for the holotype.

D.4238. Paratype. Zoarial fragment encrusting a piece of *Inoceramus* shell. Horizon, locality and collection as for the holotype.

D.4969–70, D.4974–76, D.27978–79. Seven paratypes. Zoarial fragments encrusting pieces of echinoid tests. Horizon, locality and collection as for the holotype.

D.2633. Paratype. Zoarial fragment encrusting part of an erect cheilostome polyzoan. Labelled " Syntype of *Cribrilina nitidiformis* Vine 1893 ". Horizon and locality as for the holotype. G. R. Vine Collection.

D.2640. Paratype. Zoarial fragment encrusting a piece of *Inoceramus* shell. Labelled " Syntype of *Cribrilina nitidiformis* Vine 1893 ". Horizon and locality as for the holotype. G. R. Vine Collection.

D.11280. Paratype. Zoarial fragment encrusting a piece of test of *Micraster precursor* Rowe. Senonian, top of zone of *M. cortestudinarium* or base of zone of *M. cor-anguinum*. Chatham, Kent. W. Gamble Collection.

D.21209. Paratype. Zoarial fragment with young zooecia possessing six oral spines encrusting a small fragment of echinoid test. Senonian, zone of *M. cor-anguinum*. Woburn Green, south-west of Beaconsfield, Buckinghamshire. L. Treacher Collection.

D.24236–38. Three paratypes. Zoarial fragments encrusting pieces of echinoid test. Senonian, high in zone of *M. cortestudinarium*. Luton, south-east of Chatham, Kent. W. Gamble Collection.

D.24385. Paratype. Extensive zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Locality and collection as for D.24236–38.

D.24519–20. Two paratypes. Zoarial fragments encrusting pieces of echinoid test, the former with small avicularia of type (*b*). Senonian, zone of *M. cor-anguinum*. Gillingham, north-east of Chatham, Kent. W. Gamble Collection.

D.24525. Paratype. Zoarial fragment with numerous avicularia of type (*a*) and with a regenerated and reversed zooecium, encrusting a piece of echinoid test. Horizon, locality and collection as for D.24519–20.

D.8135, D.24136, D.24541. Three paratypes. Zoarial fragments; D.8135 encrusting an erect polyzoan, the remainder encrusting pieces of echinoid test. Horizon, locality and collection as for D.24519–20.

D.24539, D.24592. Two paratypes. Zoarial fragments, the former encrusting an erect cyclostome polyzoan, the latter encrusting an erect cheilostome polyzoan. Horizon, locality and collection as for D.24519–20.

D.27039–40. Two paratypes. Zoarial fragments encrusting the dorsal surface of pieces of unattached cheilostome polyzoan zoaria. Senonian, zone of *M. cortestudinarium*. Lower Pit, Slines Oak, Worms Heath, Warlingham, Surrey. F. Möckler Collection.

D.27854–56. Three paratypes Zoarial fragments encrusting pieces of echinoid tests. Senonian, zone of *M. cortestudinarium* (upper part). Locality and collection as for D.24519–20.

D.29102. Paratype. Zoarial fragment encrusting three pieces of echinoid test. Senonian, zone of *M. cor-anguinum*. Keston, Kent (south-east of Croydon, Surrey). R. Jones Collection.

11. ***Lagynopora urceolus*** Lang

(Pl. 4, fig. 7; Pl. 5, fig. 1; Text-figs. 38, 39)

1916*a* *Lagynopora urceolus* Lang, pp. 395, 396.
1921 *Lagynopora urceolus* Lang: Lang, p. 65, pl. 2, fig. 5; text-fig. 32.
1930 *Lagynopora urceolus* Lang: Gaster, table opp. p. 340.

HOLOTYPE. D.28889. Zoarial fragment with ancestrula and ovicelled adult zooecia encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs between last groyne east of Rottingdean Gap and Saltdean, Sussex.

EMENDED DIAGNOSIS. *Lagynopora* with zooecia of large size (Lz = 0·65 to 0·78 mm., lz = 0·35 to 0·50 mm.); orifice just wider than high; frontal shield well arched; costae 12–18, closely placed; no marked intercostal spaces; avicularia larger than the orifice, interzooecial, fairly frequent, directed just proximally and outwards from the zooecia they accompany; ovicells large.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, each with a large, circular *distal communication pore* and one or more smaller, oval *lateral communication pores*. *Secondary orifice* just wider than high, rounded distally and laterally with a straight proximal margin. *Oral spines* 4, occurring in two distinct pairs, the proximal pair larger. *Proximal oral shield* obtusely " Y "-shaped, formed by the fusion of the proximal oral spines with the bifurcated median process of the apertural bar. *Frontal shield* well arched, but flattened in the median area. *Costae* 12–18, closely placed, separated by furrows rather than definite *intercostal spaces*, but with their outlines distinct. *Median area of fusion* not very wide, extending along the inner half of the costae at the maximum. The mid-line of the frontal shield may be marked by a slight, irregular ridge. No *lateral costal fusions*. Minute *pelmatidia* are visible on the upper surfaces of well-preserved costae. *Apertural bar* widely triangular, the apex pointing proximally, with a median ridge and a prominent *median process* which forms part of the proximal oral shield. *Gymnocyst* smooth, almost completely concealed by the close placing of the zooecia and by ovicells and avicularia. No *interzooecial secondary tissue.*



FIG. 38. *Lagynopora urceolus* Lang, D. 40944. Adult ovicelled zooecium and interzooecial avicularium. The proximal oral shield is destroyed ; the avicularium, with proximal-lateral constrictions, is directed typically outwards from the zooecium it accompanies.

Avicularia very prominent, larger than the orifice, interzooecial, directed just proximally and outwards from the zooecia they accompany, broadly oval with proximal-lateral constrictions dividing the wider semicircular proximal portion from the somewhat pointed longer *rostrum*. *Ovicells* very prominent, frequent, hyperstomial, large, smooth, globular, without *lateral slits*. *Young zooecia* similar to the adult zooecia, but smaller and with 5 or 6 oral spines and 12–16 costae. *Ancestrula* (of holotype) very small, with 7 oral spines. Number of costae unknown.

MEASUREMENTS. *Adult zooecia* Lz = 0·65 to 0·78 mm.
lz = 0·35 to 0·50 mm.
hr = 0·13 to 0·15 mm.
lr = 0·15 to 0·17 mm.
Avicularia Lz = 0·16 to 0·21 mm.
lz = 0·14 to 0·17 mm.
Young zooecia Lz = 0·35 to 0·47 mm.
lz = 0·23 to 0·30 mm.
hr = 0·10 mm.
lr = 0·10 to 0·12 mm.
Ancestrula (of holotype) Lz = 0·30 mm.
lz = 0·18 mm.
hr = 0·09 mm.
lr = 0·10 mm.



FIG. 39. *Lagynopora urceolus* Lang, D.28889. Holotype. Ovicell and orifice of adult zooecium with proximal oral shield broken on one side.

REMARKS. Lang (1921 : 65) apparently placed very little diagnostic value on the very distinctive avicularia of *L. urceolus* for he did not mention them in his diagnosis of the species. *L. urceolus* is distinguished particularly by its avicularia and to a lesser extent by the dimensions of its zooecia, its number of costae and its well-arched frontal shield.

The outline of individual costae may be traced across the median area of fusion which is narrow rather than " wide " as Lang has stated.

In D.40944, a well-preserved specimen, an avicularium is apparently replaced by a small, reversed, but otherwise normal zooecium.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*, and zone of *Marsupites*, subzone of *Uintacrinus*.

SPECIMENS. D.28889. Holotype—see above.

D.29898. Ovicelled zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs west of Chimney Shaft, Roedean, east of Brighton, Sussex.

D.40944. Ovicelled zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.29898.

D.41064. *Lagynopora* cf. *urceolus*. Somewhat damaged ovicelled zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites*, subzone of *Uintacrinus*. Pit 49 of G. W. Young—Medical College Pit, Epsom, Surrey.

D.41067. Well-preserved, ovicelled zoarial fragment possibly close to part of the growing edge, encrusting a fragment of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Section 113 of Gaster, cliffs east of Rottingdean Gap to Saltdean, Sussex.

12. ***Lagynopora vasculum*** Lang

(Pl. 5, fig. 2; Text-figs. 40, 41)

1910 *Cribrilina furcifera* Brydone, p. 391 [*partim*—pl. 30, fig. 6 only; *non* fig. 7 (= *Lagynopora furcifera*); *non* fig. 8 (= *L. ollula*)].

1916*a* *Lagynopora* [*Cribrilina*] *furcifera* (Brydone): Lang, p. 395 [*partim*—Brydone, 1910, pl. 30, fig. 6 only].

1921 *Lagynopora vasculum* Lang, pp. xciii, 66, text-fig. 33.

LECTOTYPE. (Chosen by Lang, 1921 : 67.) S.M., B.36164. A well-preserved zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Lower part of old pit and road cutting at Mount Pleasant, Andover, Hampshire. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Lagynopora* with large zooecia (Lz = 0·73 to 0·80 mm., lz = 0·45 to 0·50 mm.); proximal oral shield very prominent, obtusely "Y"-shaped; median process of the apertural bar very wide; costae numerous, 15–17, indistinct except towards the margins of the otherwise consolidated frontal shield; intercostal spaces between the outer ends of the costae only; ovicells hyperstomial, large, globular, smooth.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* large, broadly oval, very closely placed, each with a large, prominent, round *distal communication pore* and two or more smaller, oval *lateral communication pores*. *Secondary orifice* wider than high, rounded distally and laterally with a straight proximal margin. *Oral spines* 4, arranged symmetrically round the orifice, the proximal-lateral pair slightly larger than the distal pair. *Proximal oral shield* very prominent, consisting of a very wide median process of the apertural bar; the process bifurcates and fuses with the proximal-lateral pair of oral spines producing a very obtusely "Y"-shaped oral shield. The proximal oral shield when fully developed obscures most of the orifice. *Frontal shield* flatly arched. *Costae* 15–17, fairly narrow. At their outer ends the costae are separated by distinct *intercostal spaces* which extend about one-third of the distance towards the mid-line of the frontal shield. On either side of the mid-line a wide *median area of fusion* is developed, but the outlines of the inner ends of individual costae can be seen in this well-consolidated part of the frontal shield. Minute *pelmatidia* are visible on the upper surface of some unworn costae. No *lateral costal fusions*. No *median ridge* on the frontal shield. *Apertural bar* somewhat swollen, straight except for a slight distal bend at either end, wider than the most distal costae, with a strong, wide *median process* which forms part of the proximal oral shield. *Gymnocyst* smooth, only very occasionally exposed in small,



FIGS. 40, 41. *Lagynopora vasculum* Lang, S.M., B.36164. Lectotype.
(40) Adult ovicelled zooecium with complete proximal oral shield.
(41) Orifice of another adult zooecium in which the proximal oral shield is broken; *d.c.p.* distal communication pore in sloping distal wall of zooecium; *m.p*, median process of apertural bar.

triangular areas proximal to the frontal shields, generally obscured by the very close placing of the zooecia. No *interzooecial secondary tissue*. *Avicularia* are present occasionally. They are apparently interzooecial and fairly large, but in the lectotype only three worn avicularian-like structures have been seen. *Ovicells* hyperstomial, large, prominent, smooth, globular, occasionally slightly pointed distally, their proximal-lateral margins usually incorporating the distal pair of oral spines. *Young zooecia* smaller than the adult zooecia but otherwise similar. Only the first ring of young zooecia have 5 or 6 oral spines ; all successive young and adult zooecia possess only 4 oral spines. *Ancestrula* (almost obscured in the lectotype), with 6 long, narrow oral spines.

MEASUREMENTS. *Adult zooecia* Lz = 0·73 to 0·80 mm.
lz = 0·45 to 0·50 mm.
hr = 0·12 to 0·15 mm.
lr = 0·15 to 0·18 mm.
?*Avicularia* Not measurable.
Young zooecia Lz = 0·43 to 0·46 mm.
lz = 0·30 mm.
hr = 0·10 to 0·11 mm.
lr = 0·11 to 0·12 mm.
Ancestrula Lz, lz, hr = Not measurable.
lr = 0·12 mm.

REMARKS. Lang (1921 : 66) based his diagnosis of *Lagynopora vasculum* on Brydone's figure of the species (1910, pl. 30, fig. 6). The median process of the apertural bar is much wider than is shown in Lang's text-figure 33 (1921 : 66), and the frontal shield is less consolidated.

As in *L. ollula* Lang, there is a rapid reduction in the number of oral spines during the growth of the zoarium, but, unlike *L. ollula*, this is apparently accompanied by a slight increase in the number of costae.

L. vasculum is distinguished mainly by its large size, its numerous costae, and the width of its median area of fusion which extends across about two-thirds of the frontal shield. Avicularia are apparently rare, contrasting with the prominent presence of avicularia in *L. ollula*.

See also remarks under *L. furcifera* (Brydone), p. 82.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*.

MATERIAL. S.M., B.36164. Lectotype—see above.

Species of Uncertain Systematic Position Previously Assigned to *LAGYNOPORA*

(1) ***Lepralia pediculus*** Reuss

1874 *Lepralia pediculus* Reuss, p. 129, pl. 24, fig. 16.
1921 [*Lagynopora*] *pediculus* (von Reuss) Lang, p. 61.

REMARKS. Lang himself (1921 : 61) regarded the systematic position of this species as uncertain, for he placed the name in square brackets and stated (p. 62)

"*Lepralia pediculus*, if a *Lagynopora*, appears to be closely related to *L. lagena*; but the widely-spaced oecia [zooecia] at once distinguish it from that species". In his diagnosis (1921 : 61) he described the species as a "[*Lagynopora*] of large size (oecia about ·66 mm. long); with about 16 closely-approximated costae; lateral costal fusions rare or absent; intraterminal front-wall [frontal shield] fairly well-arched; oecia widely-spaced".

This description was presumably based on that of Reuss (1874 : 129, pl. 24, fig. 16). There are no indications of lateral costal fusions in the figure, nor is there the slightest suggestion of a proximal oral shield—an essential generic character of *Lagynopora*. There is no evidence to suggest that *Lepralia pediculus* Reuss should be placed in *Lagynopora*. The spacing of the zooecia is not of diagnostic value, and Reuss's figure is very stylized showing, at the most, a multiserial cribrimorph polyzoan with a well-arched frontal shield, numerous costae, 6 or 7 oral spines, and hyperstomial, smooth, globular ovicells.

Gillard (1942 : 91) has referred a specimen from a boring at Montbazon, France, to "*Lagynopora pediculus* var. Reuss". Its horizon may be Upper Turonian.

(2) ***Lagynopora ? prima*** (Kühn) nov. comb.

1930	*Canupora prima* Kühn, p. 65, pl. 26, fig. 13.
1935	*Canupora prima* Kühn : Bassler, p. 64.
1953	*Canupora prima* Kühn : Bassler, p. G188.

LECTOTYPE. (Here chosen.) The specimen figured by Kühn, 1930, pl. 26, fig. 13. Danian, Lithothamnienkalk. Haidhof, near Ernstbrunn, Austria.

DIAGNOSIS. (Based on Kühn's description and figure.) ?*Lagynopora* with encrusting zoarium; zooecia of large size ($Lz = 0{\cdot}90$ to $1{\cdot}00$ mm., $lz = 0{\cdot}56$ to $0{\cdot}58$ mm.); orifice very large ($hr = 0{\cdot}28$ mm., $lr = 0{\cdot}25$ to $0{\cdot}28$ mm.), rounded distally and laterally with a straight proximal margin; oral spines 6; frontal shield fairly well arched; costae 8–11, separated for half their length by wide intercostal spaces; median area of fusion extends about half-way along the costae; apertural bar fairly wide, straight; avicularia rare—their form unknown.

REMARKS. Kühn (1930 : 65) has commented "... our genus [*Canupora*] has great similarity to *Prodromopora* [= *Lagynopora*], but is distinguished by the presence of 6 oral spines, which is a more primitive condition than the 4 oral spines of *Prodromopora*, so that one may regard it as the most primitive genus of the Lagynoporidae. Always assuming that Lang's developmental series is correct" [translation].

It is difficult to interpret Kühn's figure, and in his description he did not state whether or not his material possessed proximal oral shields or large distal communication pores. The occurrence of 6 oral spines in the adult zooecia by no means justifies placing the species in a separate genus, and it is therefore provisionally referred to *Lagynopora*, especially as this, in its young stages, often has 6 oral spines. Also, if the age of the Lithothamnian Chalk is correctly established as Danian, it is very unlikely that *Lagynopora*? *prima* may be regarded as the most primitive unit in a developmental series of species which are otherwise confined to the Senonian.

No specimens of *Lagynopora*? *prima* are available in the collections of the British Museum (Natural History).

Genus *HEXACANTHOPORA* Lang

1916*a* *Hexacanthopora* Lang, p. 394.
1921 *Hexacanthopora* : Lang, pp. lxi, xcii, xciii, 44
1924 *Hexacanthopora* : Gaster, second of three tables opp. p. 110.
1927 *Hexacanthopora* : Canu & Bassler, pp. 17, 31.
1930 *Hexacanthopora* : Gaster, table opp. p. 340.
1930 *Hexacanthopora* : Voigt, pp. 495, 542, 560.
1935 *Hexacanthopora* : Bassler, pp. 30, 126.
1949 *Hexacanthopora* : Voigt, pp. 37, 43.
1953 *Hexacanthopora* : Bassler, p. G188.

TYPE SPECIES. *Hexacanthopora sexspinosa* Lang, 1916*a* : 394. Senonian, zone of *Marsupites* [subzone unspecified]. Roke Farm, south-east of Odiham, Hampshire. L. Treacher Collection.

EMENDED DIAGNOSIS. Pelmatoporinae with a small median process on the apertural bar—this process never fuses with the proximal-lateral oral spines to form a proximal oral shield ; oral spines often 6 ; avicularia large, prominent, often frequent.

REMARKS. *Hexacanthopora* Lang is distinguished from *Lagynopora* Lang primarily by the absence of a proximal oral shield, and from *Leptocheilopora* by the presence of a median process on the apertural bar and numerous, large avicularia.

Hexacanthopora is better distinguished on these characters than solely on the number of oral spines in its adult zooecia. Lang (1921 : 44) diagnosed *Hexacanthopora* as " Lagynoporinae [= Pelmatoporinae] in which (at least) some epheboecia [adult zooecia] have six apertural [oral] spines ". The genus may possess from 4 to 6 oral spines in the adult zooecia : in two specimens, D.28890 and D.29897, it was not possible to find any zooecium with 6 oral spines. It is, however, still true to say that the 4–6 (commonly 6) oral spines which occur in *Hexacanthopora* contrast with the number of oral spines in *Leptocheilopora*. In the latter there are 4 oral spines only, except in *L. vulnerata* (Brydone) where there are consistently 7 in each adult zooecium. *Lagynopora*, with the exception of *L. ollula* (4, occasionally 5) and *L.? prima* (6), consistently has only 4 oral spines in the adult zooecia.

Only two species of *Hexacanthopora* are recognized here : *H. sexspinosa* Lang, and *H. viginticostata* Voigt from the Senonian of Germany.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *M. cor-anguinum* to *B. mucronata*.

KEY TO THE SPECIES OF *HEXACANTHOPORA* LANG

(I) *Hexacanthopora* with 4–6 oral spines in the adult zooecia ; costae 11–18 ; median area of fusion not obscuring the inner ends of the costae ; avicularia usually frequent, of two sizes in the adult zoarial stage . . 1. ***H. sexspinosa*** Lang

(II) *Hexacanthopora* always with 6 oral spines in the adult zooecia ; costae 20 (average) ; median area of fusion wide, obscuring the outlines of the inner ends of the costae ; avicularia large, frequent, apparently of one size in the adult zoarial stage 2. ***H. viginticostata*** Voigt

1. *Hexacanthopora sexspinosa* Lang

(Pl. 5, figs. 3–6)

1916*a* *Hexacanthopora sexspinosa* Lang, p. 394.
1916*a* *Hexacanthopora kintburiensis* Lang, p. 394.
1916*a* *Hexacanthopora brightonensis* Lang, p. 394.
1921 *Hexacanthopora sexspinosa* Lang : Lang, lxi, xciii, 45, pl. 1, fig. 8 ; text-fig. 21.
1921 *Hexacanthopora kintburiensis* Lang : Lang, pp. xcii, 47, pl. 1, fig. 9 ; text-fig. 23.
1921 *Hexacanthopora brightonensis* Lang : Lang, pp. xcii, 48, text-fig. 24.
1924 *Hexacanthopora kintburiensis* Lang : Gaster, second of three tables opp. p. 110.
1924 *Hexacanthopora brightonensis* Lang : Gaster, second of three tables opp. p. 110.
1930 *Hexacanthopora sexspinosa* Lang : Gaster, table opp. p. 340.
1930 *Hexacanthopora kintburiensis* Lang : Gaster, table opp. p. 340.
1930 *Hexacanthopora brightonensis* Lang : Gaster, table opp. p. 340.
1930 *Hexacanthopora brightonensis* Lang : Voigt, pp. 495, 542, 560, pl. 26, fig. 11.
1930 *Hexacanthopora sexspinosa* Lang : Voigt, pp. 495, 542, 560, pl. 26, fig. 12.
1935 *Hexacanthopora sexspinosa* Lang : Bassler, p. 126.
1953 *Hexacanthopora sexspinosa* Lang : Bassler, p. G188, text-fig. 142, 11.

HOLOTYPE. D. 21205. Well-preserved zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites* [subzone unspecified]. Roke Farm, south-east of Odiham, Hampshire. L. Treacher Collection.

EMENDED DIAGNOSIS. *Hexacanthopora* with fairly large zooecia (Lz = 0·55 to 0·80 mm., lz = 0·35 to 0·55 mm.) ; orifice wider than high (hr = 0·10 to 0·15 mm., lr = 0·12 to 0·17 mm.) ; oral spines 4–6 ; costae 11–18, visible individually up to the mid-line of the frontal shield ; avicularia of two kinds, (*a*) a large oval form with small constrictions near the proximal end, (*b*) a smaller, similar form ; one type of avicularium may be more numerous than the other, or occasionally only one type may be visible, or, even more rarely, both types may be absent.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* fairly large, broadly oval, each with a very prominent, large *distal communication pore* and smaller *lateral communication pores*. *Dietellae* are present in the distal-basal angles of the zooecia, and revealed where the upper part of the zooecium has been worn away (see Pl. 5, fig. 3). *Orifice* wider than high, evenly rounded distally and laterally, with a straight or slightly proximally curved proximal margin. *Oral spines* 4–6, symmetrically arranged round the orifice. In a given zoarium 6 oral spines may be consistently developed in the adult zooecia, as in the holotype ; or 4, 5 or 6 oral spines may be present, as in D.21204 ; or only 4 or 5 may occur, as in D.28890. The proximal-lateral pair of oral spines are not enlarged as in *Lagynopora*. *Frontal shield* well arched, but flattened in the median area. *Costae* 11–18, separated for two-thirds of their length by distinct *intercostal spaces*. The costae are united in the median area of fusion, but the outlines of the inner ends of individual costae remain clearly visible. *Median area of fusion narrow*. Minute *pelmatidia* may be seen along the mid-line of well-preserved costae. *Apertural bar* straight, somewhat curved distally at either end, with a distinct, though fairly small, *median process* which apparently never fuses with the proximal-lateral pair of oral spines to form a proximal oral shield as in *Lagynopora*. *Gymnocyst* smooth, often slightly exposed proximally to the frontal shields and occasionally very narrowly along their sides. No *interzooecial secondary tissue.*

Avicularia of two sizes. Type *a* is a larger form, usually a little longer than wide, but occasionally the reverse is true, oval, or slightly pointed distally, with slight proximal-lateral constrictions dividing a small proximal portion from a very large *rostrum.* Tybe *b* is smaller, but otherwise similar to type *a.* Both types may be variously orientated—usually distally and very obliquely outwards from the zooecia they appear to accompany. None has been seen which is directed proximally. In a given zoarium one type or the other may be more frequent, or, very occasionally, only one form may be developed. Even more rarely avicularia may be absent.

Ovicells hyperstomial, prominent, globular, smooth, without *lateral slits.* A pair of oral spines may flank the entrance to the ovicell.

Young zooecia smaller than the adult zooecia, often with fewer (11–14) costae and more numerous oral spines. Paratype D.21207 possesses at least one young zooecium with 7 oral spines, and D.29895–96 have young zooecia with 6 oral spines and adult zooecia with only 4 or 5.

Ancestrula small, often obscured. In the holotype the ancestrula has 6 oral spines and 12 costae.

MEASUREMENTS. *Adult zooecia* Lz = 0·50 to 0·80 mm.
lz = 0·35 to 0·55 mm.
hr = 0·10 to 0·15 mm.
lr = 0·10 to 0·15 mm.
Young zooecia Lz = 0·35 to 0·50 mm.
lz = 0·21 to 0·33 mm.
hr = 0·09 to 0·11 mm.
lr = 0·10 to 0·13 mm.
Ancestrula (of holotype) Lz = 0·32 mm.
lz = 0·20 mm.
hr = 0·07 mm.
lr = 0·09 mm.
Avicularia :
Type *a* Lz = 0·18 to 0·25 mm.
lz = 0·17 to 0·25 mm.
Type *b* Lz = 0·11 to 0·18 mm.
lz = 0·09 to 0·16 mm.

REMARKS. *Hexacanthopora kintburiensis* Lang and *H. brightonensis* Lang are synonymous with *H. sexspinosa* Lang. Thirteen specimens of *Hexacanthopora,* labelled by Lang, are available in the collections of the British Museum (Natural History). Lang (1916*a* : 394 ; 1921 : 45) divided this material into three species, basing his division principally on the number of oral spines commonly developed in adult zooecia, the number of costae, and the length of zooecia. In addition he stated (1921 : 45) that *H. sexspinosa* possesses two types of avicularia, " both kinds, especially the smaller, rare ; the smaller kind are more or less distally directed, and have somewhat elongate, slightly constricted, and blunt apertures ; the larger vary in size, but generally have an aperture as large as that of an orthoecium [zooecium]; they are indifferently oriented, little longer than wide, with a somewhat pointed aperture divided by a constriction into a very much smaller proximal portion and a very large rostrum ". The avicularia of *H. kintburiensis* he described (1921 : 47) as " occasional, large, with blunt, indifferently orientated apertures, nearly as large as those of the orthoecia, with a very slight constriction near the proximal end ". *H. brightonensis* (1921 : 49) was stated to have " aviculoecia [avicularia] apparently of the large kind only, fairly numerous, irregularly distributed, with indifferently orientated, nearly circular apertures of nearly the same size as those of the orthoecia, and with a constriction nearer the proximal end ".

Both large and small avicularia are, however, present in each of the species of *Hexacanthopora* recognized by Lang. The larger avicularia (type *a*) may be more numerous than the smaller (type *b*). Normally avicularia occur infrequently, but in one specimen, D.28890—the holotype of *H. brightonensis* Lang, they are unusually abundant.

The type and variable number of the avicularia do not aid specific division of the material. Similarly the dimensions of the zooecia and the ranges in number of costae, in the three species recognized by Lang, overlap or are the same. These characters and the number of oral spines are compared in the following table :

Specimens labelled by Lang as :	Measurements in mm.			
	Lz	lz	hr	lr
1. *H. sexspinosa*	0·55 to 0·75	0·35 to 0·45	0·10 to 0·13	0·12 to 0·15
2. *H. kintburiensis*	0·55 to 0·80	0·35 to 0·55	0·10 to 0·15	0·14 to 0·17
3. *H. brightonensis*	0·60 to 0·67	0·38 to 0·50	0·13	0·15

	Costae	Oral spines
1.	14–18	6
2.	11–18	4–6
3.	11–13	4–5

All adult zooecia in the specimens labelled by Lang " *H. sexspinosa* " possess 6 oral spines, whereas those labelled " *H. kintburiensis* " have adult zooecia with 4 or 5 as well as commonly 6 oral spines. In the specimens labelled " *H. brightonensis* " a few adult zooecia have only 4 oral spines, the majority have 5 and no definite instance could be found of 6 oral spines.

Voigt (1930 : 495) has recorded specimens of *H. sexspinosa* and *H. brightonensis* from Germany in chalk equivalent in age to the zones of *Marsupites* and *G.* [*A.*] *quadrata*.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *Marsupites* to *G.* [*A.*] *quadrata* in England and Germany, ?also zone of *M. cor-anguinum* in England.

SPECIMENS. D.21205. Holotype—see above.

D.21206. Paratype. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for the holotype.

D.21207. Paratype. Zoarial fragment with ancestrula and mainly young zooecia encrusting a piece of echinoid test. Senonian, ? zone of *Marsupites*. Odiham, Hampshire. L. Treacher Collection.

D.29101. Paratype. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites* [subzone unspecified]. Margate, Kent. N. T. Wetherell Collection.

D.21204. Holotype of *Hexacanthopora kintburiensis* Lang. Well-preserved zoarium, two zooecia with undamaged ovicells, encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata.* Kintbury, west of Newbury, Berkshire. L. Treacher Collection.

D.29070. Paratype of *H. kintburiensis* Lang. Zoarial fragment encrusting a piece of ?*Inoceramus* shell. Senonian, zone of *G.* [*A.*] *quadrata.* Romsey, south-west of Winchester, Hampshire. H. P. Blackmore Collection.

?D.2631 Paratype of *H. kintburiensis* Lang. A worn zoarial fragment with about ten abraded zooecia only. It is not possible to distinguish the number of oral spines, nor to measure the orifice. Encrusting a shell fragment. Upper Senonian [zone unspecified]. Salisbury, Wiltshire. G. R. Vine Collection.

D.28890. Holotype of *H. brightonensis* Lang. Zoarial fragment with fairly well preserved zooecia and very frequent avicularia, encrusting a piece of echinoid test. Senonian, zone of *Marsupites* [subzone unspecified]. Brighton, Sussex.

?D.28891. Paratype of *H. brightonensis* Lang. A very small, worn zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula,* subzone of *E. scutata* var. *depressula.* Cliffs between last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.

D.29896. Zoarial fragment with ancestrula and numerous adult ovicelled zooecia. Encrusting a piece of echinoid test. Labelled by Lang "*H. kintburiensis*". Senonian, zone and subzone of *Marsupites.* West pit, in top of Hangleton Lane, Highdown Hill, east of Angmering, Sussex.

?D.21208. Very poorly preserved zoarial fragment encrusting a piece of echinoid test. Labelled by Lang "*H. kintburiensis*". Senonian, zone of *M. coranguinum.* Woburn Green, south-west of Beaconsfield, Buckinghamshire. L. Treacher Collection.

D.29895. Zoarial fragment with ancestrula and adult ovicelled zooecia encrusting a piece of echinoid test. Labelled by Lang "*H. kintburiensis*". Senonian, zone of *O. pilula,* subzone of *E. scutata* var. *depressula.* Pit 4 of Gaster, western pit, south of Lancing Ring and west of Boundstone Lane, north-east of Worthing, Sussex.

D.29897. Worn zoarial fragment with obscured ancestrula. Encrusting a piece of echinoid test. Senonian, zone of *O. pilula,* subzone of *E. scutata* var. *depressula.* Cliffs, west of Chimney Shaft, Roedean, east of Brighton, Sussex.

D.41065. Very small zoarial fragment with young zooecia encrusting a piece of echinoid test. Senonian, zone and subzone of *Marsupites.* Section 75 of Gaster, Brighton Cliffs, Sussex.

2. *Hexacanthopora viginticostata* Voigt

1930 *Hexacanthopora viginticostata* Voigt, pp. 495, 560, pl. 26, fig. 13.
1949 *Hexacanthopora viginticostata* Voigt : Voigt, pp. 37, 43, pl. 9, figs. 1, 2.

Neotype. (Designated by Voigt, 1949 : 37, pl. 9, fig. 2.) Sample 142. Senonian, zone of *G.* [*A.*] *quadrata*. Lägerdorf, Holstein, Germany.

Diagnosis. (Based on Voigt's descriptions and figures, 1930 ; 1949.) *Hexacanthopora* with zooecia of fairly large size (Lz = 0·60 to 0·70 mm., lz = about 0·40 mm.) ; orifice small, higher than wide (hr = about 0·12 mm.) ; oral spines constantly 6 ; costae 20 (average) ; median area of fusion wide, obscuring the outlines of the inner two-thirds of the costae ; avicularia variable in occurrence, large (0·20 to 0·30 mm. diameter), with a slit-like or irregularly polygonal aperture, smaller in the young than in the adult stages of zoarial growth ; ovicells hyperstomial, large, globular ; young zooecia with fewer (about 14–16) costae ; ancestrula about 0·30 mm. long.

Remarks. Voigt recently (1949 : 37) designated a neotype for the species, the holotype having been lost in a fire. The well-preserved new material, from the same horizon and the same and different localities as the original specimens, differs mainly in the number of avicularia present. They are apparently much more numerous than in the original material in which they were " rare ".

H. viginticostata Voigt is distinguished from *H. sexspinosa* Lang by its larger number of costae and its wide median area of fusion. Also the orifice of the former is apparently higher than wide, contrasting with that of *H. sexspinosa* which is consistently wider than high.

Stratigraphical distribution. Senonian, zones of *G.* [*A.*] *quadrata* and *B. mucronata* in Germany.

Specimens. None in the collections.

Genus ***LEPTOCHEILOPORA*** Lang

?1852 *Reptoporella* d'Orbigny, pl. 717, figs. 6, 7 ; 1853, p. 474 ; 1854, p. 1097.
?1857 *Repteporella* [*sic*] : Pictet, p. 111.
1900*b* *Cribrilina* (*Typica*) : Canu, p. 447 [*partim*].
1900*b* *non Membranipora* : Canu, p. 449.
1910 *Cribrilina* : Brydone, p. 391 [*partim*].
1912 *Cribrilina* : White, p. 43 [*partim*].
1914*b* *Cribrilina* : Brydone, p. 97 [*partim*].
1916*a* *Leptocheilopora* Lang, p. 396.
1917 *Cribrilina* : Brydone, p. 51 [*partim*—*C. tumuliformis* only].
?1917 *Reptoporella* : Lang, p. 172.
1917 *Leptocheilopora* : Lang, p. 172.
1921 *Leptocheilopora* : Lang, pp. lxii, lxv, 68.
?1921 *Reptoporella* : Lang, p. lxv.
1925 *Barroisina* : Levinsen, p. 384 [*partim*—*B. galeata* only].
1930 *Leptocheilopora* : Voigt, p. 496.
1930 *Leptocheilopora* : Gaster, table opp. p. 340.
1935 *Leptocheilopra* : Bassler, pp. 30, 138.
1943 *Leptocheilopora* : Gillard, pp. 186, 187 [probably *partim*—? *non L. regularis* (d'Orbigny)].
1953 *Leptocheilopora* : Bassler, p. G188.

TYPE SPECIES. *Leptocheilopora tenuilabrosa* Lang, 1916*a* : 397. Senonian, zone of *Marsupites testudinarius* (subzone unspecified). Brighton, Sussex.

EMENDED DIAGNOSIS. Pelmatoporinae with no median process on the apertural bar; no proximal oral shield; avicularia absent or only very occasionally developed.

REMARKS. Lang (1921 : 71), followed by Bassler (1953 : G188), suggested that *Reptoporella* d'Orbigny may be a synonym of *Leptocheilopora* Lang, but this cannot be decided until d'Orbigny's material of *Reptoporella regularis* d'Orbigny (1853 : 475, pl. 717, figs. 6, 7), the type species of *Reptoporella*, has been restudied. (See discussion of species of uncertain systematic position p. 118.)

Leptocheilopora as here defined is placed in the subfamily Pelmatoporinae. Four species are recognized: *L. magna* Lang and *L. vulnerata* (Brydone) are, as yet, known only from the Lower Maastrichtian or highest Senonian, *L. filliozati* (Brydone) occurs throughout the zones of *G.* [*A.*] *quadrata* and *O. pilula*, and *L. tenuilabrosa* Lang is recorded from the zones of *Marsupites* and *Micraster cor-anguinum*.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum* to zone of *B. mucronata*, and Maastrichtian, zone of *Ostrea lunata* in England, zone of *Belemnella lanceolata* mut. *sumensis* or zone of *B. junior* in Germany.

KEY TO THE SPECIES OF *LEPTOCHEILOPORA*

(I) *Leptocheilopora* with zooecia of large size (Lz = 0·80 to 1·00 mm., lz = 0·50 to 0·66 mm.); orifice of constant form with marked proximal-lateral constrictions 1. **L. magna** Lang

(II) *Leptocheilopora* with zooecia of small size (Lz = less than 0·75 mm.).

  (A) Primary ovicells costate; costae 17–22, (11–15 in young zooecia); intercostal spaces distinct; apertural bar broadly triangular, usually somewhat curved proximally; large, vicarious, spatulate avicularia may occur in the adult stages of some zoaria . . . . 2. **L. tenuilabrosa** Lang

  (B) Primary ovicells smooth.

    (1) Lz = 0·55 to 0·75 mm., lz = 0·35 to 0·55 mm.; orifice of variable shape, but often with proximal-lateral constrictions (hr = 0·12 to 0·17 mm., lr = 0·12 to 0·16 mm.); oral spines 4; costae 12–18, well defined; often with a median ridge on the frontal shield; ovicells without lateral slits . . . . 3. **L. filliozati** (Brydone)

    (2) Lz = 0·42 to 0·50 mm., lz = 0·31 to 0·35 mm.; orifice small, of fairly constant size (hr = 0·10 to 0·12 mm., lr = 0·09 to 0·11 mm.); usually lacking proximal-lateral constrictions; oral spines 7; frontal shield flatly arched without a median ridge; costae 11–15, flat, closely placed; ovicell with small lateral slits . . . . 4. **L. vulnerata** (Brydone)

Reptoporella regularis d'Orbigny, previously assigned to *Leptocheilopora*, and *Leptocheilopora longuessensis* Lang are of uncertain systematic position and are omitted from the above key.

1. *Leptocheilopora magna* Lang

(Pl. 6, figs. 1–7; Text-figs. 42, 43)

1916*a* *Leptocheilopora magna* Lang, p. 396.
?1916*a* *Leptocheilopora languessensis* [*sic*] Lang, pp. 396, 397.

?1921 *Leptocheilopora longuessensis* Lang: Lang, p. 76.
1921 *Leptocheilopora magna* Lang: Lang, p. 77, pl. 2, fig. 9; text-fig. 38.
1925 *Barroisina galeata* Levinsen, p. 385, pl. 6, fig. 67.
1930 *Leptocheilopora magna* Lang: Voigt, p. 496 [*partim*], pl. 27, fig. 5 [probably *non* fig. 4].

HOLOTYPE. D.19623. Incomplete zoarium with ancestrula, encrusting a fragment of echinoid test. Horizon given as " Senonian, Campanian, zone of *B. mucronata*, Lüneburg, Hanover ", Germany—very probably Maastrichtian. F. Möckler Collection.



FIG. 42. *Leptocheilopora magna* Lang, D.38919. Adult zooecium with widely exposed gymnocyst and marked proximal-lateral constrictions in the orifice.

EMENDED DIAGNOSIS. *Leptocheilopora* with zooecia of large size (Lz = 0·80 to 1·00 mm.); costae 12–17; orifice with marked proximal-lateral constrictions.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* large, broadly oval, each with a single, fairly large, *distal communication pore*. *Orifice* higher than wide, rounded distally and laterally with marked proximal-lateral constrictions, the proximal margin often well curved proximally. *Oral spines* 4, small, unmodified. *Frontal shield* wide and flat. *Costae* 12–17, flat, tapering inwards where their distal ends may be traced across the wide *median area of fusion*. No *lateral costal fusions*. Very small *pelmata*, usually obscured by wear, along the mid-line of each costa. *Aperturral bar* usually a little wider than the most distal costae, flat, smooth and curved proximally, with no median process. *Gymnocyst* often almost obscured by the close placing of the zooecia, but usually visible proximally to the frontal shield, and occasionally widely exposed all round the frontal shield (Text-fig. 42). No *interzooecial secondary tissue.*

Avicularia absent. *Primary ovicells* large, prominent, hyperstomial, smooth, globular, apparently composed of a smooth endooecium and a costate ectooecium which is usually incomplete and marginal. *Young zooecia* small, with only 10 costae. The orifice lacks the proximal-lateral constrictions characteristic of the orifices of adult zooecia, and is less markedly higher than wide. Oral spines 4, more prominent than in the adult zooecia. Frontal shield flatly arched, with only a very small amount of gymnocyst exposed proximally to it.



Fig. 43. *Leptocheilopora magna* Lang, D.38918. Orifice and broken ovicell resting on the proximal part of frontal shield of succeeding zooecium. The specimen is partly obscured by matrix.

Measurements. *Adult zooecia* Lz = 0·80 to 1·00 mm.
lz = 0·50 to 0·66 mm.
hr = 0·21 to 0·24 mm.
lr = 0·18 to 0·21 mm.
Young zooecia Lz = 0·50 mm.
lz = 0·27 mm.
hr = 0·15 mm.
lr = 0·12 mm.

REMARKS. The holotype, D.19623, and paratype, D.19622, have been re-examined. The former has the ancestrula imperfectly preserved, and outside the first ring of zooecia the range in number of costae, 12–17, is the maximum yet recorded for *L. magna*. In this specimen the measurements are : Lz = 0·90 to 1·00 mm., lz = 0·56 to 0·66 mm., hr = 0·21 mm., lr = 0·18 mm. The dimensions of the orifice are very constant. All zooecia of the holotype are apparently non-ovicelled.

Paratype D.19622 is a much damaged and worn zoarium with only the ancestrula and surrounding rings of young zooecia preserved ; the majority of the zooecia are thus small.

Two specimens, D.38918 and D.38919, labelled by Voigt as *L. magna* Lang, have also been re-examined. The zooecia of the former are slightly larger and possess a few more costae than those of the latter. Imperfectly preserved ovicells are present in both specimens.

Voigt (1930 : 496, pl. 27, figs. 4, 5) assigned two specimens to *L. magna* Lang. I accept the specimen shown in fig. 5 as belonging to the species, but that shown in fig. 4 has a very marked series of short, slot-like intercostal spaces, a feature which I have not found in the specimens of *L. magna* available to me. I have therefore tentatively excluded that specimen from the synonymy of the species.

In agreement with Voigt (1930 : 496), *Barroisina galeata* Levinsen is regarded as synonymous with *Leptocheilopora magna* Lang. Levinsen (1925 : 385, pl. 6, fig. 67) referred to the " rib area [frontal shield] . . . of 7–9 pairs of flat ribs ", to its " high, rounded, strongly arched " ovicells, and to the absence of avicularia—all characteristic of *L. magna*.

The imperfectly preserved ovicells which occur in D.38918 and D.38919 are apparently ribbed or costate marginally, and seem to consist of two layers interpreted as a smooth endooecium partly covered by a costate ectooecium. The primary ovicell of *L. magna* Lang is smooth, but this primary layer (the endooecium) may be partly covered secondarily by an outer layer (the costate ectooecium). This feature could lead to confusion with *L. tenuilabrosa* Lang, in which the primary ovicell is costate, consisting of one layer only. However, the very large size of *L. magna* Lang and its very distinctively shaped orifice (pliophloean)[1] serve to distinguish it clearly from other species of *Leptocheilopora*.

Leptocheilopora longuessensis Lang is provisionally included in the synonymy of *L. magna* Lang. The diagnoses given by Lang (1916*a* : 396 ; 1921 : 76) are based only on the interpretation of a poor photograph of the holotype in the Canu Collection. From the photograph it is possible to determine only that the zoarium is that of an encrusting, unilaminar *Leptocheilopora* ; the zooecia are broadly oval with flatly arched frontal shields possessing fairly numerous (probable about 16) costae ; the orifice is longitudinally elongate and distally and laterally rounded with proximal-lateral constrictions and 4 somewhat enlarged oral spines. It is very probable that the zooecia shown in the photograph are close to the ancestrula, as those near the base of the zoarial fragment are much smaller than those near the outer edges. This being so it is very uncertain whether the smaller size of the zooecia can have much diagnostic significance—Lang (1921 : 76) gave Lz as about 0·60 mm. and lz as 0·48 mm. The specimen could well be part of the young stage of zoarium of *Leptocheilopora magna* Lang. However, this opinion can only be tentative.

---

[1] See Lang, 1921 : xlvii, fig. 109.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian, either zone of *Belemnella lanceolata* mut. *sumensis* or zone of *B. junior*, and Senonian, zone of *Belemnitella mucronata*.

SPECIMENS. D.19623. Holotype—see above.

D.19622. Paratype. Damaged and worn zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for the holotype.

D.38918. Zoarial fragment encrusting a piece of siliceous sponge. Maastrichtian, either zone *Belemnella lanceolata* mut. *sumensis* or zone of *B. junior*. Cement factory (loose on floor), Hemmoor, Hanover, Germany. E. Voigt Collection.

D.38919. Zoarial fragment encrusting a belemnite guard. Horizon, locality and collection as for D.38918.

D.39431. Two damaged zooecia only, encrusting a fragment of echinoid test. Senonian, zone of *Belemnitella mucronata*. Felpham Chalk [foreshore], Bognor, Sussex.

2. ***Leptocheilopora tenuilabrosa*** Lang

(Pl. 6, figs. 8–11 ; Pl. 7, figs. 1, 2 ; Text-figs. 44–48)

1916*a* *Leptocheilopora tenuilabrosa* Lang. pp. 396 397.

1917 *Leptocheilopora tenuilabrosa* Lang : Lang p. 172.

1917 *Cribrilina tumuliformis* Brydone, p. 51, pl. 3, figs. 7, 8.

1921 *Leptocheilopora tenuilabrosa* Lang : Lang, pp. lxii, xciii, 73, pl. 2, fig. 7 ; text-fig. 35.

1921 *Leptocheilopora gasteri* Lang, pp. xciv, 71, pl. 2, fig. 6 ; text-fig. 34.

1930 *Leptocheilopora tenuilabrosa* Lang : Gaster, table opp. p. 340.

1935 *Leptocheilopora tenuilabrosa* Lang : Bassler, p. 138.

1953 *Leptocheilopora tenuilabrosa* Lang : Bassler, p. G188, text-fig. 142, 10.

HOLOTYPE. D.28892. Zoarial fragment, probably with fairly young zooecia, encrusting a piece of echinoid test. Senonian, zone of *Marsupites testudinarius* (subzone unspecified). Brighton, Sussex.

EMENDED DIAGNOSIS. *Leptocheilopora* with costate primary ovicell ; costae 17–22, (11–15 in younger zooecia) ; intercostal spaces distinct ; large, spatulate, vicarious avicularia may occur in the adult stages of some zoaria.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, with a single, oval *distal communication pore*. *Orifice* small, of variable shape ; always rounded distally and usually rounded laterally, but occasionally with straight sides, and often with slight, but distinct, proximal-lateral constrictions ; proximal margin straight or slightly curved proximally. A distinct median notch, probably produced by wear, may be present in the proximal margin of the orifice. *Oral spines* 4, very small, indistinct, grouped on the distal margin of the orifice. *Frontal shield* flatly arched, the costae curving down gently from the mid-line to meet the gymnocyst. *Costae* 17–22, flat, tapering inwards, and separated by narrow, but distinct, *intercostal* *spaces.* No *lateral costal fusions.* Very small *pelmata* occur along the mid-line of the upper surface of each costa. *Apertural bar* straight, or slightly curved proximally when it is very broadly triangular (the apex of the triangle directed proximally), flat, with no *median process*, and only slightly wider than the distal costae ; a very slight median ridge may be present and this may be prolonged on to the frontal shield as an irregular median ridge. *Gymnocyst* smooth, usually well exposed, except where the close placing of zooecia may obscure it. No *interzooecial secondary tissue.*



FIG. 44. *Leptocheilopora tenuilabrosa* Lang, D.28892. Holotype. Adult zooecium with broken, worn, costate ovicell ; the apertural bar is slightly damaged.

Avicularia very large, occasional, sporadic, vicarious and spatulate with costate frontal shields covering the proximal part of the aperture (about 8–13 costae are present). The distal end of the spatulate *rostrum* may include a narrow calcareous shelf (Text-fig. 47), but, in some cases, this may be absent (Text-fig. 48). Avicularia have been seen only in the adult stages of two zoaria : they apparently do not occur in the younger stages of zoarial growth and may be absent even in the adult zones of a zoarium. *Primary ovicells* prominent, frequent, hyperstomial, costate, tending to be longer than wide. The 11–13 flat, hollow costae which cover the ovicell apparently possess occasional minute pores along their mid-lines : the costae may form an *ectooecium*, but this is uncertain. *Young zooecia* smaller than the adult zooecia, with fewer (11–15) costae, and apparently not associated with vicarious avicularia.



FIG. 45. *Leptocheilopora tenuilabrosa* Lang, D. 28892. Holotype. Orifice and complete costate ovicell of adult zooecium ; the apertural bar is broken in the middle.

MEASUREMENTS. *Adult zooecia* Lz = 0·57 to 0·66 mm.
lz = 0·32 to 0·50 mm.
hr = 0·12 to 0·13 mm.
lr = 0·13 t9 0·14 mm.
Avicularia Lz = 0·80 to 0·83 mm.
lz = 0·33 to 0·36 mm.
Young zooecia Lz = 0·43 to 0·52 mm.
lz = 0·30 to 0·45 mm.
hr = 0·08 to 0·10 mm.
lr = 0·10 to 0·12 mm.

REMARKS. *Leptocheilopora gasteri* Lang is synonymous with *L. tenuilabrosa* Lang. The only apparent difference between the two species, which may be established on the specimens available in the collections of the British Museum (Natural History), is in the range in number of costae : 16–18 for the former and 11–15 for the latter. The dimensions of the zooecia, and particularly of the orifices, are very similar and both species have costate ovicells. Lang (1921 : 71, 72) mentioned and figured what he regarded as "rather obvious, irregular lateral costal fusions" in *L. gasteri*. The available specimens labelled by him as *L. gasteri* are worn, but the costae are clearly separated by narrow slit-like intercostal spaces (Text-fig. 46). Uneven wear, possibly combined with a certain amount of secondary calcification during fossilization, has apparently produced small hollows and slight, irregular accumulations of calcium-carbonate in the narrow spaces between the costae. These might easily be misinterpreted as incipient lateral costal fusions The same feature occurs occasionally in the specimens which he assigned to *L tenuilabrosa*



FIG. 46. *Leptocheilopora tenuilabrosa* Lang, D.29901. Holotype of *L. gasteri* Lang. Adult zooecium with costate ovicell, with four minute oral spine-bases grouped on distal margin of orifice.

Lang (1921 : 73) regarded *Cribrilina tumuliformis* Brydone (1917 : 51, pl. 3, figs. 7, 8) as synonymous with *Leptocheilopora tenuilabrosa*. Examination of Brydone's figured specimens of *Cribrilina tumuliformis* confirms this opinion. The range in number of costae in Brydone's specimens, S.M., B.36338–39, is, however, 17–22, and in both zoaria large, spatulate, vicarious avicularia are present. Young zooecia are also preserved and these are more closely similar to the zooecia of the holotype of *L. tenuilabrosa* (D.28892) and to those of the paratype (D.21210) than are the adult zooecia. It seems probably that Lang's specimens lack the fully developed adult stages of zoarial growth. In S.M., B.36339 a complete but small normal zooecium is regenerated inside a larger, damaged individual.

The very unusual vicarious, costate avicularia which may be present are particularly diagnostic of *L. tenuilabrosa*. Even where avicularia are absent, as they may be in some zoaria, the costate primary ovicell clearly distinguishes the species. Dr. Hastings kindly sent me the following note : " Vicarious avicularia with a costulate frontal shield are found in some recent species of *Figularia*. I have seen them in the type material of *F. philomela* Busk sp., where they were also observed by Harmer (1902 : 239) ; and in the specimen of *F. spatulata* (Calvet) collected by the *Discovery* Expedition (Stat. 47, 1 mile west of Shag Rocks, South Georgia, 199 m.)." Osburn (1950, p. 188, pl. 29, fig. 1) described and figured a new Recent species, *Colletosia bellula*, in which " the primary ovicell is small ... smooth ... ; it soon becomes covered by an ectooecium which is composed of three or four pairs of radiating costules similar to those of the front and with similar lacunae [intercostal spaces]. ... "



Figs. 47, 48. *Leptocheilopora tenuilabrosa* Lang.

(47) S.M., B.36339. Figured Brydone (1917, pl. 3, fig. 8) as *Cribrilina tumuliformis*. Adult zooecium with costate ovicell (the ovicell of the preceding zooecium is also shown covering the proximal end of the zooecium) and a large, vicarious avicularium with costate frontal shield.

(48) S.M., B.36338. Figured Brydone (1917, pl. 3, fig. 7) as *Cribrilina tumuliformis*. Vicarious avicularium with typically spatulate rostrum and costate frontal shield. Small spine-bases may occur near the distal end of the aperture.

Stratigraphical distribution. Senonian, zones of *M. cor-anguinum* and *Marsupites*.

Specimens. D.28892. Holotype—see above.

D.21210. Paratype. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites* [subzone unspecified]. Roke Farm, south-east of Odiham, Hampshire. L. Treacher Collection.

D.29901. Holotype of *L. gasteri* Lang. Zoarial fragment encrusting a piece of *Inoceramus* shell. Senonian, zone of *M. cor-anguinum*. Pit in Houndean Bottom, west of Lewes, Sussex.

D.29902. Paratype of *L. gasteri* Lang. Zoarial fragment encrusting a piece of *Inoceramus* shell. Horizon, locality and collection as for D.29901.

D.40931. Another fragment of D.29902.

OTHER MATERIAL

S.M., B.36338. Large, well-preserved zoarial fragment, encrusting a piece of echinoid test. Figured by Brydone (1917, pl. 3, fig. 7) as *Cribrilina tumuliformis*. Senonian, zone of *M. cor-anguinum*. Leaves Green, Kent. R. M. Brydone Collection.

S.M., B.36339. Large, well-preserved zoarial fragment encrusting a piece of echinoid test. Figured by Brydone (1917, pl. 3, fig. 8) as *Cribrilina tumuliformis*. Senonian, horizon and collection as for S.M., B.36338. Gravesend, Kent.

3. *Leptocheilopora filliozati* (Brydone)

(Pl. 7, figs. 3, 4; Pl. 8; Text-figs. 49, 50)

1910 *Cribrilina filliozati* Brydone, p. 391, pl. 30, figs. 9, 10.
1912 *Cribrilina filliozati* Brydone: White, p. 43.
1916*a* *Leptocheilopora filliozati* (Brydone) Lang, p. 396.
1916*a* *Leptocheilopora arcuata* Lang, pp. 396, 397.
1921 *Leptocheilopora filliozati* (Brydone): Lang, pp. xcii, 76.
1930 *Leptocheilopora arcuata* Lang: Gaster, table opp. p. 340.
1930 *Leptocheilopora filliozati* (Brydone): Gaster, table opp. p. 340.
1943 *Leptocheilopora arcuata* Lang: Gillard, p. 187.

LECTOTYPE. (Chosen by Lang, 1921 : 77.) S.M., B.36167, the specimen figured by Brydone, 1910, pl. 30, fig. 9. Zoarium, with ancestrula, encrusting a fragment of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Locality 932 of Brydone, lower part of old pit and road cutting at Mount Pleasant, on the road to Micheldever, Andover, Hampshire. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Leptocheilopora* with zooecia of variable size ($Lz = 0{\cdot}55$ to $0{\cdot}75$ mm., $lz = 0{\cdot}35$ to $0{\cdot}55$ mm.); orifice of variable shape but usually with proximal-lateral constrictions; frontal shield fairly well arched with 12–18 well-defined costae and often with a median ridge; ovicells smooth, without lateral slits.

DESCRIPTION. *Zoarium* unilaminar, encrusting. *Adult zooecia* broadly oval, closely placed. A large *distal communication pore* and two or three round *lateral communication pores* are present. *Orifice* always slightly higher than wide, of fairly constant dimensions, rounded distally and laterally, with proximal-lateral constrictions present but seldom pronounced as in *Leptocheilopora magna*, proximal margin usually gently curved proximally. The shape of the orifice varies proximally, where the proximal-lateral constrictions may not be distinct and the proximal margin may vary from straight to proximally curved or, occasionally prominently " V "-shaped : the absence of the constrictions may be due to wear. *Oral spines* 4, slightly thickened, equally spaced on the margin of the orifice. *Frontal shield* gently arched. *Costae* 12–18, distinct, but often very closely placed, rounded at their outer ends and tapering inwards where they may be traced across the *median area of fusion* which is of variable width. No *lateral costal fusions*. In well-preserved specimens some costae show three (possibly four) very small *pelmata* along their mid-line. This feature is easily obscured by wear and possibly also by deposition of calcium-carbonate during fossilization, and may thus not be visible. *Apertural bar* often " V "-shaped, the apex pointing proximally, not much wider than the distal costae, smooth except for a slight median ridge which is often prolonged down the mid-line of the frontal shield. *Gymnocyst* not much exposed laterally, but often visible proximally to each frontal shield. No *interzooecial secondary tissue*.



FIG. 49. *Leptocheilopora filliozati* (Brydone), D. 28294. Holotype of *L. arcuata* Lang. Adult zooecium with median ridge on frontal shield and large distal and lateral communication pores.

Avicularia apparently absent. *Ovicells* prominent, globular, but tending to be slightly pointed distally and flattened in front, hyperstomial, smooth, without *lateral slits*, and generally longer than wide. The lateral pair of oral spines may flank the entrance to the ovicell. *Young zooecia* and *ancestrulae* smaller than the adult zooecia, usually with 6 oral spines, and less often 5 or 4 in the slightly later stages of zoarial growth. There are generally fewer costae (8–12) than in the adult. The orifices, in the young stages, tend to have less marked proximal-lateral constrictions.

MEASUREMENTS. *Adult zooecia* Lz = 0·55 to 0·75 mm.
lz = 0·35 to 0·55 mm.
hr = 0·12 to 0·17 mm.
lr = 0·12 to 0·16 mm.
Young zooecia Lz = 0·33 to 0·54 mm.
lz = 0·22 to 0·37 mm.
hr = 0·09 to 0·15 mm.
lr = 0·08 to 0·13 mm.
Ancestrulae Lz = 0·28 to 0·36 mm.
lz = 0·17 to 0·25 mm.
hr = 0·09 to 0·10 mm.
lr = 0·07 to 0·10 mm.

REMARKS. *Leptocheilopora arcuata* Lang is a junior synonym of *L. filliozati* (Brydone). Of the specimens in the British museum (Natural History) collections available to Lang, he assigned only D.29900 to *L. filliozati* (Brydone). The specimen was presented in December, 1919 but was not mentioned by Lang in his 1921 catalogue. Measurement and comparison of this specimen with those assigned by Lang to *L. arcuata* suggested their synonymy, and this was confirmed by a detailed examination of Brydone's figured specimens of *L. filliozati*, S.M., B.36167–68. Among the nineteen zoaria examined, the adult zooecia of D.39425 and D.39432 were found to be slightly larger than normal and to possess slightly more costae (16–18). These specimens are therefore only tentatively referred to *L. filliozati* (Brydone).

It is of interest to note that the number of costae is not necessarily greater in more adult zooecia, but that occasionally the reverse may be true. For example, in the ancestrula of D.28894 there are apparently 12 costae, but the zooecia of the first growth ring, although larger, have only 9 or 10 costae each ; again, the largest zooecia of the holotype, and of paratypes D.4279, D.28893 and D.28895 have the fewest costae. The zooecium shown in Pl. 8, fig. 1 is distorted and of exceptional length (Lz = 1·30 mm., lz = 0·30 mm., hr = 0·16 mm., lr = 0·14 mm.), in this case there are 35 costae.

In *Leptocheilopora filliozati* (Brydone) Lz appears to vary less than lz, possibly because the width of a zooecium is more directly affected by restrictions imposed on it over a limited substratum. Apparently the development of the orifice is not usually restricted by normal growth conditions : thus, not only does hr slightly exceed lr, except in a very few instances where the two measurements are equal, but the maximum measurements also vary only a little and usually in proportion to one another. Even in the case of the abnormally long zooecium mentioned above the dimensions of the orifice are normal. The measurements of the orifice are therefore important diagnostically, although the orifices of zooecia of the same and of different zoaria may apparently vary in shape. This feature is illustrated by the following diagram (Text-fig. 50) of orifices from some of the specimens examined. Variation also occurs in the shape of the apertural bars. In both cases, some of the variation may be due to wear.

Leptocheilopora filliozati (Brydone) is smaller than *L. magna*, and lacks the constant form of the orifice of that species. It is easily distinguished from *L. tenuilabrosa* which has costate ovicells. The greater size and number of costae, and the fewer (4) oral spines in the adult zooecia of *L. filliozati* distinguish it from *L. vulnerata* (Brydone), which also has a different apertural bar and small lateral slits in the ovicell. *L. filliozati* is also characterized by the frequent development of an irregular median ridge on the frontal shield.



FIG. 50. *Leptocheilopora filliozati* (Brydone)—variation in the shape of the orifice.

STRATIGRAPHICAL DISTRIBUTION. Senonian, (? zone of *B. mucronata*), zones of *G.* [*A.*] *quadrata* and *O. pilula*, subzones of *E. scutata* var. *cincta* and *E. scutata* var. *depressula*.

SPECIMENS. D.28294. Holotype of *L. arcuata* Lang. Incomplete zoarium with ancestrula encrusting a fragment of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Cliffs between Rottingdean and Saltdean, east of Brighton, Sussex.

D.4279. Paratype of *L. arcuata* Lang. Zoarial fragment with ancestrula encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. East Harnham, south of Salisbury, Wiltshire. W. Gamble Collection.

D.4290. Paratype of *L. arcuata* Lang. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.4279.

D.4296. Paratype of *L. arcuata* Lang. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.4279.

D.28893–94. Two paratypes of *L. arcuata* Lang. Zoarial fragments encrusting pieces of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs between last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.

D.28895. Paratype of *L. arcuata* Lang. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit on East Hill, Rottingdean, east of Brighton, Sussex.

D.29900. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.28893–94.

D.39424, D.39426–28. Four zoarial fragments encrusting pieces of echinoid test (D.39424 with ancestrula). Senonian, zone of *G.* [*A.*] *quadrata*.

Pit 15 of Gaster, about one-third of a mile south-south-east of Salvington Windmill, and north of Hill Cottages, Salvington, Sussex.

?D.39425. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.39424.

D.39429. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 26 of Gaster, South Woodleighs Pit, east of gravel pit, north-east of Woodleighs Lodge, about half a mile west of Pit 24 of Gaster, Warningcamp, north-east of Arundel, Sussex.

D.39430. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.39424.

?D.39432. Incomplete zoarium encrusting a piece of echinoid test. Senonian, zone of *B. mucronata.* Pit 8 of Rowe, south of Nodewell, Freshwater, Isle of Wight.

D.40932. Incomplete zoarium encrusting a piece of echinoid test. Senonian, zone of *O. pilula,* subzone of *E. scutata* var. *cincta.* Pit 87 of Gaster, near cart-track to Newlands Barn, north end of East Hill, Rottingdean, Sussex.

D.40933–35. Three incomplete zoaria encrusting fragments of echinoid test, the latter poorly preserved. Senonian, zone of *O. pilula,* subzone of *E. scutata* var. *depressula.* Section 113 of Gaster, cliffs between last groyne east of Rottingdean Gap and Saltdean, Sussex.

D.40936–38. Three incomplete zoaria encrusting fragments of echinoid tests. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 200 yards north of Trumley Copse, Lavant, Sussex.

D.40939. Incomplete zoarium encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata.* Halnaker Chalk Pit, half a mile south of Halnaker Windmill, Boxgrove, Sussex.

OTHER MATERIAL. S.M., B.36167. Lectotype, labelled by Mr A. G. Brighton. "*Cribrilina filliozati* Brydone [Syntype]". See above.

S.M., B.36168. Zoarial fragment encrusting a piece of echinoid test. Labelled by Mr. A. G. Brighton "*Cribrilina filliozati* Brydone [Syntype]". Senonian, zone of *G.* [*A.*] *quadrata.* Locality 1054 of Brydone, Pit near a lane about one-eighth of a mile north-east of the Black Horse Inn, West Titherley, Hampshire. R. M. Brydone Collection.

4. *Leptocheilopora vulnerata* (Brydone)

(Pl. 9, figs. 1, 2; Text-fig. 51)

1914*b* *Cribrilina vulnerata* Brydone, p. 97, pl. 4, figs. 3, 4.
1916*a* *Leptocheilopora vulnerata* (Brydone) Lang, p. 396.
1921 *Leptocheilopora vulnerata* (Brydone): Lang, pp. xc, 70.

LECTOTYPE. (Chosen by Lang, 1921 : 70.) S.M., B.36271, the specimen figured by Brydone, 1914, pl. 4, fig. 3. Maastrichtian, zone of *Belemnella lanceolata* (*O. lunata* Chalk). Trimingham, Norfolk. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Leptocheilopora* with small zooecia (Lz = 0·42 to 0·50 mm., lz = 0·31 to 0·35 mm.) ; orifice small, of fairly constant size (hr = 0·10 to 0·12 mm., lr = 0·09 to 0·11 mm.) ; oral spines 7 ; costae 11–13, wide, distinct ; no intercostal spaces ; apertural bar triangular, produced slightly in the mid-line and sloping down to the proximal margin of the orifice ; ovicells smooth with small lateral slits.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* broadly oval, closely placed, with a single, large, oval *distal communication pore* and one or more smaller, oval *lateral communication pores.* *Orifice* small, distally and laterally rounded with a straight proximal margin ; slight proximal-lateral constrictions may be present. *Oral spines* 7, symmetrically and evenly spaced. *Frontal shield* flat or very gently arched, but descending steeply at the margins to meet the gymnocyst. *Costae* 11–13, wide, tapering inwards, distinct, closely placed. No *intercostal spaces* or *lateral costal fusions.* Three or four small *pelmata* with distinct, low rims occur on each costa ; they decrease in size inwards from the outer ends. *Apertural bar* triangular, with the apex pointing proximally and produced slightly in the mid-line : there is a prominent, curved facet on its descending distal side. *Gymnocyst* occasionally slightly exposed proximally to the frontal shield, but often obscured by the close placing of the zooecia.



FIG. 51. *Leptocheilopora vulnerata* (Brydone), S.M., B.36271. Lectotype. Adult zooecium with seven oral spine-bases, prominent curved facet on apertural bar, distinctly rimmed small pelmata, and large distal—and smaller lateral—communication pores.

Avicularia apparently absent. *Ovicells* smooth, large, prominent, hyperstomial, globular, with small lateral slits. *Young zooecia* smaller than the adult zooecia, but otherwise similar, with 7 oral spines and 9–12 costae; the orifice is only slightly smaller than in the adult zooecia. *Ancestrula* very small, generally obscured.

MEASUREMENTS. *Adult zooecia* Lz = 0·42 to 0·50 mm.
lz = 0·31 to 0·35 mm.
hr = 0·10 to 0·12 mm.
lr = 0·09 to 0·11 mm.
Young zooecia Lz = 0·30 to 0·32 mm.
lz = 0·19 to 0·25 mm.
hr = 0·10 mm.
lr = 0·07 to 0·10 mm.
Ancestrula Not measurable.

REMARKS. *Leptocheilopora vulnerata* (Brydone) is distinguished from other species of the genus particularly by its small zooecial size, its constant number of oral spines, its relatively constant number of costae, and by the form of its apertural bar. The small lateral slits in the ovicell are also apparently peculiar to *L. vulnerata* (Brydone).

The lectotype, S.M., B.36271, and the other specimen originally figured by Brydone (1914*b*, pl. 4, fig. 4) S.M., B.36272, have been examined. Both are exceptionally well preserved and therefore retain very detailed characters such as the small, rimmed pelmata on the costae.

Lang (1921 : 70) doubted whether this species should be included in *Leptocheilopora*, but examination of Brydone's specimens shows that the species is accurately placed there and that Lang was mistaken in classifying the genus in the group of the cribrimorph Polyzoa lacking pelmata.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian, zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk) and ? upper part of zone of *B. mucronata*—Brydone (1914*b* : 97) stated " This species is abundant in the Trimingham Chalk and occurs sparingly in the Weybourne Chalk ".

MATERIAL. S.M., B.36271. Lectotype, labelled " *Cribrilina vulnerata* (Brydone) [Syntype] ". See above.

S.M., B.36272. Zoarium, with ancestrula, encrusting a piece of echinoid test. Labelled " *Cribrilina vulnerata* (Brydone) [Syntype] ". Horizon, locality and collection as for the lectotype.

?N.C.M.76.937 (28/1). Zoarial fragment with young zooecia encrusting a test of *Cardiaster ananchytis* (Leske). Senonian, zone of *B. mucronata.* Thorpe St. Andrews, Norwich, Norfolk. R. M. Brydone Collection.

Species of Uncertain Systematic Position Previously Assigned to *LEPTOCHEILOPORA*

(1) **Leptocheilopora longuessensis** Lang.

1916*a* *Leptocheilopora languessensis* [*sic*] Lang, pp. 396, 397.

See remarks under *Leptocheilopora magna* Lang (p. 106).

(2) ***Reptoporella regularis*** d'Orbigny.

1852 *Reptoporella regularis* d'Orbigny, p. 717, figs. 6, 7.
1853 *Reptoporella regularis* d'Orbigny : d'Orbigny, p. 475.
1854 *Reptoporella regularis* d'Orbigny : d'Orbigny, p. 1097.

REMARKS. Lang (1916*a* : 397 ; 1921 : 70) tentatively referred *Reptoporella regularis* d'Orbigny to *Leptocheilopora*. However, the placing of d'Orbigny's species cannot be at all certain without actual material being examined. The figures show a cribrimorph polyzoan with prominent hyperstomial ovicells and distal to each a smaller round structure with a median proximal pore. Lang (1916*a*) regarded this as the distal communication pore, but this is usually only visible at the edges of a zoarium where the distal walls of the zooecia may be clearly exposed. The structure might even be avicularian. As Lang stated (1921 : 71), " The structure and systematic position of the species, . . . as of the genus *Reptoporella* of which *R. regularis* is the genotype, remains conjectural ".

I do not agree that *Membranipora crenulata* d'Orbigny (1852, pl. 728, figs. 13–15 ; 1853 : 547 ; 1854 : 1093) is, as suggested by Canu (1900*b* : 449) synonymous with *Reptoporella regularis* d'Orbigny. Gillard (1943 : 186) also disagreed with Canu on this point, and stated " la figure de d'Orbigny à été exagérément retouchée au point de prêter aux plus graves confusions. Les septules distales ont été reproduites à l'éxtrémité de chaque zoécie, si bien que Canu a pu supposer que chaque lignée zoéciale débutait par un avicellaire intercalé."

Evidently Gillard assigned several specimens, from the Upper Campanian of the Royan area, to *Leptocheilopora regularis* (d'Orbigny) as he also stated (1943 : 186), " Plusieurs zoaria recueillis à Port-Marant et à la base de la falaise E. de Talmont, se rapportent à cette espèce dont les zoécies montrent des traces d'épines aréales très ténues. Lz = 700–800 μ, lz = 350–375 μ, La [=hr] = 80 μ, la [= lr] = 100 u."

Genus ***PELMATOPORA*** Lang

1851 *Escharipora* d'Orbigny, pls. 603, 684–687, 700, 703 [*partim*].
1852 *Escharipora* : d'Orbigny, p. 221 [*partim*].
1852 *Semiescharipora* d'Orbigny, pls. 717–719 [*partim*].
1853 *Semiescharipora* : d'Orbigny, p. 479 [*partim*].
1854 *Escharipora* : d'Orbigny, p. 1097 [*partim*].
1854 *Semiescharipora* : d'Orbigny, p. 1097 [*partim*].
1860 *Escharipora* : Coquand, p. 121.
1860 *Semiescharipora* : Coquand, p. 150.
1885 *Semiescharipora* : Vine, pp. 116, 156.
?1889*a* *Cribrilina* : Pergens, pp. 61, 69.
1893 *Cribrilina* : Pergens, pp. 202, 216.
1893 *Cribrilina* : Vine, pp. 316, 323, 336.
1900*a* *Cribrilina* : Canu, p. 409.
1900*b* *Cribrilina* : Canu, p. 445 [*partim*].
1900*b* *Escharipora* : Canu, p. 457 [*partim*].
1900 *Cribrilina* : Rowe, p. 341.
1904 *Cribrilina* : Jukes-Browne, p. 490.
1906 *Cribrilina* : Brydone, p. 300.
1910 *Cribrilina* : White, pp. 55, 56.

1911 *Cribrilina*: Canu, pp. 252 [*partim*], 280.
1912 *Cribrilina*: White, pp. 35, 43.
1912 *Cribrilina*: Brydone, pp. 20, 53, 70–89, 95.
1913 *Cribrilina*: Brydone, p. 436 [*partim*].
?1913 *Cribrilina*: White, pp. 24, 27, 30, 32, 38 [*partim*].
1913 *Cribrilina*: Lang, p. 171.
1916*a* *Carydiopora* Lang, p. 94 [*partim*].
1916*a* *Pelmatopora* Lang, p. 101.
1917 *Cribrilina*: Brydone, p. 495.
1919*c* *Pelmatopora*: Lang, p. 107.
1919*d* *Pelmatopora*: Lang, pp. 191, 197, 204, 210.
1921 *Pelmatopora*: Lang, pp xcvi–xcv.
1921 *Cribrilina*: Lang, p. lvii [*partim*].
1922 *Pelmatopora*: Lang, p. 240.
?1923 *Cribrilina*: Waters, p. 546 [*partim*].
1923 *Pelmatopora*: Waters, p. 555.
?1924 *Cribrillina* [*sic*]: Böhm, p. 101.
1924 *Pelmatopora*: Gaster, second of three tables opp. p. 110.
1924*c* *Pelmatopora*: Voigt, p. 110.
?1924*d* *Cribrilina*: Voigt, p. 236 [*partim*].
1924 *Cribrilina*: White, pp. 55, 62.
?1925*a* *Cribrilina*: Voigt, p. 25 [*partim*].
?1925*b* *non Pelmatopora*: Voigt, pp. 98–102.
1927 *Pelmatopora*: Canu & Bassler, pp. 18, 35.
1930 *Pelmatopora*: Gaster, table opp. p. 340.
1930 *non Pelmatopora*: Voigt, pp. 515, 544, 561.
1935 *Pelmatopora*: Bassler, pp. 31, 165.
1939 *non Pelmatopora*: Voigt, p. 93.
1942 *Pelmatopora*: Gillard, p. 90.
1953 *Pelmatopora*: Bassler, p. G192.
1958 *Pelmatopora*: Didon, pp. 80, 82–85.

Type species. *Pelmatopora pero* Lang, 1916*a* : 101. Senonian, top of zone of *M. cor-anguinum*. Medical College Pit, Epsom, Surrey. F. Möckler Collection.

Emended diagnosis. Pelmatoporinae with or without a secondary orifice—where present it comprises a distal oral shield of secondary tissue tending to envelope the unmodified oral spines, or the oral shield may be partly formed by a pair, occasionally two pairs, of enlarged oral spines; apertural bar lacking a median process; avicularia variable, relatively small, rounded, sporadically distributed in the species in which the distal oral shield is not developed, but commonly paired near the orifice, large, distally pointed and directed in the species with distal oral shields.

Remarks. All the material originally assigned to *Pelmatopora* by Lang has been re-examined. Similarly, all the specimens figured by Brydone which Lang regarded as *Pelmatopora* species have been examined. In addition a large number of specimens, particularly from the upper zones of the Senonian, in the C. T. A. Gaster Collection, have been incorporated in the revision.

Lang (1922 : 241), in his remarks on *Pelmatopora*, was concerned primarily with tracing the stratigraphical relationships of the species which he recognized. The present revision shows that several of the species which he described are synonymous, and that their stratigraphical relationships are, in some cases, different from those on which Lang based some of his evolutionary theory.

Lang (1916*a* : 102 ; 1922 : 251) originally distinguished 38 species of *Pelmatopora.* Voigt (1925*b* : 99, 100 ; 1930 : 514) subsequently described two more. Voigt's species are here not definitely assigned to *Pelmatopora* (see p. 200), and only 22 of the species recognized by Lang are maintained here. It has not proved necessary to establish any new species of *Pelmatopora.*

The revision is summarized by the following comparative list of Lang's and present identifications. Only those specimens probably available to Lang when he prepared volume 4 of the *Catalogue of Cretaceous Bryozoa* (1922) are considered. The species are listed in the same order as that used by Lang in 1922.

Lang's identifications (1922)	Present identifications
1. *P. calceata,* Lang	*P. calceata* Lang, p. 123.
2. *P. crepidaria* Lang	*P. crepidaria* Lang, p. 125.
3. *P. solearis* Lang	*P. solearis* Lang, p. 127.
4. *P. larva,* sp. n.	*P. larva* Lang, p. 134 ; and *P. d'orbignyi* Lang [*partim*], p. 140.
5. *P. chrysalis* (d'Orbigny)	*P. chrysalis* (d'Orbigny), p. 132 ; and *P. larva* Lang [*partim*], p. 134.
6. *P. striata* (d'Orbigny)	*P. striata* (d'Orbigny), p. 136.
7. *P. d'orbignyi* Lang	*P. d'orbignyi* Lang, p. 140.
8. *P. pauciclavia* Lang	*P. fragilis* (d'Orbigny), p. 138.
9. *P. fragilis* (d'Orbigny)	*P. fragilis* (d'Orbigny), p. 138.
10. *P. quadrata* Lang	*P. quadrata* Lang, p. 143.
11. *P. filliozati* Lang	*P. quadrata* Lang, p. 143.
12. *P. insignis* (Canu)	? *P. suffulta* (Brydone), p. 151.
13. *P. suffulta* (Brydone)	*P. suffulta* (Brydone), p. 151.
14. *P. gasteri* Lang	*P. gasteri* Lang, p. 130.
15. *P. repleta* (Brydone)	*P. suffulta* (Brydone), p. 151.
16. *P. interrupta* (d'Orbigny)	*P. chrysalis* (d'Orbigny), p. 132 ; and *P. larva* Lang, p. 134.
17. *P. simplex* Lang	*P. simplex* Lang, p. 157.
18. *P. coryli* Lang	*P. coryli* Lang, p. 168.
19. *P. fecampensis* Lang	*P. fecampensis* Lang, p. 145.
20. *P. plantaris* Lang	*P. plantaris* Lang, p. 159.
21. *P. pero* Lang	*P. pero* Lang, p. 148 ; and *P. brydonei* Lang [*partim*], p. 164.
22. *P. brydonei* Lang	*P. brydonei* Lang, p. 164.
23. *P. quadrivolucris* Lang	*P. quadrivolucris* Lang, p. 161.
24. *P. marsupitum* [*sic*] Lang	*P. marsupitorum* Lang, p. 171.
25. *P. roedeanensis* Lang	*P. marsupitorum* Lang, p. 171.
26. *P. somptingensis* Lang	*P. somptingensis* Lang, p. 188.
27. *P. gregoryi* (Brydone)	*P. gregoryi* (Brydone), p. 175 ; and *P. palmata* Lang [*partim*], p. 196.
28. *P. palmata* Lang	*P. palmata* Lang, p. 196.
29. *P. damicornis* Lang	*P. palmata* Lang, p. 196.
30. *P. danktonensis* Lang	*P. somptingensis* Lang, p. 188.
31. *P. bidens* Lang	*P. gregoryi* (Brydone), p. 175.
32. *P. lancingensis* Lang	*P. gregoryi* (Brydone), p. 175.
33. *P. saltdeanensis* Lang	*P. gregoryi* (Brydone), p. 175.

34. *P. collium* Lang	*P. gregoryi* (Brydone), p. 175.
35. *P. promontoriorum* Lang	*P. gregoryi* (Brydone), p. 175.
36. *P. ranunculoides* Lang	*P. gregoryi* (Brydone), p. 175.
37. *P. lacuum* Lang	*P. gregoryi* (Brydone), p. 175.
38. *P. gyrinoides* Lang	*P. gregoryi* (Brydone), p. 175.

Pelmatopora is distinguished from other genera of the Pelmatoporinae by the type of avicularia which are commonly present, and particularly by the absence of prominent distal or proximal oral shields.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *M. cortestudinarium* to *G.* [*A.*] *quadrata*, ?zone of *B. mucronata* and ?Danian.

KEY TO THE SPECIES OF *PELMATOPORA* LANG

(I) *Pelmatopora* with unmodified oral spines.

(A) Median area of fusion narrow; frontal shield arched; orifice often with proximal-lateral constrictions; no lateral costal fusions; gymnocyst well exposed; large pelmata only.

(1) Interzooecial secondary tissue entirely absent; avicularia small, rounded, sporadic . . . . 1. ***P. calceata*** Lang

(2) Interzooecial secondary tissue present.

(*a*) Interzooecial secondary tissue only slightly developed; avicularia small, rounded, tubular, in two pairs near the orifice 2. ***P. crepidaria*** Lang

(*b*) Interzooecial secondary tissue abundant; avicularia small, rounded, one pair near orifice . . . . 3. ***P. solearis*** Lang

(B) Median area of fusion wider.

(1) Orifice small (hr = lr), with marked proximal-lateral constrictions; lateral costal fusions present . . . . 4. ***P. gasteri*** Lang

(2) Orifice without marked proximal-lateral constrictions.

(*a*) Lateral costal fusions absent; interzooecial secondary tissue abundant with marked lacunae; gymnocyst concealed.

(i) Large pelmata only; frontal shield flat.

(*a*′) Orifice higher than wide; avicularia small, rounded, sporadic 5. ***P. chrysalis*** (d'Orbigny)

(*b*′) Orifice wider than higher.

(*a*″) Avicularia large, pointed . . . . 6. ***P. larva*** Lang

(*b*″) Avicularia small, rounded, sporadic, occasionally tending to be paired; 25 or more costae 7. ***P. striata*** (d'Orbigny)

(ii) Pelmata and pelmatidia present; frontal shield arched; 10–15 costae . . . . 8. ***P. fragilis*** (d'Orbigny)

(*b*) Lateral costal fusions present; interzooecial secondary tissue abundant.

(i) Orifice higher than wide; frontal shield flat; avicularia relatively large, pointed, paired . . . . . 9. ***P. d'orbignyi*** Lang

(ii) Orifice wider than high.

(*a*′) Avicularia large, pointed, paired.

(*a*″) Frontal shield arched . . . 10. ***P. quadrata*** Lang

(*b*″) Frontal shield relatively flat.

(*a*‴) Apertural bar wide with slight median furrow; zooecia small, costae well spaced 11. ***P. fecampensis*** Lang (*b'''*) Apertural bar narrow with slight median ridge; zooecia large; costae closely placed; pematidia occasionally present . . . 12. ***P. pero*** Lang

(*b'*) Avicularia small, rounded, paired.

(*a''*) Pelmata and pelmatidia present; 15–20 costae; large zooecia; amount of interzooecial secondary tissue variable but never markedly developed 13. ***P. suffulta*** (Brydone)

(*b''*) Pelmatidia more numerous; Lz less than 1·00 mm.; interzooecial secondary tissue markedly developed with conspicuous lacunae.

(*a'''*) Costae narrow, 16–18 . . 14. ***P. simplex*** Lang

(*b'''*) Costae wide, 13–16 . . 15. ***P. plantaris*** Lang

(II) *Pelmatopora* with enlarged oral spines.

(A) Two pairs of enlarged oral spines present . . 16. ***P. quadrivolucris*** Lang

(B) Only the distal pair of oral spines enlarged.

(1) Proximal pair of unmodified oral spine bases often still visible 17. ***P. brydonei*** Lang

(2) Unmodified oral spines entirely obscured by secondary tissue accumulated round the margin of the orifice (may be visible in early growth stages).

(*a*) Avicularia small, rounded, occasional, sporadic; zooecia small; 11–14 costae. . . . . 18. ***P. coryli*** Lang

(*b*) Avicularia large, distally pointed and directed usually paired.

(i) Enlarged distal oral spines relatively short and peg-like; 14–20 costae . . . . . . 19. ***P. marsupitorum*** Lang

(ii) Enlarged distal oral spines elongate, pointed, relatively slender, with a strong tendency to bifurcate—the outer horn of the bifurcation curving away laterally and proximally . 20. ***P. gregoryi*** (Brydone)

(iii) Enlarged distal oral spines laterally flattened and distally expanded, with a marked tendency to bifurcate . 21. ***P. somptingensis*** Lang

(iv) Enlarged distal oral spines very broad distally and distinctly bilobed . . . . . . 22. ***P. palmata*** Lang

The following species are omitted from the above key and are regarded as of uncertain systematic position: *Pelmatopora incerta* Voigt (1925*b* : 99, 100), *P. grandiporosa* Voigt (1930 : 514), *Pelmatopora* sp. Voigt (1930 : 514), *Pelmatopora daniensis* Voigt (1925*b* : 99, 100). They are discussed on p. 200.

1. *Pelmatopora calceata* Lang

(Pl. 9, fig. 3; Text-fig. 52)

1916*a* *Pelmatopora calceata* Lang, p. 102.
1919*d* *Pelmatopora calceata* Lang: Lang, pp. 191, 196, 210, 218, 223, text-figs. 1, 49.
1921 *Pelmatopora calceata* Lang: Lang, p. xciv.
1922 *Pelmatopora calceata* Lang: Lang, p. 254, pl. 5, fig. 6; text-fig. 74.

HOLOTYPE. D.4032. Well preserved zoarial fragment encrusting a piece of echinoid test. Senonian, either base of zone of *M. cor-anguinum* or top of zone of *M. cortestudinarium*. Chatham, Kent. W. Gamble Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with no distal oral shield; oral spine bases unmodified, in two pairs; median area of fusion narrow; pelmata present but no pelmatidia; frontal shield strongly arched; no interzooecial secondary tissue; avicularia small, apparently rounded, sporadic.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* broadly oval, small. *Orifice* evenly rounded distally, the sides and proximal margin straight, occasionally with proximal-lateral constrictions, the rim not secondarily thickened. *Oral spine bases* 4, not enlarged, unmodified, the proximal-lateral pair placed on the distal-lateral curves of the orifice and not indenting its outline. *Frontal shield* strongly arched. *Costae* 10–14, widely spaced at the outer ends, with no *lateral costal fusions*. Each costa bears a prominent, large *pelma* near its inner end. The outlines of the inner ends of individual costae may be traced in the narrow *median area of fusion*. *Apertural bar* straight, slightly thickened, with two prominent pelmata near the mid-line; no *median process*. *Gymnocyst* well exposed laterally and particularly proximally to the frontal shield. No *interzooecial secondary tissue*.



FIG. 52. *Pelmatopora calceata* Lang, D.4032. Holotype. Adult zooecium and three small, oval, sporadic avicularia.

Avicularia numerous, probably variously directed, sporadic, small, rounded, possibly adventitious on the sides of the zooecia rather than interzooecial, though this is not certain. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·54 to 0·75 mm.
lz = 0·33 to 0·39 mm.
hr = 0·12 mm.
lr = 0·12 mm.
Avicularia Not measurable.

REMARKS. The only available specimen of *P. calceata*, the holotype, has been re-examined. Lang slightly underestimated the dimensions, those of the orifice are particularly constant. The range in number of costae is 10–14 and not 14–16 as stated (Lang, 1922 : 254). The avicularia are very small and it is not possible to determine whether they are proximally or distally directed; there is a slight suggestion of proximal-lateral constrictions in some avicularian apertures, possibly indicating distally directed rostra which are larger than the proximal portions of the apertures. Lang regarded this species as a radical form.

STRATIGRAPHICAL DISTRIBUTION. Senonian, either base of zone of *M. cor-anguinum* or top of zone of *M. cortestudinarium.*

SPECIMEN. D.4032. Holotype—see above.

2. *Pelmatopora crepidaria* Lang

(Pl. 9, fig. 4 ; Text-figs. 53, 54)

1916*a* *Pelmatopora crepidaria* Lang, p. 102.
1919*d* *Pelmatopora crepidaria* Lang : Lang, pp. 211, 222, 224, text-fig. 47.
1921 *Pelmatopora crepidaria* Lang : Lang, p. xciv.
1922 *Pelmatopora crepidaria* Lang : Lang, p. 255, pl. 5, fig. 7 ; text-fig. 75.

HOLOTYPE. D.21200. Well-preserved zoarial fragment. Senonian, zone of *M. cor-anguinum.* Woburn Green, south-west of Beaconsfield, Buckinghamshire. L. Treacher Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; no pelmatidia ; frontal shield very well arched ; interzooecial secondary tissue irregularly developed in small amounts particularly near the orifices ; avicularia small, rounded, prominent, tubular, occurring in one or two pairs near the orifice.

DESCRIPTION. *Zoarium* unilaminar, probably always encrusting. *Zooecia* somewhat elongate, oval. *Orifice* of equal height and width, large, evenly rounded distally and laterally with slight proximal-lateral constrictions and a straight proximal margin ; the outline of the orifice is not obscured by the accumulation of secondary calcareous tissue. *Oral spine-bases* 4, in two pairs, widely placed, well developed, but unmodified and not enlarged, not indenting the outline of the orifice. Secondary closure of the orifice may occur, as in the holotype. An irregular, curved, calcareous shelf may be formed deep in the distal half of the orifice and extend proximally along the lateral margins. In other zooecia the orifice may be completely closed by an irregular lamina of secondary calcareous tissue, see Pl. 9, fig. 4 and Text-fig. 54. *Frontal shield* well arched, elongate. *Costae* 11–14, closely spaced ; no *lateral costal fusions.* Each costa bears a large *pelma* very close to its inner end. *Median area of fusion* narrow. *Apertural bar* prominent, very wide, thickened, with a very slight median ridge and two prominent, large pelmata, one on either side of and close to the mid-line ; no *median process* developed. *Gymnocyst* smooth, somewhat obscured by the irregular, small amounts of *interzooecial secondary tissue* which are developed particularly near the orifices. The gymnocyst is usually exposed laterally and especially proximally to the frontal shields. The interzooecial valleys are deep and sharply defined.

Avicularia very prominent but small, apparently directed distally and obliquely outwards from the zooecia they accompany, rounded and raised on elongate, stout, tubular stalks which project above the general level of the arched frontal shields. The avicularia are usually arranged in a proximal-lateral and a distal-lateral pair near the margins of the orifice : but proximal-lateral pairs occur more frequently than distal-lateral pairs. Other similar, sporadic avicularia are occasionally present. *Ovicells* not seen.

MEASUREMENTS. Zooecia Lz = 0·66 to 0·81 mm.
lz = 0·35 to 0·39 mm.
hr = 0·15 to 0·18 mm.
lr = 0·15 to 0·18 mm.
Avicularia Not measurable.

REMARKS. The holotype, D.21200, and the paratype, D.4110, have been re-examined. The former is a very well preserved zoarial fragment which is now unattached, the latter is a worn zoarial fragment encrusting a piece of echinoid test.



FIGS. 53, 54. *Pelmatopora crepidaria* Lang, D.21200. Holotype.

(53) Adult zooecium with two pairs of tubular avicularia near the orifice.

(54) Distal part of adult zooecium, the tubular avicularia abraded and an irregular calcareous lamina developed in the distal and distal-lateral parts of the orifice.

Lang's measurements for the species are, apparently, maximum dimensions; generally the zooecia are smaller, usually about 0·35 mm. wide and about twice as long. The range in number of costae is 11–14, but usually less than 14. It is not possible to determine whether all the avicularia are distally directed outwards from the zooecia they accompany. The avicularia project and are thus often worn at their upper ends.

Compared with the zooecia of *P. calceata* those of *P. crepidaria* are more closely spaced, their median areas of fusion are not so wide, their costae have less rounded outer ends, and their intercostal spaces are narrower and longer.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum*.

SPECIMENS. D.21200. Holotype—see above.

D.4110. Paratype. Worn zoarial fragment encrusting a piece of echinoid test. Senonian, (zone unspecified). Chatham, Kent. W. Gamble Collection.

3. *Pelmatopora solearis* Lang

(Pl. 9, figs. 5, 6 ; Text-figs. 55, 56)

1916*a* *Pelmatopora solearis* Lang, p. 102.
1919*d* *Pelmatopora solearis* Lang : Lang, pp. 210, 213, 218, 223, text-fig. 50.
1921 *Pelmatopora solearis* Lang : Lang, pp. xciv, xcv.
1922 *Pelmatopora solearis* Lang : Lang, p. 257, pl. 5, fig. 8 ; text-fig. 76.
1930 *Pelmatopora solearis* Lang : Gaster, table opp. p. 340.

HOLOTYPE. D.21211. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Hurley Bottom, Berkshire, south-east of Henley, Oxfordshire. L. Treacher Collection.



FIG. 55. *Pelmatopora solearis* Lang, D. 2763. Paratype. Adult zooecium and two worn avicularia ; there is no interzooecial secondary tissue round the zooecium.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; median area of fusion may be narrow or wide ; no well-developed pelmatidia ; interzooecial secondary tissue relatively abundant ; avicularia small, rounded, sporadic, tending to be paired, one near each proximal-lateral corner of the orifice.

DESCRIPTION. *Zoarium* unilaminar, probably always encrusting. *Adult zooecia* broadly oval, often nearly parallel-sided and elongate. *Orifice* generally a little wider than high, evenly rounded distally, with straight sides and a straight proximal margin. Occasionally very slight proximal-lateral constrictions occur. The outline of the orifice is not obscured by the accretion of secondary calcareous tissue. *Oral spine-bases* 4, unmodified, the lateral pair tending to be larger than the distal pair, but not indenting the outline of the orifice. *Frontal shield* well arched, but often flat in the median area. *Costae* 14-24, usually 14–19, fairly closely spaced with no *lateral costal fusions*. Each costa bears a large *pelma* very close to its inner end.

Very occasionally a small *pelmatidium* may also occur. The pelmata often form a prominent " rim " to the *median area of fusion* which may be narrow and elongate, with the double row of large pelmata lying close together, or it may be wide, with the two rows of pelmata further away from the mid-line of the frontal shield. When the median area of fusion is wide the outlines of the inner ends of individual costae may be traced across it. *Apertural bar* straight and somewhat triangular, prominent, wide, rounded on its upper surface and thickened, with two large, but shallow, pelmata, one on either side of, and close to, the mid-line; no *median process*.



FIG. 56. *Pelmatopora solearis* Lang, D.19510. Paratype. Two adult zooecia and orifice of a proximally adjacent third zooecium and five worn avicularia. The contour-like development of interzooecial secondary tissue is apparent. The frontal shield of the zooecium on the right has a wide median area of fusion, contrasting with that of the zooecium on the left where the pelmata are close to the mid-line.

Gymnocyst smooth, often well exposed, particularly proximally to the frontal shield, but it may be considerably, and even completely, obscured by *interzooecial secondary tissue* accumulated in contour-like ridges and eventually filling the interzooecial furrows. Irregular *lacunae* occur in this secondary tissue.

Avicularia numerous, small, oval to rounded, occasionally somewhat pointed distally, sporadic, or tending to occur in pairs, one avicularium occurring near each proximal-lateral corner of the orifice. *Ovicells* not seen. *Young zooecia* smaller than the adult zooecia. Orifice of the same size and shape as in the adult zooecia. Oral spine-bases 4. Frontal shield more arched than in the adult stages, with about 12 costae. Each costa has a large pelma at its inner end close to the mid-line, the median area of fusion always being very narrow. The apertural bar is stout with two prominent pelmata near the mid-line, and the gymnocyst is well exposed. Interzooecial secondary tissue is scanty or absent in the young stages. The avicularia are small, rounded and generally sporadic, but tend to occur in the paired arrangement found more commonly in the adult stages.

MEASUREMENTS.	*Adult zooecia*	Lz = 0·63 to 1·00 mm.
		lz = 0·32 to 0·50 mm.
		hr = 0·12 to 0·18 mm.
		lr = 0·12 to 0·21 mm.
	Young zooecia	Lz = 0·50 mm.
		lz = 0·30 mm.
		hr = 0·12 to 0·15 mm.
		lr = 0·15 to 0·18 mm.
	Avicularia	Not measurable.

REMARKS. Lang's description was based mainly on the holotype. This, with all paratypes, has been re-examined, as well as several other specimens. In the many zooecia measured, costae were found to be more numerous in the larger than in the smaller ones. D.2813 possesses one abnormally long zooecium (Lz = 1·45 mm., lz = 0·33 mm.) with about 35 costae. The most variable feature is the amount of interzooecial secondary tissue which may be developed. It may be absent or very abundant in different specimens, and occasionally varies from scarce to abundant in the same zoarium. As Lang has suggested, the young zooecia of *P. solearis* are similar to the adult zooecia of *P. calceata*, the radical form, according to Lang, from which *P. solearis* may be derived. Most of the specimens assigned by Lang to *P. solearis* encrust fragments of echinoid test. D.24542, D.27043–46, D.28265 and D.4221 are now unattached.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *M. cortestudinarium* and *M. cor-anguinum*.

SPECIMENS. D.21211. Holotype—see above.

D.28267. Paratype. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed (about 20 ft. above base of zone). Hindover, north-east of Seaford, Sussex.

D.28268. Paratype. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed. Cliff End, east side of Cuckmere Haven, Sussex.

D.24542. Paratype. Zoarial fragment, now unattached. Senonian, zone of *M. cor-anguinum*. Gillingham, north-east of Chatham, Kent. W. Gamble Collection.

D.27042–46. Paratypes. Five fragments of the same unattached zoarium. Senonian, zone of *M. cortestudinarium*. Lower pit, Slines Oak, Worms Heath, Warlingham, Surrey. F. Möckler Collection.

D.19510. Paratype. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cortestudinarium*. Seaford, Sussex. F. Möckler Collection.

D.28269. Paratype. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cortestudinarium*. Cliffs between Hope Gap and Cuckmere Haven, Sussex.

D.2763, D.2813, D.2817, D.27980. Four paratypes. Zoarial fragments now unattached. Senonian, either base of zone of *M. cor-anguinum* or top of zone of *M. cortestudinarium.* Chatham, Kent. G. R. Vine Collection.

D.4026, D. 4971, D.4978. Three paratypes. Zoarial fragments, now unattached. Horizon and locality as for D.2763. W. Gamble Collection.

D.4030, D.4221, D.4965. Three poorly preserved zoarial fragments, now unattached. Horizon and locality as for D.2763. W. Gamble Collection.

D.28265–66. Two zoarial fragments, the former now unattached, the latter encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum, Trochiliopora* Bed, Summit of the Downs, Mount Harry, near Lewes, Sussex.

D.40134. Zoarial fragment encrusting a piece of echinoid test. Lower Senonian, zone unspecified. ?Chatham, Kent. W. Gamble Collection.

D.40135. Zoarial fragment encrusting a piece of echinoid test. Lower Senonian, zone unspecified. ?Chatham, Kent. Collection unspecified.

D.40722–27. Six unattached zoarial fragments. Horizon, locality and collection as for D.28265—see above.

D.40728–31. Four unattached zoarial fragments. Senonian, zone of *M. cor-anguinum, Trochiliopora* Bed. Cliffs, Beltout, west of Eastbourne, Sussex.

D.40732–33. Two unattached zoarial fragments. Senonian, zone of *M. cor-anguinum, Trochiliopora* Bed. Cliffs, east side of Cuckmere Haven, east of Seaford, Sussex.

D.40734–35. Two zoarial fragments encrusting pieces of echinoid tests. Senonian, zone of *M. cortestudinarium.* Cliffs between Hope Gap and Cuckmere Haven, east of Seaford, Sussex.

4. ***Pelmatopora gasteri*** Lang

(Pl. 9, fig. 8 ; Text-fig. 57)

1916*a* *Pelmatopora gasteri* Lang, pp. 102, 105.
1919*d* *Pelmatopora gasteri* Lang : Lang, pp. 211, 218, 222, 226, text-fig. 44.
1921 *Pelmatopora gasteri* Lang : Lang, p. cxiv.
1922 *Pelmatopora gasteri* Lang : Lang, p. 278, pl. 5, fig. 12 ; text-fig. 84.

HOLOTYPE. D.28274. The only specimen. Fragment of a well-preserved zoarium, now unattached. Senonian, zone of *M. cor-anguinum, Trochiliopora* Bed, about 20 ft. above base of zone. Cliffs, east side of Cuckmere Haven, Sussex.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; pelmatidia present ; avicularia tend to be paired but are occasionally sporadic ; costae 15–17 ; zooecia small (Lz = 0·60 to 0·72 mm., lz = 0·33 to 0·36 mm.) ; orifice with marked proximal-lateral constrictions.

DESCRIPTION. *Zoarium* unilaminar, now unattached. *Zooecia* oval, small. *Orifice* of equal height and width, rounded distally and laterally, with a straight proximal margin and with marked proximal-lateral constrictions. *Oral spine-bases* unmodified, occurring in two pairs, the proximal-lateral pair slightly larger. The outline of the orifice may be very slightly indented by the oral spine bases. *Frontal shield* steeply arched, but with a flat median area of fusion. *Costae* 15–17, each with a fairly prominent *pelma* near its inner end, and, closer to the mid-line of the frontal shield, often with an obscure *pelmatidium*. *Lateral costal fusions* occur at the level of the pelmata only. *Median area of fusion* fairly narrow, spindle-shaped. *Apertural bar* prominent, flat, very wide, with two pelmata, one on each side of and close to the mid-line. *Gymnocyst* obscured by *interzooecial secondary tissue*, which, though abundant, does not encroach on to the frontal shields. Slight, irregularly elongate depressions are present in the interzooecial secondary tissue.



FIG. 57. *Pelmatopora gasteri* Lang, D.28274. Holotype. Adult zooecium and paired, worn avicularia. The two avicularia at the proximal end belong to adjacent zooecia.

Avicularia tend to occur just proximally to the proximal-lateral corners of the orifice, often paired. Other occasional, sporadic, interzooecial avicularia occur. All are small, elongate-oval and probably distally directed. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·60 to 0·72 mm.
lz = 0·33 to 0·36 mm.
hr = 0·12 mm.
lr = 0·12 mm.
Avicularia Not measurable.

REMARKS. The unique holotype has been re-examined. The oral spine-bases are larger than is suggested by Lang's figure (1922, text-fig. 84), and the avicularia tend to be paired rather than sporadic. The pelmatidia lie very close to the mid-line in the median area of fusion and are generally obscured by slight extra-calcification or by wear along the mid-line. The form and constant small size of the orifice, the relatively small zooecia, and the restricted range in number of costae particularly distinguish this species. No other specimens have been found in the large amount of material examined from the C. T. A. Gaster Collection.

Stratigraphical distribution. Senonian, zone of *M. cor-anguinum.*

Specimen. D.28274. Holotype—see above.

5. ***Pelmatopora chrysalis*** (d'Orbigny)

(Pl. 9, fig. 7; Text-fig. 58)

1851 *Escharipora chrysalis* d'Orbigny, pl. 686, figs. 6–8 (as *chrisalis* [*sic*]); 1852, p. 228; 1854, p. 1097.
1852 *Semiescharipora interrupta* d'Orbigny, pl. 719, figs. 5–8; 1853, p. 487; 1854, p. 1098.
1860 *Escharipora chrysalis* d'Orbigny: Coquand, p. 121.
1860 *Semiescharipora interrupta* d'Orbigny: Coquand, p. 150.
1900*a* *Cribrilina interrupta* (d'Orbigny): Canu, p. 409.
1900*b* *Cribrilina interrupta* (d'Orbigny): Canu, p. 449.
1900*b* *Escharipora chrysalis* d'Orbigny: Canu, p. 449.
1916*a* *non Pelmatopora interrupta* (d'Orbigny): Lang, p. 102.
1916*a* *Pelmatopora chrysalis* (d'Orbigny) Lang, p. 102.
1917 *Semiescharipora interrupta* d'Orbigny: Brydone, p. 495.
1919*d* *non Pelmatopora interrupta* (d'Orbigny): Lang, pp. 211, 223, text-fig. 42.
1919*d* *Pelmatopora chrysalis* (d'Orbigny): Lang, pp. 212, 223.
1921 *Pelmatopora chrysalis* (d'Orbigny): Lang, p. xciii.
1921 *Pelmatopora interrupta* (d'Orbigny): Lang, p. xciii.
1922 *Pelmatopora chrysalis* (d'Orbigny): Lang, p. 263 [*partim*—D.28438 and D.28439 *excl.*].
1922 *Pelmatopora interrupta* (d'Orbigny): Lang, p. 281, text-fig. 85.
?1942 *Pelmatopora chrysalis* (d'Orbigny): Gillard, p. 90.

Emended diagnosis. *Pelmatopora* with unmodified oral spine-bases; median area of fusion narrow; pelmatidia apparently absent; frontal shield flat; interzooecial secondary tissue abundant, with marked lacunae; avicularia small, rounded, sporadic, unpaired.

Description. *Zoarium* uni- or bilaminar, now unattached. *Zooecia* elongate, elliptical. *Orifice* higher than wide, evenly rounded distally with straight sides and a straight proximal margin; slight proximal-lateral constrictions may occur. The outline of the margin of the orifice is apparently somewhat obscured by the accretion of secondary calcareous tissue. Number of *oral spine-bases* indeterminate, probably 4, but obscured by secondary calcareous tissue. *Frontal shield* flat. *Costae* 19–22, closely spaced. No *lateral costal fusions*. Each costa bears a prominent *pelma* near its inner end: these pelmata tend to form a slight "ridge" on either side of the narrow, lens-shaped *median area of fusion*. No *pelmatidia* visible. *Apertural bar* wide, apparently straight and thickened, with a slight median furrow, and probably carrying two large pelmata near the mid-line; no *median process* present. *Gymnocyst* apparently obscured by plentiful *interzooecial secondary tissue* which fills the interzooecial valleys and contains rather regular, elongate, narrow *lacunae* which are wider at their rounded proximal ends.

Avicularia not very frequent, unpaired, sporadic, small, rounded to slightly elongate. *Ovicells* not seen.

MEASUREMENTS. [Based on Canu's measurements for "*Cribrilina interrupta* (d'Orbigny)", (Canu, 1900*b* : 449).]

Zooecia Lz = 0·92 to 1·00 mm.
lz = 0·42 to 0·50 mm.
hr = 0·24 to 0·26 mm.
lr = 0·17 to 0·21 mm.

REMARKS. Only a photograph (according to Lang, 1922 : 264, the middle specimen of three on a slide labelled "*Escharipora striata* d'Orbigny, Coniacian, Villedieu" in the Canu Collection) is available. The diagnosis and description are based on this photograph (Pl. 9, fig. 7). This specimen compares closely with d'Orbigny's figure of *Escharipora chrysalis* except for the very regular pairing of the avicularia on each zooecium in d'Orbigny's figure.



FIG. 58. *Pelmatopora chrysalis* (d'Orbigny). Drawn from a photograph, in the British Museum (Natural History), of the middle specimen of three on a slide labelled "*Escharipora striata* d'Orbigny, Coniacian, Villedieu." F. Canu Collection, Paris.
An adult zooecium and one worn avicularium surrounded by interzooecial secondary tissue with characteristically elongate lacunae.

Canu (1900*b* : 449), who regarded *Semiescharipora interrupta* d'Orbigny and *Escharipora chrysalis* d'Orbigny as synonymous, based his opinion on a comparison of d'Orbigny's actual specimens and gave measurements for the material. Unfortunately he retained the name *interrupta* for the species, although *chrysalis* has priority by one year.

d'Orbigny's figures of both species are highly stylized, but Lang (1922) rigidly interpreted them and disagreed with Canu's conclusions, keeping the two species separate. Lang did not choose a lectotype for *Escharipora chrysalis*, but did so for *Semiescharipora interrupta*. Such rigid interpretation of stylized figures is not desirable, nor is it advisable to choose a lectotype from them in the absence of actual specimens. Canu's conclusions (1900*b*) based on an examination of d'Orbigny's specimens, are therefore accepted in preference to those founded on such questionable interpretations.

D.28438 and D.28439 were labelled by Lang " *Pelmatopora chrysalis* (d'Orbigny) ", but the former has prominently paired, distally pointed and directed avicularia and is here assigned to *P. larva* Lang : the latter has very numerous narrow costae and has been assigned also, with a query, to *P. larva* Lang.

STRATIGRAPHICAL DISTRIBUTION. Senonian, Coniacian of France.

SPECIMENS. None in the collections.

Photograph—the middle specimen of three on a slide labelled " *Escharipora striata* d'Orbigny, Coniacian, Villedieu ". F. Canu Collection.

6. *Pelmatopora larva* Lang

(Pl. 9, fig. 9 ; Text-figs. 59, 60)

1852	*non Semiescharipora interrupta* d'Orbigny, pl. 719, figs. 5–8 ; 1853, p. 487 ; 1854, p. 1098.
?1916*a*	*Pelmatopora interrupta* (d'Orbigny) : Lang, p. 102.
?1919*d*	*Pelmatopora interrupta* (d'Orbigny) : Lang, pp. 211, 223, text-fig. 42.
1921	*Pelmatopora larva* Lang, p. xcv [*nom nud.*].
1922	*Pelmatopora larva* Lang, p. 260, pl. 5, fig. 9 ; text-fig. 77.
1922	*Pelmatopora chrysalis* (d'Orbigny) : Lang, p. 263 [*partim*—D.28438–39 only].

HOLOTYPE. D.28440. Small fragment of unilaminar zoarium, now unattached. Senonian, Coniacian. La Ribochère, Loir et Cher, east of La Chartres-sur-le-Loir, Sarthe, France. F. Canu Collection.



FIG. 59. *Pelmatopora larva* Lang, D. 28438. Distal end of adult zooecium with two distally pointed avicularia, showing their paired occurrence proximal to the orifice.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; median area of fusion narrow ; no pelmatidia ; frontal shield flat ; interzooecial secondary tissue abundant ; avicularia elongate, distally pointed and directed, tending to be paired, one near each proximal-lateral corner of the orifice.

DESCRIPTION. *Zoarium* uni- or bilaminar, now unattached. *Zooecia* elliptical, long and narrow. *Orifice* wider than high, rounded distally with a straight proximal margin, the sides straight or converging slightly proximally. The outline of the margin of the orifice may be slightly obscured by accretion of secondary calcareous tissue. *Oral spine-bases* apparently 4, unmodified, often obscured by secondary calcareous tissue, and not indenting the outline of the orifice. *Frontal shield* flat, of variable length. *Costae* about 17, fairly closely spaced. No *lateral costal fusions.* Each costa bears a large *pelma* near its inner end close to the narrow *median area of fusion.* No *pelmatidia. Apertural bar* wide, but not very thick, bearing two large pelmata very close to the mid-line; there is no *median process. Gymnocyst* almost entirely obscured by plentiful *interzooecial secondary tissue* which fills the intezooecial valleys to a level generally equal to, or a little above, that of the flat frontal shields. Irregular, elongate, shallow *lacunae* occur frequently in the interzooecial tissue, occasionally descending deeply to expose the gymnocyst.



FIG. 60. *Pelmatopora larva* Lang, D. 28440. Holotype. Adult zooecium and two distally pointed and directed avicularia.

Avicularia relatively large, prominent, elongate, distally pointed and directed; generally paired, one near each proximal-lateral corner of the orifice or a little proximal to this position. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·87 to 1·00 mm.
lz = 0·33 to 0·36 mm.
hr = 0·18 mm.
lr = 0·18 to 0·21 mm.
Avicularia Lz = 0·12 to 0·15 mm.
lz = 0·06 to 0·09 mm.

REMARKS. *P. larva* Lang, *P. interrupta* (d'Orbigny) and *P. chrysalis* (d'Orbigny), as interpreted by Lang (1922), have been reconsidered. *P. interrupta* and *P. chrysalis* are synonymous, and *P. larva* is distinguished from them mainly by the type and position of the avicularia. In the holotype of *P. larva*, D.28440, the avicularia always lie proximally to the apertural bar at or near the proximal-lateral corners of the orifice. Nowhere do they occur on the lateral margins of the orifice as in *P. d'orbignyi* Lang, nor are they of the smaller, rounded, sporadic type which are seen in photographs of a specimen here assigned to *P. chrysalis* (d'Orbigny).

D.28438, labelled by Lang " *P. chrysalis* (d'Orbigny) ", also belongs to *P. larva*, for it possesses paired, distally pointed and directed avicularia which lie proximally to the proximal-lateral corners of the orifice. D.28439, similarly labelled by Lang, is also probably better assigned to *P. larva*. Although its costae are constant in number at 21, the dimensions of the zooecia, the well-developed, distally pointed and directed, paired avicularia (one proximal to each proximal-lateral corner of the orifice), and the abundant interzooecial secondary tissue with shallow, elongate lacunae are all features which are more characteristic of *P. larva* than of *P. chrysalis*. The measurements of the specimen are as follows :

Zooecia Lz = 0·87 mm.
lz = 0·36 mm.
hr = 0·15 to 0·18 mm.
lr = 0·18 to 0·21 mm.
Avicularia Lz = 0·15 mm.
lz = 0·09 mm.

STRATIGRAPHICAL DISTRIBUTION. Senonian, Coniacian of France.

SPECIMENS. D.28440. Holotype—see above.

D.28438. Small unattached fragment of bilaminar zoarium, labelled by Lang " *P. chrysalis* (d'Orbigny) ". Horizon, locality and collection as for the holotype.

?D.28439. Small zoarial fragment, now unattached, labelled by Lang " *P. chrysalis* (d'Orbigny) ". Horizon, locality and collection as for the holotype.

7. ***Pelmatopora striata*** (d'Orbigny)

(Pl. 9, fig. 10)

1851	*Escharipora striata* d'Orbigny, pl. 686, figs. 9–12 ; 1852, p. 229 ; 1854, p. 1097.
1851	*Escharipora mumia* d'Orbigny, pl. 687, figs. 4–6 ; 1852, p. 233 ; 1854, p. 1097.
1851	*non Escharipora elegans* d'Orbingy, pl. 684, figs. 13–15 ; 1852, p. 222 ; 1854, p. 1097.
1900*b*	*Cribrilina* (*Decurtaria*) *striata* (d'Orbigny) Canu, p. 450.
1900*b*	*non Escharipora elegans* (d'Orbigny) : Canu, p. 457.
1916*a*	*Pelmatopora striata* (d'Orbigny) Lang, pp. 102, 104.
1919*d*	*Pelmatopora striata* (d'Orbigny) : Lang, pp. 212, 222.
1921	*Pelmatopora striata* (d'Orbigny) : Lang, p. xciii.
1922	*Pelmatopora striata* (d'Orbigny) : Lang, p. 264.
?1942	*Pelmatopora striata* (d'Orbigny) : Gillard, p. 90.

LECTOTYPE. (Chosen by Lang, 1922 : 265.) The specimen figured by d'Orbigny, 1851, pl. 686, fig. 10. Néhou, south of Valoignes, east of Ste. Colombe, Manche, France. Senonian.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; apparently no pelmatidia ; frontal shield flat ; about 25 costae.

DESCRIPTION. *Zoarium* uni- or bilaminar, now unattached. *Zooecia* large, elliptical. *Orifice* rounded distally and laterally, with a straight proximal margin, apparently higher than wide. Number and form of *oral spine-bases* not determinable, but definitely not enlarged, probably obscured by the accretion of secondary calcareous tissue. *Frontal shield* apparently flat. *Costae* about 25 or 26, narrow, closely spaced, each with a large *pelma* near its inner end close to the *median area of fusion*. This is somewhat wider than in *P. chrysalis* (d'Orbigny). The outlines of the inner ends of individual costae may be traced into the narrow median area of fusion. No *pelmatidia* and no *lateral costal fusions* visible. *Apertural bar* prominent, rather wide, with a slight median furrow, and with two large pelmata, one on either side of the mid-line. *Gymnocyst* apparently entirely concealed by the very abundant *interzooecial secondary tissue* which completely fills the interzooecial valleys and stands up as ridges between the zooecia. Irregular elongate *lacunae* occur in the secondary tissue.

Avicularia are possibly present but the photograph on which the present diagnosis and description are based is not clear enough to show them with certainty. *Ovicells* not seen.

MEASUREMENTS. [From Canu (1900*b* : 450).]
Zooecia Lz = 1·14 to 1·35 mm.
lz = 0·50 to 0·64 mm.
hr = 0·20 to 0·30 mm.
lr = 0·21 to 0·25 mm.

REMARKS. The diagnosis and description for *P. striata* (d'Orbigny) are only tentative. Canu (1900*b* : 450) based his conclusions on an examination of d'Orbigny's actual specimens and regarded *Escharipora striata* and *E. mumia* as synonymous. This interpretation is followed here, since d'Orbigny's stylized figures have little systematic value.

As Lang has remarked (1922 : 265), there are no indications of pelmata in either of d'Orbigny's figures of the two species which Canu regarded as synonymous, and therefore the placing of *E. striata* in the genus *Pelmatopora* can only be tentative. However, the slide in the Canu Collection, labelled " *Escharipora striata* d'Orbigny ", holds three specimens : the outermost specimens compare closely with the d'Orbigny figures and are definitely *Pelmatopora*. A photograph of one of these outer specimens is in the collections of the British Museum (Natural History). The middle specimen on the slide has been assigned to *P. chrysalis* (d'Orbigny).

STRATIGRAPHICAL DISTRIBUTION. Senonian, Coniacian of France.

SPECIMENS. A photograph only—one of the outer specimens of three on a slide labelled " *Escharipora striata* d'Orbigny, Coniacian, Villedieu ". F. Canu Collection.

8. *Pelmatopora fragilis* (d'Orbigny)

(Pl. 10, figs. 1, 2; Text-fig. 61)

1851 *Semiescharipora fragilis* d'Orbigny, pl. 717, figs. 8–11; 1853, p. 480; 1854, p. 1097.
1885 *Semiescharipora fragilis* d'Orbigny: Vine, pp. 116, 156.
1893 *Cribrilina fragilis* (d'Orbigny) Pergens, pp. 202, 216.
1893 *Cribrilina fragilis* (d'Orbigny): Vine, pp. 316, 323, 336.
1900*b* *Cribrilina fragilis* (d'Orbigny): Canu, pp. 448, 458.
1904 *Cribrilina fragilis* (d'Orbigny): Jukes-Browne, p. 490.
1916*a* *Pelmatopora fragilis* (d'Orbigny) Lang, pp. 102, 104.
1916*a* *Pelmatopora pauciclavia* Lang, pp. 102, 104.
1919*d* *Pelmatopora fragilis* (d'Orbigny): Lang, pp. 212, 222.
1919*d* *Pelmatopora pauciclavia* Lang: Lang, p. 211, text-fig. 41.
1921 *Pelmatopora pauciclavia* Lang: Lang, p. xciv.
1921 *Pelmatopora fragilis* (d'Orbigny): Lang, p. xcv.
1922 *Pelmatopora pauciclavia* Lang: Lang, p. 267, pl. 5, fig. 11; text-fig. 80.
1922 *Pelmatopora fragilis* (d'Orbigny): Lang, p. 269.
1942 *Pelmatopora fragilis* (d'Orbigny): Gillard, p. 90.

LECTOTYPE. (Chosen by Lang, 1922: 269.) The specimen figured by d'Orbigny, 1851, pl. 717, fig. 9. Senonian, Coniacian. Fécamp, Seine Inférieure, France.



FIG. 61. *Pelmatopora fragilis* (d'Orbigny), D. 28273. Holotype of *P. pauciclavia* Lang. Adult zooecium and paired avicularia. The avicularia at the proximal end belong to adjacent zooecia.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases; median area of fusion narrow; pelmatidia often present; frontal shield high, flat on top; costae 9–15; orifice relatively large; avicularia often paired.

DESCRIPTION. *Zoarium* unilaminar, encrusting or now unattached. *Zooecia* broadly oval or with somewhat parallel sides, of variable length. *Orifice* relatively large, usually higher than wide, rounded distally with straight sides and a straight proximal margin. Occasionally the orifice may narrow slightly at the proximal end; there are no proximal-lateral constrictions. *Oral spine-bases* 4, widely spaced, unmodified, possibly very slightly enlarged, but not indenting the outline of the margin of the orifice. *Frontal shield* high, flat on top. *Costae* wide, 9–15 (often 10–11), each with a prominent, raised *pelma* near the inner end at the edge of the median area of fusion: an inner *pelmatidium* is often also present on each costa. *Lateral costal fusions* occur at the level of the pelmata. The area within the pelmata is flat; outside it the costae slope down fairly steeply. *Median area of fusion* narrow. *Apertural bar* thick, prominent, with two large pelmata, one immediately on either side of the mid-line which is marked by a slight median ridge. *Gymnocyst* largely concealed by fairly abundant *interzooecial secondary tissue* which contains irregularly elongate, well-defined *lacunae.*

Avicularia interzooecial, tubular, prominent, often paired one on either side of each zooecium and placed just proximally to the apertural bar, or well proximal to this position. Apparently the avicularian apertures are small, rounded to oval, and, in less worn examples, may be slightly pointed distally. *Ovicells* not seen.

MEASUREMENTS. [Based partly on Canu's measurements for " *Cribrilina fragilis* (d'Orbigny) ", 1900*b* : 448.]

Zooecia Lz = 0·71 to 0·81 mm.
lz = 0·39 to 0·50 mm.
hr = 0·14 to 0·18 mm.
lr = 0·11 to 0·18 mm.

REMARKS. *P. fragilis* was poorly figured by d'Orbigny. Lang (1922 : 269) interpreted the species from a photograph provided by Canu of a specimen in the d'Orbigny Collection. That specimen was one of three on a slide: one of the other two specimens is specifically distinct and was made the type of *P. filliozati* Lang.

Although it is difficult to interpret the photograph of *P. fragilis* it is apparent that the frontal shields are not flat, as suggested by Lang in his diagnosis of the species, but are high and sloping outwards, and that the avicularia tend to occur one on either side of each frontal wall near the apertural bar.

The measurements of the holotype of *P. pauciclavia* Lang, D.28273, are as follows:

Zooecia Lz = 0·78 to 0·81 mm.
lz = 0·39 to 0·47 mm.
hr = 0·18 mm.
lr = 0·15 to 0·18 mm.

The range in number of costae in the specimen is 9–12. Comparing this with the measurements given by Canu (1900*b* : 448) for " *Cribrilina fragilis* (d'Orbigny) ", it is apparent that the dimensions of the two are closely similar. Canu recorded the dimensions of the latter species as Lz = 0·71 to 0·78 mm., lz = 0·42 to 0·50 mm., hr = 0·14 mm. and lr = 0·11 mm. The range in number of costae discernible in the photograph of *Pelmatopora fragilis* is 13–15, and the amount of interzooecial secondary tissue present also compares closely with that in D.28273. D.28452, labelled by Lang " *Pelmatopora ?fragilis* (d'Orbigny) ", is rather badly worn but is tentatively retained in the species. D.28441–42, included in *P. fragilis* by Lang, are, however, here excluded from the species. The former has a flat frontal shield with at least 16 narrow costae, and an amount of interzooecial secondary tissue different from that in *P. fragilis*. Other details are destroyed in this badly damaged specimen and, like D.28442, it is indeterminate.

STRATIGRAPHICAL DISTRIBUTION. Senonian, Coniacian in France, zone of *M. cor-anguinum* in England.

SPECIMENS. D.28452. Worn zoarial fragment encrusting a piece of molluscan shell. Senonian, Coniacian. St. Avertin, south-east of Tours, Indre-et-Loire, France. F. Canu Collection.

D.28273. Holotype of *P. pauciclavia* Lang. Very well preserved zoarial fragment. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed (about 20 ft. above base of zone). East side of Cuckmere Haven, cliffs, Sussex.

Photograph—one of three specimens on a slide labelled " *Cribrilina* [*Semiescharipora*] *fragilis* d'O.". Senonian, Emscherian. Fécamp, Seine Inférieure, France. F. Canu Collection.

Excluded from the species—indeterminate—D.28441, D.28442.

9. *Pelmatopora d'orbignyi* Lang

(Pl. 10, fig. 3 ; Text-figs. 62, 63)

1916*a* *Pelmatopora d'orbignyi* Lang, pp. 102, 104.
1919*d* *Pelmatopora d'orbignyi* Lang : Lang, pp. 211, 222, 225, text-fig. 48.
1921 *Pelmatopora d'orbignyi* Lang : Lang, p. xcv.
1922 *Pelmatopora larva* Lang, p. 260 [*partim*—paratype D.28443 only].
1922 *Pelmatopora d'orbignyi* Lang : Lang, p. 266, pl. 5, fig. 10 ; text-fig. 79.

HOLOTYPE. D.28453. Small fragment of erect, bilaminar zoarium. Senonian, Coniacian. St. Avertin, south-east of Tours, Indre-et-Loire, France. F. Canu Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; pelmatidia occasionally present ; frontal shield flat ; interzooecial secondary tissue abundant ; avicularia prominent, large, distally pointed and directed, paired at the proximal-lateral corners of the orifice.

DESCRIPTION. *Zoarium* unilaminar and probably encrusting, or bilaminar and erect. *Zooecia* long, elliptical, somewhat pointed proximally. *Orifice* higher than wide, evenly rounded distally and laterally with a straight proximal margin, proximal-lateral constrictions variably developed. The outline of the margin of the orifice is somewhat obscured by secondary tissue and is indented laterally by a pair of large avicularia. *Oral spine-bases* probably 4, the distal pair are often visible though somewhat immersed in secondary tissue : the lateral pair, if present, are concealed by the large, lateral, oral avicularia or by secondary tissue. *Frontal shield* apparently flat, the outer ends of the costae descend sharply, but are largely obliterated by the abundant interzooecial secondary tissue. *Costae* 14–20, each with a large *pelma* near the inner end. Some costae may also possess a *pelmatidium* nearer to the mid-line of the zooecium. *Lateral costal fusions* are often present at the level of the pelmata, producing an irregularly developed row of *intercostal spaces* very close to the mid-line and a series of outer intercostal spaces which are slot-like. *Median area of fusion* very narrow and irregular. *Apertural bar* somewhat triangular, wide, but not very thick, with two prominent pelmata one on each side of and close to the mid-line; a slight median ridge is present, but there is no *median process*. *Gymnocyst* almost entirely concealed by plentiful *interzooecial secondary tissue* which contains irregular but sharply defined elongate *lacunae* through which the smooth gymnocyst may be locally exposed.



FIG. 62. *Pelmatopora d'orbignyi* Lang, D.28443. Paratype of *P. larva* Lang. Adult zooecium with two worn lateral oral avicularia and adjacent broken zooecium with two well-preserved lateral oral avicularia.

Avicularia prominent, large, paired at the proximal-lateral corners of the orifice, almost as long as the orifice, raised, elongate, pointed, directed distally and obliquely inwards, divided by a *transverse bar* into a larger, rounded, proximal portion and a small, triangular *rostrum*. They are not deeply immersed in the interzooecial secondary tissue. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·94 to 1·15 mm.
lz = 0·46 to 0·50 mm.
hr = 0·21 to 0·24 mm.
lr = 0·15 to 0·18 mm.
Avicularia Lz = 0·15 to 0·24 mm.
lz = 0·12 mm.



FIG. 63. *Pelmatopora d'orbignyi* Lang, D.28453. Holotype. Adult zooecium and three avicularia surrounded by very abundant interzooecial secondary tissue.

REMARKS. The holotype (D.28453) has been re-examined. D.28443, a paratype of *P. larva* Lang, has been assigned to *P. d'orbignyi* also. Occasional pelmatidia are present on the costae in the median area of the frontal shield in D.28443, and its dimensions and range in number of costae are similar to those of the holotype of *P. d'orbignyi*. D.28443 also has two particularly well-preserved lateral oral avicularia which are large, raised, and distally pointed and directed. Lang described the specimen as " much worn " ; however, it is not as worn as the holotype of *P. d'orbignyi*, in which the avicularia are considerably abraded but regularly paired close to the proximal-lateral corners of the orifice. The position of these avicularia was regarded by Lang as diagnostic for the species, and D.28443 accordingly may be placed in *P. d'orbignyi* Lang.

The holotype has a bilaminar zoarium, but the two laminae of zooecia are not aligned, as is usually the case with normally bilaminar forms, but cross obliquely.

STRATIGRAPHICAL DISTRIBUTION. Senonian, Coniacian of France.

SPECIMENS. D.28453. Holotype—see above.

D.28443. Paratype of *P. larva* Lang. Small fragment of unilaminar zoarium. Coniacian. La Ribochère, Loir-et-Cher, east of La Chartre-sur-le-Loir, Sarthe, France. F. Canu Collection.

10. ***Pelmatopora quadrata*** Lang

(Pl. 10, figs. 4, 5 ; Text-fig. 64)

1916*a*	*Pelmatopora quadrata* Lang, pp. 102, 104.
1916*a*	*Pelmatopora filliozati* Lang, pp. 102, 105.
1919*d*	*Pelmatopora quadrata* Lang : Lang, pp. 197, 211, 222, text-fig. 40.
1919*d*	*Pelmatopora filliozati* Lang : Lang, pp. 197, 212, 222.
1921	*Pelmatopora quadrata* Lang : Lang, p. xciv.
1921	*Pelmatopora filliozati* Lang : Lang, p. xciv.
1922	*Pelmatopora quadrata* Lang : Lang, p. 270, text-fig. 81*a*, *b*.
1922	*Pelmatopora filliozati* Lang : Lang, p. 272, text-fig. 81*c*.

HOLOTYPE. D.28271. Well-preserved zoarial fragment with numerous ovicells. The fragment is now unattached. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed (about 20 ft. above base of zone). Cliffs, east side of Cuckmere Haven, Sussex.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; pelmatidia present ; costae 15–20 ; frontal shield well arched ; avicularia elongate, distally pointed and directed, tending to be paired.

DESCRIPTION. *Zoarium* unilaminar, now unattached. *Zooecia* broadly oval, of variable length. *Orifice* wider than high, evenly rounded distally and laterally, with a straight proximal margin ; there are no proximal-lateral constrictions. Where an ovicell is present the orifice is transversely rectangular in outline. *Oral spine-bases* 4, often obscured by the accretion of secondary tissue near the orifice or by the development of ovicells. The oral spines do not indent the outline of the margin of the orifice. *Frontal shield* well arched, but slightly flattened on top in the median area. *Costae* 13–20 (often 15 or 20), each with a prominent *pelma* near the inner end at the edge of the median area of fusion and a *pelmatidium* almost at the extreme inner end. *Lateral costal fusions* occur at the level of the pelmata. *Median area of fusion* of variable width. *Apertural bar* not very wide, relatively flat, with a marked median furrow and carrying a pair of prominent pelmata, one on either side of the mid-line. *Gymnocyst* almost completely concealed by abundant *interzooecial secondary tissue* which lacks well-formed lacunae.

Avicularia not very large, but prominent, distally pointed and directed; a *transverse bar* divides a small, semicircular proximal portion from a long, narrow *rostrum*. There is a strong tendency for the avicularia to be paired, one near each proximal-lateral corner of the orifice; occasional sporadic avicularia also occur.

Ovicells apparently endozooecial, well immersed. Where ovicells are developed a distal oral shield is formed by the infilling of the space between the distal oral spines by a transverse shelf of calcareous tissue. This transverse shelf reduces the size of the orifice which becomes transversely oblong. The lateral pair of oral spines may be displaced proximally, appearing at the distal-lateral corners of the ovicelled orifice and not, as is sometimes the case, obscured by the calcareous tissue.



FIG. 64. *Pelmatopora quadrata* Lang, D.28271. Holotype. Ovicelled adult zooecium with paired avicularia. The avicularium at the proximal end belongs to an adjacent zooecium.

MEASUREMENTS. *Zooecia* Lz = 0·81 to 0·85 mm.
lz = 0·39 to 0·43 mm.
hr = 0·12 mm.
lr = 0·15 mm.
Avicularia Lz = 0·15 mm.
lz = 0·09 mm.

REMARKS. The holotype of *P. filliozati* Lang is the specimen nearest to the label on a slide with three specimens from the F. Canu Collection labelled "*Cribrilina* [*Semiescharipora*] *fragilis* d'O." (see Lang, 1922 : 273). A photograph of this specimen is in the collections of the British Museum (Natural History). Careful comparison of this photograph with the holotype of *P. quadrata* Lang shows that the two species are synonymous; in the photograph most zooecia possess 18–20 costae, and in the holotype of *P. quadrata* the range is apparently 13–16, most zooecia having 15 costae. This is the principal difference between the two. Ovicells are developed in the Canu specimen producing the same oblong, modified orifice. The median area of fusion is similar and the avicularia apparently pointed and tending to be paired. It has not been possible to discover the evidence for the measurements which Lang gave for the Canu specimen. The photograph has no scale.

STRATIGRAPHICAL DISTRIBUTION. Senonian, Emscherian in France; zone of *M. cor-anguinum*, *Trochiliopora* Bed, in England.

SPECIMENS. D.28271. Holotype—see above.

D.28272. Paratype. Fragment of a well-preserved, now unattached, zoarium. Horizon, locality and collection as for the holotype.

Photograph of a specimen in the F. Canu Collection. The specimen is mounted nearest to the label on a slide containing three specimens labelled "*Cribrilina* [*Semiescharipora*] *fragilis* d'O.". The specimen was chosen by Lang as the holotype of his species *Pelmatopora filliozati*. Senonian, Emscherian. Fécamp, north-east of Le Havre, Seine Inférieure, France.

11. ***Pelmatopora fecampensis*** Lang

(Pl. 10, fig. 6; Text-figs. 65–68)

1916*a*	*Pelmatopora fecampensis* Lang, pp. 102, 105.
1919*d*	*Pelmatopora fecampensis* Lang: Lang, pp. 211, 222, text-fig. 43.
1921	*Pelmatopora fecampensis* Lang: Lang, p. xcv.
1922	*Pelamtopora fecampensis* Lang: Lang, p. 286, pl. 6, fig. 3; text-fig. 88.
1958	*Pelmatopora fecampensis* Lang: Didon, pp. 82, 84, text-fig. 9.

HOLOTYPE. D.28473. Zoarial fragment, now unattached. Senonian, Coniacian. Fécamp, north-east of Le Harvre, Seine Inférieure, France. F. Canu Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases; orifice wider than high, larger than in *P. gasteri*; costae 16–21 (often 18 or 19); pelmatidia present; outermost intercostal spaces long and slit-like; interzooecial secondary tissues abundant, lacking well-marked lacunae; avicularia distally pointed and directed, generally two to each zooecium.

DESCRIPTION. *Zoarium* unilaminar, probably encrusting but now unattached. *Zooecia* elliptical, the frontal shields fairly closely spaced. *Orifice* a little wider than high, of very constant size, lacking proximal-lateral constrictions, rounded distally and laterally with a straight proximal margin, its outline not obscured by secondary calcareous tissue. *Oral spine-bases* 4, small and unmodified. *Frontal shield* flatly arched, the *median area of fusion* depressed and narrow. *Costae* 16–21 (often 18 or 19), each with a prominent *pelma* about half-way along its length, coinciding with the edge of the median area. In addition, each costa usually carries a *pelmatidium* closer to the mid-line of the zooecium. *Lateral costal fusions* are developed at the level of the pelmata. *Outermost intercostal spaces* long and slit-like. *Apertural bar* fairly wide, with two large pelmata near the mid-line which is marked by a slight furrow. *Gymnocyst* concealed by abundant *interzooecial secondary tissue* which fills the interzooecial valleys and partly immerses some avicularia. There are usually no well-defined *lacunae* in the interzooecial tissue.

Avicularia small but prominent, distally pointed and directed, the apertures semicircular proximally, narrowing distally into very elongate triangular *rostra*. *Transverse bars* are apparently absent. Very commonly, two variably placed avicularia are associated with each zooecium; occasionally other sporadic avicularia of the same type occur. *Ovicells* apparently endozooecial, but globular and often prominent, tending to reduce the size of the orifice, either obliterating the oral spines or developing apparently within the orifice without displacing the oral spines (Text-figs. 66–68).



FIG. 65. *Pelmatopora fecampensis* Lang, D. 28473. Holotype. Adult zooecium and three avicularia.

MEASUREMENTS. *Zooecia* Lz = 0·75 to 0·90 mm.

lz = 0·39 to 0·50 mm.
hr = 0·15 mm.
lr = 0·18 mm.
Avicularia Lz = 0·12 to 0·15 mm.
lz = 0·06 mm.

REMARKS. The holotype and five paratype zoarial fragments of *P. fecampensis* have been re-examined. The avicularia tend to occur regularly, two associated with each zooecium, but they are not constantly developed in the same position. The ovicells, though probably endozooecial, are very prominent and lack the thickened edge characteristic of the endozooecial ovicells of other species, such as *P. somptingensis* Lang and *P. gregoryi* (Brydone).

Paratype D.28472 consists of only four zooecia of normal dimensions one of which is ovicelled. The ovicell is developed apparently within the orifice of the zooecium without displacing the oral spines (Text-fig. 68). Lacunae are quite well developed in the interzooecial secondary tissue of this specimen.



FIG. 66. *Pelmatopora fecampensis* Lang, D. 28475. Paratype. Distal end of ovicelled zooecium, three avicularia, and part of distally adjacent zooecium, the ovicell has a proximally extended calcareous lamina which almost closes the orifice.

Paratypes D.28274 and D.28477 are fragments of zoaria which are now unattached. The former is well preserved with one zoarial lamina encrusting another ; D.28477 is a very small fragment with well-preserved avicularia. D.28476 is another paratype zoarial fragment with only one complete zooecium preserved and no unworn avicularia.



FIG. 67. *Pelmatopora fecampensis* Lang, D.28475. Paratype. Distal end of ovicelled zooecium showing apparent displacement of the lateral oral spines by the ovicell.

Paratype D.28475 is a larger zoarial fragment with prominent oral spine bases and well-preserved avicularia. At least three of its zooecia have ovicells which are apparently endozooecial, but very prominent and globular. In one case the edge of the ovicell is apparently prolonged by a lamina which almost completely closes the orifice, and a thickened, transverse band of tissue runs across the ovicell, possibly marking its normal limit. Several of the zooecia have developed thin calcareous laminae entirely sealing the orifices, and it is possible that the prolongation associated with the ovicell just mentioned is part of such a lamina. In this case the development of the ovicell has obliterated the oral spines, but in an adjacent zooecium two of the oral spines, probably the lateral pair, have been displaced by the development of the ovicell but still remain visible.



FIG. 68. *Pelmatopora fecampensis* Lang, D.28472. Paratype. Distal end of ovicelled zooecium with two avicularia, showing the apparent development of an ovicell inside the orifice, but not displacing the oral spines.

STRATIGRAPHICAL DISTRIBUTION. Senonian, Coniacian of France.

SPECIMENS. D.28473. Holotype—see above.

D.28472, D.28474–77. Five paratypes. Zoarial fragments now unattached. Horizon, locality and collection as for the holotype.

12. ***Pelmatopora pero*** Lang

(Pl. 10, fig. 7; Text-fig. 69)

1916*a* *Pelmatopora pero* Lang, pp. 102, 105.
1919*c* *Pelmatopora pero* Lang: Lang, p. 107.
1919*d* *Pelmatopora pero* Lang: Lang, pp. 210, 213, 216, 218, 220, 222, text-fig. 52.
1921 *Pelmatopora pero* Lang: Lang, pp. lxiii, xciii.
1922 *Pelmatopora pero* Lang: Lang, p. 290, pl. 6, fig. 5; text-fig. 90 [*partim*—D.29861 *excl.* = *P. brydonei*].
1935 *Pelmatopora pero* Lang: Bassler, p. 165.
1953 *Pelmatopora pero* Lang: Bassler, p. G192, text-fig. 144, 7.

HOLOTYPE. D.23405. Zoarial fragments, now unattached. Senonian, extreme top of zone of *M. cor-anguinum*. Medical College Pit, Epsom, Surrey. F. Möckler Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases; zooecial length about 1·00 mm.; costae 13–20 (often 16–18); two avicularia to each zooecium, usually not symmetrically paired, large, prominent, distally pointed and directed.

DESCRIPTION. *Zoarium* unilaminar, encrusting, but may now be unattached. *Zooecia* elongate, elliptical, tending to be somewhat pointed proximally. *Orifice* wider than high, rounded distally and laterally with a straight proximal margin, of very constant size, without proximal-lateral constrictions. *Oral spine-bases* 4, unmodified, sometimes slightly obscured by secondary tissue. *Frontal shield* flat. *Costae* 13–20 (often 16–18), fairly closely spaced but well defined, each with a prominent *pelma* about half-way along its length and a *pelmatidium* nearer to its inner end. *Lateral costal fusions* are developed at the levels of both the pelmata and pelmatidia, producing a paired row of *intercostal spaces* on either side of the median area. Occasionally a second paired row of irregularly developed intercostal spaces may be present close to the mid-line, when a second series of minute pelmatidia is also developed. The *median area of fusion* is thus of variable width but generally narrow. It is commonly level with, but may occasionally be depressed slightly below, the level of the rest of the frontal shield. *Apertural bar* not very wide, with a very slight median ridge and two indistinct pelmata, one on either side of the mid-line. *Gymnocyst* concealed by plentiful *interzooecial secondary tissue* which almost fills the interzooecial valleys and contains shallow, irregularly elongate depressions.



FIG. 69. *Pelmatopora pero* Lang, D. 23411. Paratype. Adult zooecium with paired, damaged avicularia just proximal to the orifice and, at the proximal end, two undamaged avicularia belonging to adjacent zooecia.

Avicularia prominent, large, elongate, distally pointed and directed, rounded proximally with a long, triangular *rostrum*. There are two avicularia to each zooecium, but they are not symmetrically paired. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·81 to 1·12 mm.
lz = 0·39 to 0·63 mm.
hr = 0·15 to 0·18 mm.
lr = 0·18 to 0·21 mm.
Avicularia Lz = 0·15 to 0·24 mm.
lz = 0·09 mm.

REMARKS. The holotype, fifteen paratypes and two other specimens labelled by Lang *Pelmatopora pero* have been re-examined and retained in this species, but one, D.29861, also labelled as that species, is referred to *P. brydonei* Lang.

Where possible the zooecia of all specimens have been measured. Their dimensions vary considerably, but those of the orifice remain very constant. The avicularia, where unworn, are always elongate and sharply pointed distally. Well-preserved avicularia are present in several specimens, especially in the holotype, and in paratypes D.23404, D.23408–09 and D.23411. Paratype D.23410 possesses lacunae which are more deeply developed than in the other specimens. D.23414 and D.23428 have two definitely paired rows of intercostal spaces in the median area ; the inner paired row is imperfectly developed in several zooecia of paratype D.23415. The occurrence of a second paired row of intercostal spaces close to the mid-line is possibly the most variable feature of the species.

All zoarial fragments are now unattached except D.29908 which encrusts a fragment of echinoid test. No young zooecia have been seen.

D.29861 is here assigned to *Pelmatopora brydonei* Lang. It compares closely with the holotype of that species ; the distal oral spines of each zooecium are undoubtedly enlarged, the lateral oral spines are visible and unmodified, the length of the zooecia is over 1·00 mm., the paired avicularia are small and oval, and there are 16–20 costae.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum*.

SPECIMENS. D.23405. Holotype—see above.

D.23402–04, D.23406–16, D 23428. Fifteen paratypes. Zoarial fragments, all now unattached. Horizon, locality and collection as for the holotype.

D.29858. Zoarial fragment, now unattached. Senonian, zone of *M. cor-anguinum*. Pit between Cliff End and Exceat Farm, near Seaford, Sussex.

D.29908. Large fragment of zoarium encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Meopham, south of Gravesend, Kent. G. E. Dibley Collection.

D.40744. Zoarial fragment encrusting a piece of echinoid test. Senonian, high in the zone of *M. cor-anguinum*. Pit 264 of Young, Coombs Pit, West Horsley, Surrey.

D.40745–46. Two unattached zoarial fragments. Senonian, zone of *M. cor-anguinum*. Locality and collection as for D.29858.

13. *Pelmatopora suffulta* (Brydone)

(Pl. 10, figs. 8, 9; Pl. 11, fig. 1; Text-figs. 70, 71)

?1911 *Cribrilina insignis* Canu, pp. 252, 280, 282, pl. 6, figs. 7–10.
1913 *Cribrilina suffulta* Brydone, p. 436, pl. 14 [*partim*—fig. 4 only].
?1916*a* *Pelmatopora insignis* (Canu) Lang, pp. 102, 104.
1916*a* *Pelmatopora suffulta* (Brydone) Lang, pp. 102, 104.
1917 *Cribrilina suffulta* Brydone: Brydone, p. 495.
1917 *Cribrilina repleta* Brydone, p. 495 [*partim*—*non* pl. 32, fig. 9].
?1919*d* *Pelmatopora insignis* (Canu): Lang, pp. 211, 218, 222, 224, text-fig. 39.
1919*d* *Pelmatopora suffulta* (Brydone): Lang, pp. 198, 211, 218, 222, 226, text-fig. 45.
1921 *Pelmatopora suffulta* (Brydone): Lang, pp. xciii, xciv.
?1921 *Pelmatopora insignis* (Canu): Lang, p. xciv.
1921 *Pelmatopora repleta* (Brydone) Lang, p. xciv [*nom. nud.*].
1922 *Pelmatopora insignis* (Canu): Lang, p. 273, text-fig. 82.
1922 *Pelmatopora suffulta* (Brydone): Lang, p. 277, text-fig. 83.
1922 *Pelmatopora repleta* (Brydone) Lang, p. 280.
1930 *Pelmatopora repleta* (Brydone): Gaster, table opp. p. 340.

LECTOTYPE. (Chosen by Lang, 1922 : 277.) The specimen figured by Brydone (1913, pl. 14, fig. 4). S.M., B.36262. A large zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Gravesend, Kent. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases; costae 14–22; pelmatidia usually present; avicularia small, distally pointed and directed, often paired, one near each proximal-lateral corner of the orifice; interzooecial secondary tissue very variably developed, the gymnocyst may be widely exposed or secondary tissue may almost fill the interzooecial valleys.

DESCRIPTION. *Zoarium* unilaminar, encrusting. *Adult zooecia* usually elongate-oval, some with almost parallel sides; occasionally very wide zooecia occur. *Orifice* rounded distally, often with straight sides, but occasionally somewhat rounded laterally, proximal margin straight, no proximal-lateral constrictions. The outline of the orifice is not obscured either by the oral spines or by secondary calcareous tissue. *Oral spine-bases* 4, possibly very occasionally 5, arranged symmetrically on the distal margin of the orifice. *Frontal shield* variably arched, either low, with a flat, fairly wide, depressed median area, or with fairly steeply sloping costae and a narrower median area; occasionally the whole frontal shield may be flat. *Costae* 14–22, each with a prominent *pelma* about half-way along its length at the edge of the median area, and, nearer the mid-line, a *pelmatidium*. *Lateral costal fusions* always occur at the level of the pelmata producing a paired row of *intercostal spaces*; occasionally, when distinct lateral costal fusions arise also at the level of the pelmatidia, a second row of intercostal spaces may occur in the median area nearer to the mid-line. The *median area of fusion* is narrow and fusiform, or wider with an almost oval outline; it is often depressed below the general surface level of the frontal shield. *Apertural bar* prominent, but not very wide, with two large pelmata, one immediately on either side of the mid-line; there is often a very slight median ridge, but no *median process*. *Gymnocyst* very variably exposed, smooth. *Interzooecial secondary tissue* may be absent in a few cases, but is commonly abundant. Where interzooecial secondary tissue is absent the gymnocyst is widely exposed laterally, and particularly proximally, to the frontal shield. Shallow *lacunae* may be developed in the interzooecial tissue.

Avicularia small, distally pointed and directed, often paired, one near each proximal-lateral corner of the orifice, but sporadic avicularia of the same type also occur frequently. The avicularian aperture is rounded proximally, where it is almost semicircular, and is constricted distally in the triangular *rostrum*. In many specimens the avicularia are worn and apparently oval—it is only when they have not been worn that they retain distally acute rostra.



FIG. 70. *Pelmatopora suffulta* (Brydone), 50466 (old collection number). Adult zooecium and three avicularia of specimen labelled by Lang " *P. insignis* (Canu)."

Ovicells apparently endozooecial. When present they are marked by a straight, low distal oral shield which is produced by the infilling with secondary tissue of the space between the distal oral spines. The orifices of ovicelled zooecia thus have an oblong outline.

Young zooecia (of S.M., B.36263) are smaller than the adult zooecia of the same zoarium, but otherwise similar, except that there are fewer costae (10–12), a second series of lateral costal fusions is not present near the mid-line, and the avicularia are not paired.

MEASUREMENTS. *Adult zooecia* Lz = 0·77 to 1·12 mm.
lz = 0·43 to 0·65 mm.
hr = 0·14 to 0·19 mm.
lr = 0·18 to 0·22 mm.

Young zooecia
(measurements based on the young zooecia of S.M., B.36263 only):
Lz = 0·53 to 0·62 mm.
lz = 0·30 to 0·35 mm.
hr = 0·12 to 0·14 mm.
lr = 0·12 to 0·14 mm.

Avicularia
among adult zooecia: Lz = 0·09 to 0·12 mm.
lz = 0·06 mm.

REMARKS. *Pelmatopora suffulta* (Brydone) and *P. repleta* (Brydone) are synonymous. *P. insignis* (Canu) has been doubtfully included in the synonymy and is discussed more fully below.

The specimens, S.M., B.36262–63, which Brydone (1913 : 436, pl. 14, figs. 4, 5) figured as *Cribrilina suffulta*, have been re-examined. The latter specimen, shown in his fig. 5, was later separated by Brydone (1917 : 495) as *Cribrilina repleta*. Lang (1916*a* : 102, 104) placed both specimens in *Pelmatopora suffulta* (Brydone). Subsequently he followed Brydone's separation of the specimens (Lang, 1922 : 280) making S.M., B.36263 the lectotype of *P. repleta* (Brydone). The dimensions and other characters of S.M., B.36262 and S.M., B.36263 are compared below.

	S.M., B.36262 Lectotype of *Pelmatopora suffulta* (Brydone)	S.M., B.36263 Lectotype of *Pelmatopora repleta* (Brydone)
Adult zooecia	Lz = 0·95 to 1·10 mm.	Lz = 0·77 to 0·90 mm.
	lz = 0·43 to 0·65 mm.	lz = 0·43 to 0·53 mm.
	hr = 0·14 to 0·19 mm.	hr = 0·17 to 0·18 mm.
	lr = 0·18 to 0·22 mm.	lr = 0·18 to 0·20 mm.
Avicularia	Lz = Not measurable	Lz = 0·12 mm.
	lz = Not measurable	lz = 0·06 mm.
Number of costae	14–20	14–16.
Oral spine-bases	4	4.
Other characters	Gymnocyst generally widely exposed; interzooecial secondary tissue absent; one (occasionally two) lateral costal fusions on each costa; avicularia usually paired, all worn	Gymnocyst usually concealed; interzooecial secondary tissue plentiful; two (occasionally only one) lateral costal fusions on each costa; avicularia usually paired, distally pointed and directed.

Although the gymnocyst is frequently well exposed in the lectotype of *P. suffulta* there are also numerous zooecia in which it is concealed or only slightly exposed. Conversely, some zooecia in the lectotype of *P. repleta* have exposed gymnocysts. Apart from differences in the amounts of interzooecial secondary tissue which may be developed the two species are clearly similar. Brydone (1917 : 495) said of his species *Cribrilina suffulta* and *C. repleta*, " The relationship between the two is, however, so very close that each is liable to display here and there a considerable measure of the distinguishing character of the other ". This statement is confirmed by an examination of the specimens which Lang assigned to *Pelmatopora suffulta* and to *P. repleta*.

One encrusting zoarial fragment, D.29907, is labelled as an ideotype of *P. suffulta* (Brydone). As in the lectotype, each of its zooecia possesses only four oral spine-bases, not " five in number normally " as suggested by Brydone (1913 : 436). Its avicularia are all worn, but Brydone, basing his opinion on the well-preserved lectotype, rightly stated that the avicularia of *suffulta* correspond " very closely in shape, size and disposition with those of *Cribrilina* [=*Pelmatopora*] *gregoryi*." Some zooecia of D.29907 are ovicelled ; the range in number of costae is 16–18, and the dimensions of the adult zooecia are as follows :

Lz = 0·87 to 0·95 mm.
lz = 0·45 to 0·50 mm.
hr = 0·15 to 0·18 mm.
lr = 0·18 to 0·21 mm.

There are slight amounts of interzooecial secondary tissue.

50466 [Old Collection number], which Lang (1922 : 274) placed in *Pelmatopora insignis* (Canu), is regarded as *P. suffulta* (Brydone), its measurements are :

Lz = 1·00 to 1·12 mm.
lz = 0·45 to 0·59 mm.
hr = 0·15 mm.
lr = 0·18 to 0·21 mm.

The avicularia of 50466 are 0·09 mm. long and 0·06 mm. wide. The main difference from D.29907 is the absence of interzooecial secondary tissue, while the avicularia are more regularly paired. Ovicelled zooecia occur occasionally ; the zoarium is encrusting. The frontal shields of the zooecia are less closely spaced and the gymnocyst generally more widely exposed than in D.29907. Unfortunately the horizon and locality of the specimen are unknown.

D.29857 and D.28275–77, labelled by Lang " *P. repleta* (Brydone) ", are also here referred to *P. suffulta*. The first is a fragmentary encrusting zoarium with a poorly preserved ancestrula and younger zooecia only. Interzooecial secondary tissue is fairly abundant, and each zooecium has about 17 costae and paired avicularia. Describing the avicularia of *P. repleta*, Lang (1922 : 280) referred to them as " sporadically distributed ". They show, however, a strong tendency to be paired, one just proximal to each proximal-lateral corner of the orifice. D.28275–77 are not well preserved, but each zooecium has apparently 16 or 17 costae. Lang (1922 : 281) also recorded as *P. repleta* one other specimen, D.28290, which has 22 costae in some zooecia and a flat frontal shield.

The amount of interzooecial secondary tissue which may be present in a given zoarium is not a useful diagnostic character. The specimens in the British Museum (Natural History), previously recorded by Lang as *P. suffulta* or as *P. repleta*, show particularly the variable amounts of tissue which may occur, both in different parts of the same zoarium and in separate zoaria of otherwise similar specimens. In their development of interzooecial secondary tissue the lectotypes of *P. suffulta* (S.M., B.36262) and *P. repleta* (S.M., B.36263) lie near the extremes of a wide variation which encompasses 50466, D.29907, D.29857, D.28275–77 and D.28290—the other characters of all these specimens being similar, or, as with the range in number of costae which may be present, having a wide variation.

Cribrilina insignis Canu has been doubtfully included in the synonymy of *P. suffulta,* though it is certainly a *Pelmatopora.* The specimen is of Rocanean age, from the type locality of Roca, Argentine. Canu (1911 : 252, pl. 6, figs. 7–10) referred to the zooecia as distinct, separated by deep furrows, large, though the dimensions given are very variable—one of the figured zooecia is 0·50 mm. wide. The range in number of costae is apparently 16–20, and the median area [="plastron" of Canu] is fusiform and " irregularly perforate ". Canu noted, however, that the " perforations " in the median area are aligned with the " prolongation of the intercostal spaces ". There are, in fact, two paired rows of intercostal spaces with others occurring occasionally nearer to the mid-line. It is not possible to discern pelmatidia in Canu's figures—they are certainly present in 50466, which Lang assigned to Canu's species, and which is here referred to *P. suffulta.* The avicularia of *Cribrilina insignis* were described by Canu as oval, not constantly placed, and with small, elliptical somewhat elongate apertures. Although very similar to *P. suffulta, P. insignis* (Canu) has narrower zooecia and smaller orifices than those of the former species.



FIG. 71. *Pelmatopora suffulta* (Brydone), D. 29907. Labelled bv Lang " Idiotype ". Parts of five adult zooecia with five worn avicularia : showing the slight amount of interzooecial secondary tissue developed.

It is unfortunate that the horizon of *Cribrilina insignis* Canu is of uncertain age (see p. 19). Feruglio (1949) gave a very full discussion of the conflicting views on the age of the type series of the Rocanean. It has been placed as low as the Cenomanian and as high as the Eocene. The presence of a species of *Pelmatopora*, apparently of the less complex type, possibly suggests a horizon in the Lower Senonian for part, at least, of the type section. However, it is unwise to attempt too firm a comparison of material from such widely separated provinces.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum* in England; Rocanean in Argentina.

SPECIMENS. 50466. Old Collections number. A well-preserved zoarial fragment encrusting a piece of echinoid test. Labelled by Lang "*Pelmatopora insignis* (Canu)". Horizon and locality unknown, possibly Senonian of southern England. Prof. Morris Collection.

D.29907. Zoarial fragment with many broken zooecia encrusting a piece of echinoid test. Labelled by Lang "Ideotype". Senonian, zone of *M. cor-anguinum*. Grays, Essex. G. E. Dibley Collection.

D.29857. Zoarial fragment with poorly preserved ancestrula and young zooecia only, encrusting a shell fragment. Recorded by Lang as *P. repleta* (Brydone). Senonian, zone of *M. cor-anguinum*. Pit in Houndean Bottom, west of Lewes, Sussex.

D.28275–77, D.28290. Four zoarial fragments now unattached. Recorded by Lang as *P. repleta* (Brydone). Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed, about 20 ft. above base of zone. Cliff End, east side of Cuckmere Haven, Sussex.

D.40736. Zoarial fragment encrusting a piece of *Inoceramus* hinge. Senonian, zone of *M. cor-anguinum* (less than 10 ft. below the "strong *cor-anguinum* tabular flint band"). Cliffs, first of Seven Sisters, Cuckmere end, east of Seaford, Sussex.

D.40737–38. Two unattached zoarial fragments. Senonian, zone of *M. cor-anguinum*. Pit between Cliff End and Exceat Farm, east of Seaford, Sussex.

D.40739–41. Three unattached zoarial fragments. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed. Cliffs, Beltout, west of Eastbourne, Sussex.

D.40742. Unattached zoarial fragment. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed. Summit of Downs, Mount Harry, near Lewes, Sussex.

OTHER MATERIAL. S.M., B.36262. Lectotype—see above.

S.M., B.36263. Large, well-preserved zoarial fragment with young and adult zooecia, some with ovicells, encrusting a piece of echinoid test. Figured by Brydone (1913, pl. 14, fig. 5) as *Cribrilina repleta*, chosen by Lang (1922 : 280) as lectotype of *Pelmatopora repleta* (Brydone). Senonian, zone of *M. cor-anguinum*. Gravesend, Kent. R. M. Brydone Collection.

14. ***Pelmatopora simplex*** Lang

(Pl. 11, figs. 2, 3 ; Text-figs. 72, 73)

1916*a*	*Pelmatopora simplex* Lang, pp. 102, 105.	
1919*d*	*Pelmatopora simplex* Lang :	Lang, pp. 210, 218, 221, 224, 226, text-fig. 46.
1921	*Pelmatopora simplex* Lang :	Lang, p. xcii.
1922	*Pelmatopora simplex* Lang :	Lang, p. 282, pl. 6, fig. 1 ; text-fig. 86.
1924	*Pelmatopora simplex* Lang :	Gaster, second of three tables opp. p. 110.
1930	*Pelmatopora simplex* Lang :	Gaster, table opp. p. 340.

HOLOTYPE. D.28281. Well-preserved zoarium, with ancestrula, encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Cliffs, east side of Old Nore Point, west of Newhaven, Sussex.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases ; Lz less than 1·00 mm. ; costae 16–18 ; a second series of pelmatidia may be developed close to the mid-line ; interzooecial secondary tissue conspicuous, with large, well-defined lacunae ; avicularia paired, one at each proximal-lateral corner of the orifice.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* oval. *Orifice* rounded distally and laterally with a straight proximal margin, without proximal-lateral constrictions ; its outline may be obscured by a high ring of secondary tissue developed particularly on the distal margin. *Oral spine-bases* 4, tending to be obscured by the ring of secondary tissue, the more distal pair slightly enlarged, otherwise unmodified and not completely immersed. *Frontal shield* flat. *Costae* 16–18, well defined, narrow, each with a *pelma* about half-way along its length, and, nearer to the mid-line, at least one, and often two, *pelmatidia*. *Lateral costal fusions* occur at the levels of the pelmata and pelmatidia, producing two paired rows of *intercostal spaces*. *Apertural bar* fairly wide, flat, or sloping down towards the proximal edge of the orifice, with two poorly developed pelmata (one near each end), and a very slight median ridge. *Gymnocyst* frequently exposed through the very conspicuous, deep, well-defined, large and irregular *lacunae* in the *interzooecial secondary tissue*. The latter is abundant, and in some cases more than fills the interzooecial valleys, overflowing on to the frontal shields.

Avicularia paired, one near each proximal-lateral corner of the orifice, small, rounded or somewhat elongate, probably distally directed, and usually deeply immersed in the interzooecial secondary tissue. *Ovicells* not seen.

Young zooecia much smaller than the adult zooecia, with smaller, but well-defined, orifices, the margins of which are not obscured by secondary tissue. Oral spine-bases four, small and unmodified. Frontal shield slightly more arched than in the adult zooecia. Costae 11, well defined, narrow, each with a pelma about half-way along its length and a pelmatidium nearer to the mid-line. Lateral costal fusions occur at the level of the pelmata only. Apertural bar very well defined, with a marked pelma near each end. The gymnocyst is exposed through large, irregular lacunae which occur frequently in the plentiful interzooecial secondary tissue. The latter tends to fill the interzooecial valleys and to encroach on to the frontal shields. Avicularia are generally paired, small, round or slightly elongate, and are probably distally directed.



FIGS. 72, 73. *Pelmatopora simplex* Lang, D. 28281. Holotype.

(72) Adult zooecium and three avicularia, showing very deep, large lacunae in the interzooecial secondary tissue.

(73) Young zooecium and avicularium. An expansion of secondary tissue partly obscures the frontal shield.

MEASUREMENTS. *Adult zooecia* Lz = 0·69 to 0·84 mm.
lz = 0·39 to 0·45 mm.
hr = 0·15 to 0·18 mm.
lr = 0·18 mm.
Young zooecia Lz = 0·50 mm.
lz = 0·24 to 0·27 mm.
hr = 0·12 mm.
lr = 0·12 to 0·15 mm.

REMARKS. The holotype and D.29859–60 have been re-examined. The measurements given by Lang (1922 : 282) for adult zooecia of *P. simplex* were slightly overestimated. D.29859 is an encrusting, unilaminar zoarium with a poorly preserved ancestrula. D.29860 is another fragment of a unilaminar encrusting zoarium, probably a young stage. Most zooecia of this specimen are small and have rather more interzooecial secondary tissue than is present in the ancestrular area of the holotype.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *G.* [*A.*] *quadrata* and zone of *O. pilula*, subzone of *E. scutata* var. *depressula* and subzone of *E. scutata* var. *cincta*.

SPECIMENS. D.28281. Holotype—see above.

D.29860. Zoarial fragment with poorly preserved ancestrula, encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, upper part of subzone of *E. scutata* var. *depressula*. Pit 4 of Gaster, west of Boundstone Lane, south of Lancing Ring, north-east of Worthing, Sussex.

D.29859. Zoarial fragment with young zooecia, encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit on East Hill, Rottingdean, east of Brighton, Sussex.

D.40743. cf. *simplex*. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 87 of Gaster, near track to Newlands Barn, north end of East Hill, Rottingdean, east of Brighton, Sussex.

D.40945. Small, well-preserved zoarial fragment with young zooecia, encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit 1 of Gaster, pit near Bridle Road, south of Lancing College and College (or Bunwell's) Farm, west of Pad Farm, Lancing, Sussex.

15. *Pelmatopora plantaris* Lang

(Pl. 11, fig. 4; Text-fig. 74)

1916*a* *Pelmatopora plantaris* Lang, pp. 102, 105.
1919*d* *Pelmatopora plantaris* Lang: Lang, pp. 199, 210, 213, 216, 218, 222, 224, text-figs. 6–8, 51.
1921 *Pelmatopora plantaris* Lang: Lang, p. xciv.
1922 *Pelmatopora plantaris* Lang: Lang, p. 288, pl. 6, fig. 4; text-fig. 89.

HOLOTYPE. D.19620. Zoarial fragment, with young zooecia, encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Wivelrod, west of Alton, Hampshire. H. O. White Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with unmodified oral spine-bases; Lz less than 1·00 mm.; costae 13–16, slightly thicker than in *P. simplex*; a second series of pelmatidia may be developed near the mid-line; interzooecial secondary tissue conspicuous with large well-defined lacunae; two avicularia to each zooecium but not usually symmetrically paired.

DESCRIPTION. *Zoarium* unilaminar, probably encrusting but now unattached. *Adult zooecia* broadly oval. *Orifice* rounded distally and laterally, with a straight proximal margin, without proximal-lateral constrictions; its outline may be obscured by secondary tissue. *Oral spine-bases* 4, unmodified, the distal pair seldom show the slight thickening which is more characteristic of the distal oral spines of *P. simplex*. *Frontal shield* flat. *Costae* 13–16 (often 14), well-defined, slightly wider than the costae of *P. simplex*. Each costa bears a *pelma* about half-way along its length, and, nearer the inner end, one (occasionally two) *pelmatidia* very close to the mid-line. *Lateral costal fusions* occur at the levels of the pelmata and the outermost pelmatidia producing two paired rows of *intercostal spaces*. *Apertural bar* not quite so wide as in *P. simplex*, with two prominent pelmata and a very slight median furrow. *Gymnocyst* extensively exposed through very conspicuous, deep, large, well-defined, irregular *lacunae* in the abundant *interzooecial secondary tissue*, which occasionally encroaches slightly on to the edges of the frontal shields of some zooecia.

Avicularia, two to each zooecium, but not symmetrically paired as in *P. simplex*, small, rounded (probably by wear), or slightly elongate and probably distally directed; not deeply immersed in the interzooecial secondary tissue. *Ovicells* not seen.



FIG. 74. *Pelmatopora plantaris* Lang, D. 19620. Holotype. Adult zooecium and paired avicularia. The proximal pair belong to adjacent zooecia.

Young zooecia are smaller than the adult zooecia and have more arched frontal shields with fewer (11) costae and less interzooecial secondary tissue; the gymnocyst is more exposed. Avicularia occur singly in the young stages, or there are two associated with each zooecium, although they are not symmetrically paired.

MEASUREMENTS. *Adult zooecia* Lz = 0·78 to 0·87 mm.
lz = 0·42 to 0·50 mm.
hr = 0·15 mm.
lr = 0·18 mm.
Young zooecia Lz = 0·50 mm.
lz = 0·31 mm.
hr = 0·12 mm.
lr = 0·12 mm.
Avicularia Not measurable.

REMARKS. In the description emphasis has been laid on the differences between *P. simplex* and *P. plantaris*. At most these differences are relatively slight, but there is also a difference in the stratigraphical occurrence of the two species. All the material assigned to *P. simplex* is from the zones of *G.* [*A.*] *quadrata* and *O. pilula*, whereas the material referred to *P. plantaris* is of *M. cor-anguinum* age.

Until more material is available from the intervening horizons it seems advisable to retain the two species as separate, basing the division principally on (i) the lack of symmetrically paired avicularia in *P. plantaris* compared with the common occurrence of evenly placed pairs in *P. simplex*, and (ii) on the smaller number of slightly wider costae in the former compared with the higher number of narrower costae in the latter.

D.28291, an encrusting zoarium, was labelled by Lang " *Pelmatopora ?plantaris* ", but it has 15–17 costae and its avicularia tend to be symmetrically paired.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum*.

SPECIMENS. D.19620. Holotype—see above.

D.28288–89. Two paratypes, zoarial fragments, now unattached. Senonian, zone of *M. cor-anguinum*, *Trochiliopora* Bed, about 20 ft. above base of zone. Cliffs, east side of Cuckmere Haven, Sussex.

?D.28291. Zoarial fragment encrusting a piece of echinoid test. Labelled by Lang " *Pelmatopora ?plantaris* ". Senonian, zone of *M. cor-anguinum*. Cliffs, westernmost of Seven Sisters, east of Cuckmere Haven, Sussex.

D.40747. Incomplete zoarium encrusting a fragment of *Inoceramus* shell. Senonian, zone of *M. cor-anguinum*, 4 ft. below the " strong *cor-anguinum* tabular flint band ". Cliffs, Beltout, west of Eastbourne, Sussex.

16. *Pelmatopora quadrivolucris* Lang

(Pl. 11, figs. 5, 6 ; Pl. 12, fig. 1 ; Text-fig. 75)

1913	*Cribrilina gregoryi* Brydone : Brydone, p. 436 [*partim*—pl. 14, fig. 3 only].
1913	*Cribrilina suffulta* Brydone, p. 436 [*partim*—pl. 14, fig. 5 only].
1916*a*	*Pelmatopora quadrivolucris* Lang, pp. 102, 105.
1917	*Cribrilina galanthis* Brydone, p. 495.
1917	*Cribrilina repleta* Brydone : Brydone, p. 495 [*partim*—pl. 32, fig. 9 only].
1919*d*	*Pelmatopora quadrivolucris* Lang : Lang, pp. 200, 213, 216, 222.
1921	*Pelmatopora quadrivolucris* Lang : Lang, p. xciii.
1922	*Pelmatopora quadrivolucris* Lang : Lang, p. 295, text-fig. 92.

HOLOTYPE. D.28907. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum*. Coombs Pit (Pit 264 of Young), West Horsley, north-east of Guildford, Surrey.

EMENDED DIAGNOSIS. *Pelmatopora* with two pairs of exposed enlarged oral spines.

DESCRIPTION. *Zoarium* unilaminar, encrusting, although it may now be unattached. *Zooecia* long, elliptical or almost parallel-sided. *Orifice* rounded distally and laterally with a straight proximal margin ; its outline is often indented by the two pairs of enlarged oral spines, and its size may be reduced by the accumulation of secondary tissue round its margin. *Oral spines* 4, symmetrically arranged in two pairs, all enlarged, and somewhat greater in size than the peg-like distal oral spines of *P. brydonei*. *Frontal shield* flat, of variable length, often just over twice as long as wide. *Costae* 14–19, well defined, each with a *pelma* about half-way along its length and a *pelmatidium* nearer the mid-line; very occasionally a second pelmatidium may occur very close to the mid-line. *Lateral costal fusions* arise at the level of the pelmata, and less frequently at the level of the pelmatidia. *Apertural bar* wide, flat, with two well-defined pelmata. *Gymnocyst* concealed by plentiful *interzooecial secondary tissue* which fills the interzooecial valleys. Slight depressions are present in the surface of the interzooecial tissue.



FIG. 75. *Pelmatopora quadrivolucris* Lang, D.28907, Holotype. Adult zooecium with paired, worn avicularia and four enlarged oral spine-bases.

Avicularia distinct, distally pointed and directed, usually paired, one near each proximal-lateral corner of the orifice, occasionally single. The avicularian aperture is constricted distally in the triangular *rostrum* and is wider, almost semicircular, proximally. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 1·00 to 1·25 mm.
lz = 0·35 to 0·60 mm.
hr = 0·15 to 0·21 mm.
lr = 0·21 to 0·27 mm.
Avicularia Lz = 0·12 to 0·15 mm.
lz = 0·07 to 0·09 mm.

REMARKS. D.28907, the holotype, has been re-examined. Five specimens from the R. M. Brydone Collection in the Sedgwick Museum, Cambridge, have also been assigned to the species; these are S.M., B.36261, S.M.,B.36363, S.M., B.71538–39 and S.M., B.75867. The first of these is a well-preserved encrusting zoarium. Part of it was figured by Brydone (1913, pl. 14, fig. 3) originally as *Cribrilina gregoryi*, and later (1917 : 495) distinguished by him as *Cribrilina galanthis*. There is no doubt that it is a specimen of *Pelmatopora quadrivolucris*. Lang (1922 : 293) provisionally included the specimen in his synonymy of *P. brydonei*, presumably basing his opinion on Brydone's figure only. Each zooecium in S.M., B.36261 has four definitely enlarged oral spines, the range in number of costae is 14–16, there is less interzooecial secondary tissue than in the holotype of *P. quadrivolucris*, and the avicularia are more markedly paired, although single avicularia are also frequent. No ovicells or young zooecia are preserved. Part of the zoarial fragment is shown here (Pl. 11, fig. 5).

Brydone (1913 : 436, pl. 14, figs. 4, 5) originally designated *Cribrilina suffulta* [=*Pelmatopora suffulta*] to include his "coarse form" shown in fig. 5. Later (1917 : 495), Brydone separated this coarse form as *Cribrilina repleta* and figured part of another specimen which he also assigned to the latter species. Of this specimen, S.M., B.36363, he stated (1917 : 495, pl. 32, fig. 9) "I have figured an exceptionally well-preserved specimen which shows that the tubercles on the anterior lip [oral spines] are typically four in number, long and tapering, but very brittle". I have re-examined this specimen, each zooecium of which has four enlarged oral spines, and confirm Lang's opinion (1922 : 295 *in synon.*) that it belongs to *Pelmatopora quadrivolucris*. The measurements and other details of the specimen are as follows:

Zooecia Lz = 1·11 to 1·25 mm.
lz = 0·50 to 0·60 mm.
hr = 0·20 to 0·21 mm.
lr = 0·26 to 0·27 mm.
Range in number of costae = 16–19.
Number of lateral costal fusions on each costa = 1 (occasionally 2).
Avicularia Lz = 0·15 to 0·16 mm.
lz = 0·07 to 0·09 mm.

It is noticeable that the dimensions of S.M., B.36363 are generally greater than those of the holotype of *P. quadrivolucris*, suggesting that the latter may be a slightly younger stage of zoarial growth. The avicularia of S.M., B.36363, which are well preserved, are distally pointed and directed, unlike the apparently oval avicularia of the holotype which is somewhat worn. Part of S.M., B.36363 is shown in Pl. 12, fig. 1.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *M. cor-anguinum.*

SPECIMEN. D.28907. Holotype—see above.

OTHER MATERIAL. S.M., B.36261. Holotype (by monotypy) of *Cribrilina galanthis* Brydone. A well-preserved zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *M. cor-anguinum.* Locality 199 of Brydone, siding east of junction, railway cutting, Basingstoke, Hampshire. R. M. Brydone Collection.

S.M., B.36363.	Zoarial fragment, now unattached, but partly embedded in a small block of chalk. The specimen figured by Brydone (1917, pl. 32, fig. 9) as *Cribrilina repleta*. Senonian, zone of *M. cor-anguinum*. Locality 559 of Brydone, excavations for tank at fold yard, one quarter mile west of Lower Grenville Copse, Soberton, Hampshire. R. M. Brydone Collection.
S.M., B.71538.	Zoarial fragment, now unattached, but partly embedded in a small block of chalk. Senonian, zone of *M. cor-anguinum*. Locality 502 of Brydone, pit in lane one quarter mile south-east of Grasmere Cottage, Easton, Hampshire. R. M. Brydone Collection.
S.M., B.71539.	Zoarial fragment, now unattached, but partly embedded in a small block of chalk. Horizon, locality and collection as for S.M., B.36363.
S.M., B.75867.	Zoarial fragment encrusting four pieces of echinoid test. Senonian, zone of *M. cor-anguinum*. Locality 328 of Brydone, Hurstbourne Priors (Beehouse Pit), Hampshire. R. M. Brydone Collection.

17. ***Pelmatopora brydonei*** Lang

(Pl. 12, fig. 3 ; Text-figs. 76–78)

1906	*non Cribrilina gregoryi* Brydone, pp. 290, 300, text-fig. 13.
1913	*non Cribrilina gregoryi* Brydone : Brydone, p. 436, pl. 14, figs. 1–3.
1916*a*	*Pelmatopora brydonei* Lang, pp. 103, 105.
1917	*non Cribrilina galanthis* Brydone, p. 495.
1919*d*	*Pelmatopora brydonei* Lang : Lang, pp. 199, 210, 213, 220, 222, 225, text-figs. 9–11, 53, 54.
1921	*Pelmatopora brydonei* Lang : Lang, p. xciii.
1922	*Pelmatopora brydonei* Lang : Lang, p. 293, text-fig. 91.
1922	*Pelmatopora pero* Lang : Lang, p. 290, pl. 6, fig. 5, text-fig. 90 [*partim*—D.29861 only].
?1958	*Pelmatopora* cf. *brydoneis* [*sic*] Lang : Didon, pp. 83, 85, text-fig. 10.

HOLOTYPE. D.23396. Small zoarial fragment, now unattached. Senonian, top of zone of *M. cor-anguinum*. Epsom, Surrey. F. Möckler Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with a pair of enlarged distal oral spines, the unmodified pair of lateral oral spines not entirely concealed by secondary thickening on the margin of the orifice.

DESCRIPTION. *Zoarium* unilaminar, encrusting, but may now be unattached. *Zooecia* elongate, elliptical, tending to be parallel-sided and somewhat pointed proximally. *Orifice* wider than high, rounded distally and laterally, with a straight proximal margin ; the outline of the orifice often indented by the enlarged distal oral spines. Two pairs of *oral spines* visible—the distal pair are enlarged and peg-like and point proximally, the lateral pair are unmodified and almost obscured by the development of secondary tissue. *Frontal shield* variable in length, usually narrow, flat. *Costae* 15–19, well defined, each with a *pelma* about half-way along its length, a *pelmatidium* near its inner end, and, occasionally, a second very small pelmatidium very close to the mid-line. *Lateral costal fusions* occur at the levels of the pelmata and the outermost pelmatidia, producing a paired row of irregular *intercostal spaces* and a variably developed second paired row closer to the mid-line *Median area of fusion* narrow. *Apertural bar* wide and flat, with obscure pelmata.



FIGS. 76, 77. *Pelmatopora brydonei* Lang, D.23396. Holotype.

(76) Very long adult zooecium with normal, enlarged distal oral spines, unmodified lateral oral spine-bases and paired avicularia near the proximal-lateral corners of the orifice. The two more heavily calcified and worn avicularia, near the proximal end of the zooecium, belong to adjacent zooecia.

(77) Dorsal surface of adult zooecium with irregularly spaced solid bosses.

Gymnocyst concealed by plentiful *interzooecial secondary tissue* which fills the interzooecial valleys and which contains slight *lacunae*.

Avicularia distinct, large, distally directed, oval, about two-thirds as wide as long; there is a strong tendency for them to occur near the proximal-lateral corners of the orifice, either singly or paired. Usually two avicularia are associated with each zooecium. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 1·00 to 1·45 mm.
lz = 0·39 to 0·56 mm.
hr = 0·18 mm.
lr = 0·21 mm.
Avicularia Lz = 0·09 mm.
lz = 0·06 mm.

REMARKS. The holotype, eight paratypes, and one other specimen, all labelled by Lang " *P. brydonei* ", have been re-examined.



(*a*) (*b*)

FIG. 78. *Pelmatopora brydonei* Lang, D. 23397. Paratype.
(*a*) Side view of part of a damaged orifice. (*b*) Front view of same orifice. *m*, hard matrix ; *o.s*, apparently unmodified " primary " oral spine ; *r.p*, rim of primary orifice ; *r.s*, rim of secondary orifice ; *s.t*, secondary tissue of enlarged oral spine.

The dorsal surface of the zoarial lamina of the holotype is covered by small, apparently solid, conical bosses which are irregularly distributed and by which the zoarium may have been attached (Text-fig. 77). The form of these dorsal bosses contrasts with the hollow dorsal " spines " developed in *P. gregoryi*, and particularly with those illustrated (Text-figs. 91, 92).

It was stated by Lang (1922 : 294), with reference particularly to D.23397, that " both the right distal apertural [oral] spine and the right secondary aviculoecium are present, the latter lying outside the former. But the primary apertural rim in this specimen is so distinct as to suggest that the whole aperture has been renewed. If this has happened, this specimen cannot be used to demonstrate that the secondary aviculoecium is a new structure, and not a development of the distal apertural spine, as has been assumed hitherto. ... If the secondary aviculoecia are rightly regarded as developments of no pre-existing structures, the species *P. brydonei* must be considered as arising from a saltation ...". Here Lang referred to the nature of the oral spines and orifices of the specimen which is shown here in side and front views respectively (Text-fig. 78*a*, *b*). From a careful examination of the specimen under high-power magnification it is apparent that the rim of the primary orifice is undamaged but partly concealed by hard matrix. Distal to the rim of the primary orifice an apparently unmodified " primary " oral spine is preserved. This structure is, however, intimately fused to a localized accumulation of secondary calcareous tissue on its distal side, here termed " secondary tissue of the enlarged oral spine ". This enlarged oral spine protrudes above the general level of the rim of the secondary orifice which is very much broken, but which normally, in undamaged zooecia, obscures the rim of the primary orifice and the unmodified oral spines. The left-hand distal oral spine has been destroyed.

I regard the smaller " primary " unmodified right-hand distal oral spine and the localized accumulation of secondary tissue on its distal side as parts of the same enlarged distal oral spine which has been broken and worn obliquely along its length. Damage to the distal end of the zooecium has largely removed secondary tissue, exposing the sharply defined distal margin of the primary orifice and the originally small base of the oral spine.

This arrangement does not necessarily indicate renewal of the orifice, but does show that the so-called " secondary aviculoecia " of Lang are, in fact, enlarged oral spines : that is, spines which are developed round the margin of the orifice and which have increased in size during the localized accumulation of secondary tissue.

It is thus incorrect to represent *P. brydonei* (*vide* Lang, 1922 : 295, text-fig. 91) as possessing two pairs of unmodified oral spines in addition to the so-called " secondary aviculoecia ". The species has only one pair of lateral unmodified oral spines and one pair of enlarged distal oral spines. (See also the general remarks on oral spines, p. 35.)

In the adult zooecia of *P. brydonei* the number of costae varies from 15 to 19 and is most often 16. The avicularia, described by Lang as " sporadic ", occur almost invariably two to each zooecium. They are often symmetrically paired, one on either side, near the proximal-lateral corners of the orifice.

D.29861, recorded by Lang (1922 : 292) as *P. pero*, is actually a specimen of *P. brydonei*. The specimen is discussed on p. 150.

STRATIGRAPHICAL DISTRIBUTION. Senonian, upper part of the zone of *M. cor-anguinum*.

SPECIMENS. D.23396. Holotype—see above.

D.23397–401, D.23429–31. Eight paratypes. Zoarial fragments, now unattached. Senonian, extreme top of the zone of *M. cor-anguinum*. Medical College Pit, Epsom, Surrey. F. Möckler Collection.

D.23325. Small zoarial fragment, now unattached. Horizon, locality and collection as for the paratypes.

D.29861. Zoarial fragment encrusting a piece of echinoid test. Recorded by Lang as *P. pero*. Senonian, high in the zone of *M. cor-anguinum*. Coombs Pit (Pit 264 of Young), West Horsley, north-east of Guildford, Surrey.

18. ***Pelmatopora coryli*** (Lang)

(Pl. 12, fig. 4; Text-fig. 79)

1916*a*	*Carydiopora coryli* Lang, p. 94.
1921	*Pelmatopora coryli* (Lang) Lang, pp. xcii, xciii [*nom. nud.*].
1922	*Pelmatopora coryli* (Lang): Lang, p. 285, pl. 6, fig. 2; text-fig. 87.
1923	*Pelmatopora corylii* [*sic*] (Lang): Waters, p. 555.
1924	*Pelmatopora coryli* (Lang): Gaster, second of three tables opp. p. 110.
1930	*Pelmatopora coryli* (Lang): Gaster, table opp. p. 340.

HOLOTYPE. D.28998. Small zoarial fragment of only eight zooecia, encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 2 of Gaster, by reservoir, near Hill Barn, north of North Lancing village, south-east of Lancing Ring, north-east of Worthing, Sussex.

EMENDED DIAGNOSIS. *Pelmatopora* with small zooecia; distal oral spines enlarged, lateral oral spines unmodified and concealed by secondary thickening on the margin of the orifice; costae 11–14; avicularia occasional, sporadic.

DESCRIPTION. *Zoarium* unilaminar, encrusting. *Adult zooecia* small, broadly oval. *Orifice* rounded distally and laterally, with a straight proximal margin, wider than high, not much obscured by secondary tissue. *Oral spines* four, but in the adult zooecia the lateral pair are unmodified and completely covered by secondary tissue. The distal pair of oral spines is enlarged. *Frontal shield* steeply sloping outwards at the margins but flat on top. *Costae* 11–14, each with a *pelma* at the middle and a *pelmatidium* near the inner end. A second pelmatidium may occur occasionally very close to the mid-line. *Lateral costal fusions* are developed at the levels of the pelmata and the outermost pelmatidia, producing two irregularly paired rows of *intercostal spaces*. *Median area of fusion* narrow. The costae are characteristically rounded where they turn down to meet the gymnocyst. *Apertural bar* fairly wide, sloping down to the proximal edge of the orifice, with two very prominent, large pelmata and a very slight median furrow. *Gymnocyst* not exposed. The interzooecial valleys are apparently lined with *secondary tissue*, the edges of this lining tissue standing up as a small but distinct ridge round each frontal shield. The "lacunae" are apparently deep hollows in the interzooecial secondary tissue; they occur only over the deepest parts of the interzooecial valleys.

Avicularia occasional, sporadic, small, rounded, possibly distally directed, frequently much obscured by secondary tissue. *Ovicells* not seen.

Young zooecia smaller than the adult zooecia, with an orifice of the same shape, but with four unmodified oral spines, none of which is obscured by secondary tissue. Frontal shield more often arched, costae 10 or 11, well defined, each with a pelma near the middle and a pelmatidium near the inner end near the mid-line. Only one lateral costal fusion occurs regularly at the level of the pelmata, though lateral fusions may occasionally be developed at the level of the pelmatidia. Apertural bar narrower than in the adult zooecia, with two very large pelmata. The gymnocyst is well exposed proximally to the frontal shield and slightly exposed laterally. Interzooecial secondary tissue, if present, is small in amount and is confined to the region of the orifice. Paired avicularia have been seen associated with only one young zooecium, where they occur one near each proximal-lateral corner of the orifice. Apart from this example, the avicularia occur sporadically in the young zoarial stages.



Fig. 79. *Pelmatopora coryli* (Lang), D.28998. Holotype. Adult zooecium with enlarged distal oral spine-bases and single avicularium near the proximal end.

Measurements. *Adult zooecia* Lz = 0·60 to 0·75 mm.
lz = 0·39 to 0·45 mm.
hr = 0·12 to 0·15 mm.
lr = 0·15 to 0·18 mm.
Young zooecia Lz = 0·51 to 0·59 mm.
lz = 0·27 to 0·33 mm.
hr = 0·09 to 0·12 mm.
lr = 0·12 to 0·15 mm.
Avicularia Not measurable.

Remarks. The holotype and three paratypes, D.28996–97 and D.28999, have been re-examined. In these specimens it is possible to find only one adult zooecium, in the holotype, in which the distal margin of the orifice is complete and unworn. Only three other adult zooecia are present in the holotype—in each case the distal margin is worn but still retains the bases of the enlarged distal oral spines which are well preserved in the unworn adult zooecium. On this evidence the specimen is here grouped with the more complex species of *Pelmatopora* which possess a pair of enlarged distal oral spines. The remaining zooecia of the holotype are smaller, and each has two pairs of unmodified oral spine-bases. It seems probable that Lang based his diagnosis and description on these young zooecia.

In paratype D.28996 only three zooecia retain complete frontal shields and all have worn or damaged distal margins. The zoarium is unilaminar and encrusting. Paired avicularia occur in two of the more complete zooecia, and there are at least two lateral costal fusions arising from each of the 12 or 13 costae which are present. The gymnocyst is variably exposed. The dimensions of this specimen are :

Lz = 0·60 to 0·69 mm.
lz = 0·39 to 0·42 mm.
hr = About 0·12 mm.
lr = 0·15 mm.

Paratype D.28997 is also worn, but there is more abundant interzooecial secondary tissue than in the holotype or in D.28996. The avicularia are generally paired. The adult zooecia are a little larger than in the holotype, with 13 or 14 costae.

D.28999 was labelled by Lang as a paratype of *P. coryli*. It is, however, very badly worn and cannot be placed with certainty in that species. It is the only specimen recorded from the zone of *Marsupites*.

Comparison has also been made between all specimens referred by Lang to *P. coryli* and all species of *Carydiopora* from which it was removed by Lang (1922). None of the remaining species of *Carydiopora* possesses pelmata or distal oral spines typical of *Pelmatopora*.

Stratigraphical distribution. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula* and subzone of *E. scutata* var. *cincta*, and ?zone of *Marsupites* (subzone unspecified).

Specimens. D.28998. Holotype—see above.

D.28996. Paratype. Worn zoarial fragment, encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 4 of Gaster, Western Pit (close to Pit 3 of Gaster), at top of Boundstone Lane, south of Lancing Ring, east of Halewick Lane, Sompting, Sussex.

D.28997. Paratype. Worn zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs, between last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.

?D.28999. Paratype. Very badly worn zoarial fragment. Senonian, zone of *Marsupites* (subzone unspecified). Cliffs, Brighton, Sussex.

19. ***Pelmatopora marsupitorum*** Lang

(Pl. 12, figs. 2, 5 ; Text-figs. 80–83)

1916*a* *Pelmatopora marsupitorum* Lang, pp. 103, 105.
1916*a* *Pelmatopora roedeanensis* Lang, pp. 103, 105.
1919*d* *Pelmatopora marsupitum* [*sic*] Lang : Lang, pp. 214, 216, 222, text-figs. 55, 56.
1919*d* *Pelmatopora roedeanensis* Lang : Lang, pp. 215, 222.
1921 *Pelmatopora marsupitum* [*sic*] Lang : Lang, pp. xcii, xciii.
1921 *Pelmatopora roedeanensis* Lang : Lang, pp. xcii, xciii.
1922 *Pelmatopora marsupitum* [*sic*] Lang : Lang, p. 297, pl. 6, fig. 6 ; text-figs. 73*b*, 93.
1922 *Pelmatopora roedeanensis* Lang : Lang, p. 299.
1924 *Pelmatopora marsupitum* [*sic*] Lang : Gaster, second of three tables opp. p. 110.
1924 *Pelmatopora roedeanensis* Lang : Gaster, second of three tables opp. p. 110.
1930 *Pelmatopora marsupitum* [*sic*] Lang : Gaster, p. 336, table opp. p. 340.
1930 *Pelmatopora roedeanensis* Lang : Gaster, table opp. p. 340.
?1958 *Pelmatopora* cf. *marsupitum* [*sic*] Lang : Didon, pp. 83, 85, text-fig. 11.

Holotype. D.28867. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites* (subzone unspecified), Brighton, Sussex.

Emended diagnosis. *Pelmatopora* with a pair of relatively short and peg-like enlarged distal oral spines ; the unmodified lateral oral spine-bases always concealed by secondary tissue ; costae 14–20.

Description. *Zoarium* unilaminar, encrusting, but may now be unattached. *Zooecia* elliptical, of variable length. *Orifice* rounded distally and laterally, with a straight proximal margin, its outline often indented by the pair of enlarged distal oral spines. *Oral spines* 4, only the enlarged distal pair visible. Compared with the enlarged distal oral spines of other species these are relatively short and peg-like, occasionally rather broad, and very occasionally tending to widen slightly at their distal ends. The unmodified pair of lateral oral spines is always obscured by secondary tissue accumulated on the margins of the orifice. *Frontal shield* flat, of variable length. *Costae* 14–20 (often 18), well defined, each with a *pelma* near the outer end, a smaller pelma towards the inner end, and often a *pelmatidium* close to the mid-line. *Lateral costal fusions* arise at the levels of the pelmata producing two paired rows of *intercostal spaces*. Occasionally a single line of intercostal spaces may be developed along the mid-line of the frontal shield when opposite intercostal spaces coalesce. Even less frequently, lateral costal fusions may arise at the levels of the pelmatidia giving a third paired row of intercostal spaces. *Median area of fusion* very narrow or absent. *Apertural bar* low and wide, with two well-defined, large pelmata and often with less obvious, smaller pelmata nearer the mid-line. The distal edge of the apertural bar is frequently indented by a minute median furrow. *Gymnocyst* frequently concealed by plentiful *interzooecial secondary tissue* which fills the interzooecial valleys and which contains irregularly elongate *lacunae*.

Avicularia large, prominent, distally pointed and directed, usually paired, one near each proximal-lateral corner of the orifice, but sporadic avicularia of the same type also occur frequently. In unworn avicularia the aperture may be either divided by a straight *transverse bar*, separating a semicircular proximal portion from a triangular *rostrum*, or there may be no transverse bar, the aperture being



80



81



82



83

FIGS. 80–83. *Pelmatopora marsupitorum* Lang.

(80) D. 39400. Normal adult zooecium with well-preserved enlarged distal oral spines and paired avicularia near the orifice. The two avicularia near the proximal end belong to adjacent zooecia.

(81) Orifices of four adult zooecia showing variation in form of enlarged distal oral spines. The variation is due largely to different degrees of wear: (*a*) D.39400; (*b*) D.39363; (*c*) D.28866, Paratype; (*d*) D.28868, holotype of *P. roedeanensis* Lang.

(82) D. 39363. Orifice with well-buttressed and thickened distal oral shield typical of ovicelled zooecia. Paired, worn, proximal-lateral avicularia are present.

(83) Avicularia. (*a*) D.39369. Somewhat worn, secondarily thickened avicularium with robust transverse bar. (*b*) D. 39400. Well-preserved stalked avicularium.

narrow and slit-like for the distal two-thirds of its length. Unworn avicularia are rare : the commonly worn avicularian aperture has an elongate-oval outline.

Ovicells apparently endozooecial. In ovicelled zooecia the space between the enlarged distal oral spines is infilled with secondary tissue which may project forwards above the distal margin of the orifice forming a prominent distal oral shield. This structure is apparently less pointed than that developed in ovicelled zooecia of *P. somptingensis* and tends to be more solidly buttressed than in that species.

MEASUREMENTS. *Zooecia* Lz = 1·00 to 1·20 mm.
lz = 0·50 to 0·57 mm.
hr = 0·18 mm.
lr = 0·21 mm.
Avicularia Lz = 0·12 mm.
lz = 0·09 mm.

REMARKS. All available specimens recorded by Lang as *P. marsupitorum* or as *P. roedeanensis* have been re-examined. Much additional material, largely from the zone of *G.* [*A.*] *quadrata*, has also been assigned to *P. marsupitorum* as here revised.

Compared with those of *P. brydonei*, the distal oral spines of the holotype of *P. marsupitorum* are more enlarged ; they show no tendency to bifurcate as in *P. gregoryi* or in *P. somptingensis*. *P. roedeanensis* is synonymous with *P. marsupitorum*. Lang distinguished the species on possible differences in zoarial growth habit only. The dimensions of the zooecia of *P. roedeanensis* fall within the range of those of *P. marsupitorum* and the two are similar in all other features.

In D.39363 several zooecia possess apparently endozooecial ovicells which are marked by prominent distal oral shields of the type described ; the orifices of these ovicelled zooecia are typically oblong.

D.39369 has well worn avicularia of a typically elongate-oval shape. One avicularium is comparatively unworn, with its stout transverse bar preserved and its aperture raised on a robust stalk, the walls of which appear to be thickened by the accretion of secondary tissue. In contrast, the occasionally preserved unworn avicularia of D.39400 apparently lack a transverse bar and their apertures are constricted distally. This form of avicularian aperture seems to be more generally typical of *P. marsupitorum*. Unworn enlarged distal oral spines occur only occasionally. They tend to be triangular in shape, the apex of the triangle pointing towards the centre of the orifice. The shape of the enlarged distal oral spines may vary considerably according to the degree of wear.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *Marsupites* to zone of *G.* [*A.*] *quadrata*, but not yet recorded from the subzone of *E. scutata* var. *cincta* in the zone of *O. pilula*.

SPECIMENS. D.28867. Holotype—see above.

D.28866. Paratype. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for the holotype.

D.28868. Holotype of *P. roedeanensis* Lang. Zoarial fragment now unattached. Horizon, locality and collection as for the holotype.

D.28282, D.28287, D.28869–70. Four zoarial fragments. Horizon, locality and collection as for the holotype.

D.29867. Zoarial fragment. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit 1 of Gaster, south of College Farm, North Lancing, Sussex.

D.29866. Zoarial fragment. Horizon and collection as for D.29867. Cliffs between last groyne east of Rottingdean Gap and Saltdean, Sussex.

?D.28283. Very worn zoarial fragment. Horizon, locality and collection as for D.29866.

D.29868–69. Two zoarial fragments labelled by Lang " *P. roedeanensis* ". Horizon and collection as for D.29867. Cliffs east of Chimney Shaft, Roedean, east of Brighton, Sussex.

D.39363, D.39408. Two zoarial fragments, the former with well-preserved ovicells. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 15 of Gaster, large pit about one-third of a mile south-south-east of Salvington Windmill and north of Hill Cottages, Salvington, Sussex.

D.39369, D.39371, D.39373, D.39376, D.39383. Five zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 20 of Gaster, lower pit on Patching Hill, just above the 200 ft. contour, north of St. John's Church, Patching, Sussex.

D.39386–88. Three zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit by track, near 200 ft. contour, south-east of Walderton, west side of Walderton Down, West Sussex.

D.39393, D.39403, D.39433–34. Four zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 26 of Gaster, South Woodleighs Pit, east of gravel pit, north-east of Woodleigh Lodge, about half a mile west of Pit 24 of Gaster, Warningcamp, north-east of Arundel, Sussex.

D.39398–99. Two zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 18 of Gaster, Old Lime Kiln, north-west corner of Munery's Copse, west of Pit 17 of Gaster, east end of Clapham Wood, Sussex.

D.39400. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 19 of Gaster, on western boundary of Church Copse, west of St. Mary's Church, Clapham, Sussex.

D.40707. Very large zoarial fragment encrusting a piece of echinoid test. Senonian, zone and subzone of *Marsupites*. Section 75 of Gaster, cliffs, east of Brighton, Sussex.

D.40708–10. Three zoarial fragments now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Section 85 of Gaster. Roedean Cliffs, east of Brighton, Sussex.

D.40711–17. Seven zoarial fragments encrusting pieces of echinoid test. Horizon, locality and collection as for D.40707.

D.40973. *Pelmatopora* cf. *marsupitorum*. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Section 113 of Gaster, cliffs, Rottingdean to Lower Bannings, east of Saltdean Cottages, Sussex.

D.41036.	Zoarial fragment encrusting large cyclostomatous polyzoan. Horizon, locality and collection as for D.40973.
D.41037.	Very large zoarial fragment encrusting part of an echinoid test. Horizon, locality and collection as for D.40973.
D.41038–40.	Three zoarial fragments encrusting pieces of echinoid test. Horizon, locality and collection as for D.40973.
D.41042.	Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.40708.
D.41047–49.	Three zoarial fragments encrusting pieces of echinoid test. Horizon, locality and collection as for D.29867.
D.41050–55.	Six zoarial fragments, now unattached. Horizon, locality and collection as for D.29867.
D.41056–57	Two zoarial fragments encrusting pieces of echinoid test Horizon, locality and collection as for D.40973.

20. ***Pelmatopora gregoryi*** (Brydone)

(Pl. 13 ; Pl. 14, figs. 1–4 ; Text-figs. 84–94)

1852	*non Semiescharipora simplex* d'Orbigny, pl. 718, figs. 1–4 ; 1853, p. 481 ; 1854, p. 1097 [*vide* Lang].
?1900	*Cribrilina* sp. : Rowe, p. 341 [*partim*—special undescribed form], [*vide* Lang].
1906	*Cribrilina gregoryi* Brydone, pp. 290, 300, text-fig. 13 [*partim*—specimens from zone of *G.* [*A.*] *quadrata* only, *caet. excl.*].
1910	*Cribrilina gregoryi* Brydone : Brydone, p. 77.
?1910	*Cribrilina* sp. : White, p. 56 [*vide* Lang].
?1910	*Cribrilina simplex* (d'Orbigny) : White, p. 55 [*vide* Lang].
?1911	*Cribrilina gregoryi* Brydone : Griffith & Brydone, p. 4.
?1912	*Cribrilina gregoryi* Brydone : Brydone, pp. 20, 53, 62, 70, 89–95.
?1912	*Cribrilina gregoryi* Brydone : White, pp. 35, 43.
1913	*Cribrilina gregoryi* Brydone : Brydone, p. 436, pl. 14, figs. 1, 2 [*partim*—*non* fig. 3].
?1913	*non Cribrilina gregoryi* Brydone : White, p. 24 [*vide* Lang].
?1913	*Cribrilina gregoryi* Brydone : White, pp. 27, 30, 32, 38 [*partim*—*vide* Lang].
1913	*Cribrilina gregoryi* Brydone : Lang, p. 171 [*partim*—specimens other than D.23963 from zone of *G.* [*A.*] *quadrata* only : *caet. excl.*].
1916*a*	*Pelmatopora gregoryi* (Brydone) Lang, pp. 103, 106.
1916*a*	*Pelmatopora bidens* Lang, pp. 103, 106.
1916*a*	*Pelmatopora lancingensis* Lang, pp. 103, 106.
1916*a*	*Pelmatopora promontoriorum* Lang, pp. 103, 106.
1916*a*	*Pelmatopora collium* Lang, pp. 103, 106.
1916*a*	*Pelmatopora saltdeanensis* Lang, pp. 103, 106.
1916*a*	*Pelmatopora ranunculoides* Lang, pp. 103, 106.
1916*a*	*Pelmatopora lacuum* Lang, pp. 103, 106.
1916*a*	*Pelmatopora gyrinoides* Lang, pp. 103, 106.
1917	*Cribrilina gregoryi* Brydone : Brydone, pp. 50, 52, 495 [*partim*].
1919*d*	*Pelmatopora gregoryi* (Brydone) : Lang, pp. 215, 221.
1919*d*	*Pelmatopora bidens* Lang : Lang, pp. 214, 221, text-figs. 61, 62.
1919*d*	*Pelmatopora lancingensis* Lang : Lang, pp. 214, 221, text-figs. 63, 64.
1919*d*	*Pelmatopora promontoriorum* Lang : Lang, pp. 214, 221, text-fig. 67.
1919*d*	*Pelmatopora collium* Lang : Lang, pp. 214, 221, text-fig. 68.
1919*d*	*Pelmatopora saltdeanensis* Lang : Lang, pp. 214, 221, 224, 226, text-figs. 65, 66.

1919*d* *Pelmatopora ranunculoides* Lang : Lang, pp. 214, 221, text-figs. 69, 70.
1919*d* *Pelmatopora lacuum* Lang : Lang, pp. 214, 221, text-fig. 71.
1919*d* *Pelmatopora gyrinoides* Lang : Lang, p. 214, text-fig. 72.
1921 *Pelmatopora gregoryi* (Brydone) : Lang, p. xci.
1921 *Pelmatopora bidens* Lang : Lang, pp. xci, xcii.
1921 *Pelmatopora lancingensis* Lang : Lang, pp. xci, xcii.
1921 *Pelmatopora promontoriorum* Lang : Lang, pp. xci, xcii.
1921 *Pelmatopora collium* Lang : Lang, p. xcii.
1921 *Pelmatopora saltdeanensis* Lang : Lang, p. xcii.
1921 *Pelmatopora ranunculoides* Lang : Lang, pp. xcii, xciii.
1921 *Pelmatopora lacuum* Lang : Lang, p. xcii.
1921 *Pelmatopora gyrinoides* Lang : Lang, p. xcii.
1922 *Pelmatopora gregoryi* (Brydone) : Lang, p. 304 [*partim—non* D. 8009, D.28537, D.21169].
1922 *Pelmatopora bidens* Lang : Lang, p. 311, pl. 6, fig. 8, text-figs. 73*f*, 97.
1922 *Pelmatopora lancingensis* Lang : Lang, p. 314, pl. 6, fig. 9 ; text-figs. 73*g*, 98.
1922 *Pelmatopora saltdeanensis* Lang : Lang, p. 316, text-fig. 99.
1922 *Pelmatopora collium* Lang : Lang, p. 319, text-fig. 100.
1922 *Pelmatopora promontoriorum* Lang : Lang, p. 320, pl. 6, fig. 10.
1922 *Pelmatopora ranunculoides* Lang : Lang, p. 321, pl. 6, fig. 11, text-fig. 101.
1922 *Pelmatopora lacuum* Lang : Lang, p. 324.
1922 *Pelmatopora gyrinoides* Lang : Lang, p. 324, pl. 6, fig. 12 ; text-fig. 102.
1924 *Pelmatopora bidens* Lang : Gaster, second of three tables opp. p. 110.
1924 *Pelmatopora lancingensis* Lang : Gaster, second of three tables opp. p. 110.
1924 *Pelmatopora collium* Lang : Gaster, second of three tables opp. p. 110.
1924 *Pelmatopora promontoriorum* Lang : Gaster, second of three tables opp. p. 110.
1924 *Pelmatopora ranunculoides* Lang : Gaster, second of three tables opp. p. 110.
1924 *Pelmatopora lacuum* Lang : Gaster, second of three tables opp. p. 110.
?1924 *Cribrilina gregoryi* Brydone : White, pp. 55, 62.
?1929 *Cribrilina gregoryi* Brydone : Brydone, pp. 9, 12, 13.
1929 *Pelmatopora gregoryi* (Brydone) : Lang, p. 437.
1930 *Pelmatopora gyrinoides* Lang : Gaster, table opp. p. 340.
1930 *Pelmatopora ranunculoides* Lang : Gaster, table opp. p. 340.
1930 *Pelmatopora bidens* Lang : Gaster, table opp. p. 340.
1930 *Pelmatopora lancingensis* Lang : Gaster, table opp. p. 340.
1930 *Pelmatopora saltdeanensis* Lang : Gaster, table opp. p. 340.
1930 *Pelmatopora collium* Lang : Gaster, table opp. p. 340.
1930 *Pelmatopora promontoriorum* Lang : Gaster, table opp. p. 340.

LECTOTYPE. (Chosen by Brydone, 1913 : 436.) S.M., B.36120. The specimen figured by Brydone (1906 : 300, text-fig. 13). Senonian, zone of *G.* [*A.*] *quadrata*. Pit 1092 of Brydone, large pit in Stakes' Lane, Upham, Hampshire. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with a pair of distally elongated, pointed, somewhat slender, enlarged distal oral spines which tend to bifurcate, the " outer horn " of the bifurcation curving away laterally and proximally ; interzooecial secondary tissue abundant, often tending to stand out as a ridge above the flat frontal shield.

DESCRIPTION. *Zoarium* unilaminar, encrusting or may now be unattached, occasionally (in parts at least) erect or cylindrical. *Adult zooecia* oval to elliptical with somewhat parallel sides. *Orifice* wider than high, rounded distally and laterally, with a straight proximal margin, its outline indented by the pair of enlarged distal oral spines. Only the distal pair of 4 *oral spines* are visible in the adult zooecia. The distal oral spines are enlarged, elongated, generally pointed and somewhat slender, lengthened in an outwardly curving fashion, or tending to bifurcate, the " outer horn " of the bifurcation curving towards the lateral or proximal margins of the orifice, and the " inner horn" only slightly developed. Other unmodified oral spines, if present, are always entirely concealed by the accumulation of secondary tissue round the margin of the orifice. *Frontal shield* of variable length, flat. *Costae* 11–20 (often 13–16), well defined, each with two *pelmata* and a *pelmatidium*. *Lateral costal fusions* occur at the levels of the pelmata, producing two paired rows of *intercostal spaces*. The *outermost intercostal spaces* vary in shape from slot-like to oval. *Median area of fusion* very narrow. *Apertural bar* wide and low, often sloping down to the proximal margin of the orifice, with conspicuous pelmata on its proximal edge. *Gymnocyst* concealed by abundant *interzooecial secondary tissue* which fills the interzooecial valleys, often standing up as ridges above the general level of the commonly flat frontal shields. Irregular *lacunae* are variably developed in the interzooecial tissue.

Avicularia prominent, distally pointed and directed, usually regularly paired, one near each proximal-lateral corner of the orifice. The avicularian aperture is almost circular proximally, narrowing distally to a triangular *rostrum*, and usually without a transverse bar ; the aperture is often raised on a stout stalk (Text-fig. 94). Occasionally sporadic avicularia of the same type also occur.

Ovicells endozooecial. In ovicelled zooecia the space between the enlarged distal oral spines is infilled with secondary tissue producing a prominent distal oral shield, which is apparently less pointed than that developed in *P. somptingensis*.

Young zooecia small, but with a relatively large orifice which is wider than high, with two pairs of oral spine-bases which are slightly enlarged but otherwise unmodified. Apertural bar more rounded and narrower than in the adult zooecia, with two very prominent pelmata and with a slight proximal bend in the mid-line. Costae 10–13, each with a pelma and, in some, an inconspicuous pelmatidium nearer to the mid-line. Frontal shield more arched than in the adult zooecia. Small, sporadic avicularia may be present in the circum-ancestrular area in which secondary interzooecial tissue is lacking. Secondary tissue may, however, encroach on to the young zooecia from the adult stages of the same zoarium.

MEASUREMENTS. *Adult zooecia* Lz = 0·75 to 1·20 mm.
lz = 0·40 to 0·50 mm.
hr = 0·15 to 0·18 mm.
lr = 0·21 mm.
Young zooecia Lz = About 0·50 mm.
lz = About 0·33 mm.
hr = 0·12 to 0·15 mm.
lr = 0·13 to 0·15 mm.

[*continued on* p. 180.



84



86

85



87



Scale as for Fig. 84.

FIGS. 84–87. *Pelmatopora gregoryi* (Brydone).

(84) S.M. B.36120. Lectotype. Adult zooecium with lengthened, enlarged distal oral spines and three avicularia surrounded by plentiful interzooecial secondary tissue with lacunae.

(85) S.M. B. 36120. Lectotype. Ancestrula with one small, rounded lateral avicularium.

(86) D.39414. Part of encrusting zoarium with plentiful interzooecial secondary tissue which is smooth with shallow lacunae. The avicularia and most of the enlarged distal oral spines are worn.

(87) D.39414. Young zooecium of the circum-ancestrular area. Two small, worn avicularia are present near the proximal-lateral corners of the orifice.



88



89



91



90



92

Figs. 88–92. *Pelmatopora gregoryi* (Brydone).

(88) Orifices of six adult zooecia showing variation in form of enlarged distal oral spines. The variation is due largely to different degrees of wear : (*a*) and (*c*) D.28947, holotype of *P. lancingensis* Lang ; (*b*), (*d*) and (*e*) D.28820, paratype of *P. promontoriorum* Lang ; (*f*) D. 28934, holotype of *P. bidens* Lang.

(89) D. 39046. Orifice of adult ovicelled zooecium with markedly thickened and projecting distal oral shield.

(90) (*a*) D. 39301. Abnormal adult zooecium produced by coalescence of buds of two adjacent rows of zooecia. The orifice of the proximally adjacent zooecium and seven worn avicularia are also shown ; (*b*) D.39301. Young zooecium from circum-ancestrular area.

(91) D.39301. Dorsal surface of normal adult zooecium with irregularly spaced small, hollow spines.

(92) D.39301. Enlarged dorsal spines developed along the line of a fracture in the basal zooecial wall.



93



94

FIGS. 93–94. *Pelmatopora gregoryi* (Brydone).

(93) D.39382. Aborted adult zooecium. *a*, avicularia; *d*, worn, enlarged distal oral spine; *f*, distorted frontal shield; *g*, very prominent gymnocyst; *o*, heavily calcified infilling of orifice; *s.t*, distal extension of tongue of interzooecial secondary tissue covering distal oral spine.

(94) D.28816. Paratype of *P. lancingensis* Lang. Unworn avicularium.

Ancestrula of lectotype Lz = 0·45 mm.
lz = 0·30 mm.
hr = 0·12 mm.
lr = 0·13 mm.
Avicularia Lz = 0·12 to 0·15 mm.
lz = 0·06 to 0·07 mm.

REMARKS. The lectotype of *Pelmatopora gregoryi* (Brydone), S.M., B.36120, is a very well preserved, unilaminar, encrusting zoarium with young zooecia. The adult zooecia possess enlarged distal oral spines of the type described. The range in number of costae is 12–15, and the outermost intercostal spaces slot-like. The avicularia are prominent, regularly paired, distally pointed and directed; the pointed rostrum may be separated from the proximal part of the aperture by a slight transverse bar which is frequently distally curved—more often there is no bar between the rostrum and the proximal part of the avicularian aperture. A few zooecia are ovicelled; in these the distal oral shield produces a markedly oblong, transversely elongate orifice. The dimensions of the lectotype are:

Adult zooecia Lz = 0·87 to 1·06 mm.
lz = 0·45 to 0·50 mm.
hr = 0·15 to 0·18 mm.
lr = 0·21 mm.

Young zooecia Lz = 0·48 to 0·50 mm.
lz = 0·32 to 0·34 mm.
hr = 0·12 to 0·14 mm.
lr = 0·13 mm.
Avicualaria Lz = 0·13 mm.
lz = 0·06 to 0·07 mm.

Pelmatopora bidens, *P. lancingensis*, *P. promontoriorum*, *P. collium*, *P. saltdeanensis*, *P. ranunculoides*, *P. lacuum* and *P. gyrinoides*, all species established by Lang, are here grouped together as synonymous with *P. gregoryi*.

Lang originally interpreted the lectotype of *P. gregoryi* (Brydone) from Brydone's 1906 and 1913 descriptions and figures. Unfortunately these do not show the form of the enlarged distal oral spines. Examination of the actual specimen shows that they are not, as Lang (1922 : 305) has stated, " very much flattened distally and distinctly bilobed ", but are of the distally elongate curved form described above. This feature, and the range in number of the costae, the form of the avicularia, and the nature and amount of secondary tissue, are all similar to those of *P. bidens* Lang, which is thus synonymous with *P. gregoryi*. In addition the other seven species of *Pelmatopora* mentioned above are all synonymous with *P. bidens* and are thus all junior synonyms of *P. gregoryi*.

P. saltdeanensis Lang is morphologically similar to *P. promontoriorum* Lang and was maintained as separate solely on supposed differences in zoarial growth habit. Similarly, Lang separated *P. lacuum* from *P. ranunculoides*.

P. gyrinoides was established by Lang as a separate encrusting species only on the evidence of a single, somewhat worn zooecium : it might equally well be a young zooecium of *P. ranunculoides*. The latter species is morphologically similar to *P. promontoriorum* : the general range in number of costae is very similar (11–16 for *ranunculoides* and 12–19 for *promontoriorum*), the zooecia of the latter are generally more slender than those of the former species, but this may be a reflection of differences in growth habit (see remarks on the Recent species *Steganoporella buskii* Harmer, p. 22).

In turn, *Pelmatopora promontoriorum* is synonymous with *P. bidens* : they have the same general range in number of costae (11–19 and 12–20 respectively) and are similar in other features.

P. collium was established by Lang on the basis of its apparently more plentiful, ridge-like interzooecial secondary tissue and on its supposedly fewer costae, but both of these features may be very variable even in a single zoarium.

Lang divided the remaining species, *P. lancingensis*, from *P. bidens* primarily on the nature of the enlarged distal oral spines. But he stated that in *P. bidens* these structures vary " much and, apparently, irregularly, in individual development " from small and peg-like to large and more elongate. In his description of *P. lancingensis*, he referred to the enlarged distal oral spines as " varying somewhat and, apparently, irregularly in amount of development, but generally large, elongate . . ". The number of costae in both species is very similar. It was suggested by Lang (1922 : 315) that the avicularia of *P. lancingensis* have " somewhat blunt apertures ", whilst those of *P. bidens* he described (1922 : 312) as having apertures

" divided by a constriction into a proximal more-or-less circular portion and a rostrum, which is more-or-less triangular, sometimes rather blunt, and sometimes rather acutely pointed ". Here Lang was evidently comparing worn and unworn avicularia. The unworn avicularia of both species are distinctly pointed and of very similar form. Apparent variation in the form of the enlarged distal oral spines is due almost entirely to different degrees of wear ; as with the avicularia, the enlarged distal oral spines of both species, when unworn, are very similar in form. The two species are therefore regarded as synonymous.

In these terms, *Pelmatopora gregoryi* is seen to be a species with a relatively wide range of morphological variation which includes the eight species discussed above, many of which have identical horizons and localities.

Many specimens from the C. T. A. Gaster Collection, particularly from the zone of *G.* [*A.*] *quadrata*, have been assigned to *Pelmatopora gregoryi* as revised here. D.39406 possesses some ovicelled zooecia with well-developed distal oral shields (Text-fig. 89). D.39414 and D.39301 are well-preserved encrusting zoarial fragments with young zooecia (Text-figs. 87, 90*b*). An abnormal growth-form of zooecium occurs in D.39301 (Text-fig. 90*a*), it is possibly produced by the coalescence of the buds of two adjacent rows of zooecia. The dorsal surface of each zooecium of D.39301 is covered by a variable number of small, hollow bosses. These may become lengthened into small, hollow spines possibly affording extra means of attachment to damaged parts of the zoarium (Text-figs. 91, 92).

Some very unusual aborted zooecia are present in D.39382 : in the figured example (Text-fig. 93), the avicularia appear to be duplicated, the orifice is almost completely sealed by secondary calcareous tissue, and the frontal shield is much distorted, with the gymnocyst showing very prominently. A worn proximally placed avicularium is also present.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *Marsupites* to *G.* [*A.*] *quadrata*.

SPECIMENS. D.28934, D.28828–35, D.28933. Holotype and nine paratypes of *P. bidens* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit 2 of Gaster, by reservoir, near Hill Barn, North Lancing, north-east of Worthing, Sussex.

D.28987. Paratype of *P. bidens* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 10 of Gaster, western pit, in lane east of Charman Dean, north of Worthing, Sussex.

D.28796, D.29873–78. Seven paratypes of *P. bidens* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 9 of Gaster, eastern pit, lane east of Charman Dean, north of Worthing, Sussex. D.28796—T. H. Withers Collection.

D.28968–70. Three paratypes of *P. bidens* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 7 of Gaster, pit in Upton Lane (Lambleys Lane), Sompting, north-east of Worthing, Sussex.

D.28804–05. Two paratypes of *P. bidens* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit on East Hill, Rottingdean, east of Brighton, Sussex.

D.21168. Paratype of *P. bidens* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit north-west of Chilton Candover, west of Alton, Hampshire. L. Treacher Collection.

D.28953–58. Six paratypes of *P. bidens* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Pit 4 of Gaster, west of Boundstone Lane, south of Lancing Ring, north-east of Worthing, Sussex.

D.28963–64. Two paratypes of *P. bidens* Lang. Horizon as for D.28953–58. Pit 5 of Gaster, Dankton Lane, north-east of Sompting Church, north-east of Worthing, Sussex.

D.7315. Large zoarial fragment labelled by Lang " *P. bidens* ". Senonian, zone of *G.* [*A.*] *quadrata*. Arreton Down, south of Newport, Isle of Wight. Miss M. Salter Collection.

D.29097. Three fragments of the same zoarium labelled by Lang " *P. bidens* ". Senonian, zone of *G.* [*A.*] *quadrata*. Portsdown, Hampshire. W. Gamble Collection.

D.23992–24001. Ten zoarial fragments labelled by Lang " *P. bidens* ". Senonian, zone of *G.* [*A.*] *quadrata*. Newhaven, Sussex. F. Möckler Collection.

D.29865. Zoarial fragment labelled by Lang " *P. bidens* ". Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Large pit on Warningcamp Hill, north-east of Arundel, Sussex.

D.28947, D.28946, D.28948–50. Holotype and four paratypes of *P. lancingensis* Lang. Senonian, zone of *O. pilula*, upper part of subzone of *E. scutata* var. *depressula*. Pit 3 of Gaster, east of Boundstone Lane, south of Lancing Ring, north-east of Worthing, Sussex.

D.28285–86. Two paratypes of *P. lancingensis* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs, east side of Rottingdean Gap, near and above the last groyne, east of Brighton, Sussex.

D.28962. Paratype of *P. lancingensis* Lang. Horizon, locality and collection as for D.28963–64.

D.28808–19, D.28923–29. Nineteen paratypes of *P. lancingensis* Lang. Horizon, locality and collection as for D.28934.

D.4337. Zoarial fragment labelled by Lang " *P. lancingensis* ". Senonian, zone of *G.* [*A.*] *quadrata*. East Harnham, south of Salisbury, Wiltshire. W. Gamble Collection.

D.29874–75. Two zoarial fragments labelled by Lang " *P. lancingensis* ". Horizon, locality and collection as for D.28953.

D.29055. Zoarial fragment with ovicelled zooecia, labelled by Lang " *P. lancingensis* ". Senonian, zone of *G.* [*A.*] *quadrata*. Locality and collection as for D.28796.

D.29871–73. Three zoarial fragments labelled " *P. lancingensis* ". Horizon, locality and collection as for D.28947.

D.28842, D.28841, D.28843–44. Holotype and three paratypes of *P. saltdeanensis* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs, between last groyne east of Rottingdean Gap and Saltdean, east of Brighton, Sussex.

D.28538. Paratype of *P. saltdeanensis* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Southern England. F. H. Butler Collection.

D.21171. Zoarial fragment labelled by Lang "*P. saltdeanensis*". Senonian, zone of *G.* [*A.*] *quadrata*. Compton, south-west of Winchester, Hampshire. L. Treacher Collection.

D.28824, D.28825–27. Holotype and three paratypes of *P. collium* Lang. Horizon, locality and collection as for D.28934.

D.28823. Paratype of *P. collium* Lang. Horizon, locality and collection as for D.28842.

D.28284. Zoarial fragment labelled by Lang "*P. collium*". Horizon, locality and collection as for D.28842.

D.28930, D.28836–40, D.28849–52, D.28871–73, D.28931–32. Holotype and fourteen paratypes of *P. promontoriorum* Lang. Horizon, locality and collection as for D.28934.

D.28959–61. Three paratypes of *P. promontoriorum* Lang. Horizon, locality and collection as for D.28953.

D.28951–52. Two paratypes of *P. promontoriorum* Lang. Horizon, locality and collection as for D.28947.

D.28806–07, D.28874, D.28877. Four paratypes of *P. promontoriorum* Lang. Horizon, locality and collection as for D.28842.

D.28876. Paratype of *P. promontoriorum* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Cliffs, east side of Telscombe Cliff Staircase, east of Brighton, Sussex.

D.28820–22. Three paratypes of *P. promontoriorum* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Cliffs, east of Old Nore Point, west of Newhaven, Sussex.

D.28967. Paratype of *P. promontoriorum* Lang. Horizon, locality and collection as for D.28968.

D.28972. Paratype of *P. promontoriorum* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 8 of Gaster, between Upton Lane (Lambleys Lane) and east of Charman Dean, north of Broadwater, Worthing, Sussex.

D.28856, D.28857–59, D.28845–48. Holotype, three paratypes and four zoarial fragments of *P. ranunculoides* Lang. Horizon, locality and collection as for D.28842.

D.28864. Paratype of *P. ranunculoides* Lang. Horizon, locality and collection as for D.28934.

D.28860. Paratype of *P. ranunculoides* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Upper part of pit on East Hill, Rottingdean, east of Brighton, Sussex.

D.28861. Paratype of *P. ranunculoides* Lang. Senonian, zone of *Marsupites* [subzone unspecified]. Brighton, Sussex.

D.28875. Zoarial fragment labelled by Lang "*P. ranunculoides*". Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Cliffs, west side of Saltdean Bottom, east of Brighton, Sussex.

D.28279. Zoarial fragment labelled by Lang " *P. ranunculoides* ". Senonian, zone of *G.* [*A.*] *quadrata*. Cliffs, between Rottingdean and Saltdean, east of Brighton, Sussex.

D.28280. Zoarial fragment labelled by Lang " *P. ranunculoides* ". Senonian, zone of *G.* [*A.*] *quadrata*. Saltdean, east of Brighton, Sussex.

D.10995. Two fragments of the same zoarium labelled by Lang " *P. ranunculoides* ". Rottingdean, east of Brighton, Sussex. H. D. Schloss Collection.

D.28862, D.28863, D.28938–43, D.28935–37, D.28853–55. Holotype and seven paratypes of *P. lacuum* Lang, and six zoarial fragments labelled by Lang " *P. lacuum* ". Horizon, locality and collection as for D.28934.

D.28270. Holotype of *P. gyrinoides* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Saltdean, west of Newhaven, Sussex.

D.39340–41, D.39344, D.39348, D.39351–52, D.39357, D.39359–60, D.39366, D.39368, D.40057. Twelve zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 15 of Gaster, large pit about one-third of a mile south-south-east of Salvington Windmill, north of Hill Cottages, Salvington, Sussex.

D.9425. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. Locality unknown. Caroline Birley Collection.

D.19564. Zoarial fragment—specimen presented by H. O. White, said to be identical with that described by him as a *Cribrilina*, resembling the sole of a hob-nailed boot (White, 1910 : 56). Senonian, zone of *G.* [*A.*] *quadrata*. Southampton Waterworks Pit, Otterbourne, south of Winchester, Hampshire. H. O. White Collection.

D.39379, D.39382. Two zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 20 of Gaster, lower pit on Patching Hill, north of St. John's Church, just above the 200 ft. contour, Patching, Sussex.

D.39389. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. Pit by track, near 200 ft. contour, south-east of Walderton, west side of Walderton Down, West Sussex.

D.39401, D.40058. Two zoarial fragments. Horizon, locality and collection as for D.28796.

D.39405, D.40059–63. Six zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. South Woodleighs Pit, east of gravel pit, north-east of Woodleighs Lodge, about half a mile west of Pit 24 of Gaster, Warningcamp, north-east of Arundel, Sussex.

D.39406. Zoarial fragment, some zooecia ovicelled. Senonian, zone of *G.* [*A.*] *quadrata*. Labelled as " Material from Pits 17, 18, 19, 24 and 32 " of Gaster—all in the zone of *G.* [*A.*] *quadrata*.

D.39412, D.40064. Two zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata*. Labelled as " Material from Pits 8, 11, 15, 16, 17 and 23 " of Gaster—all in the zone of *G.* [*A.*] *quadrata*.

D.39414–15, D.40065–66. Four zoarial fragments. Senonian, ?zone of *G.* [*A.*] *quadrata*. Locality uncertain.

D.39301. Zoarial fragment with some aborted zooecia. Senonian, very probably zone of *G.* [*A.*] *quadrata*. Specimen marked " W.p." ?Whiteparish, eight miles south-east of Salisbury, Wiltshire. H. P. Blackmore Collection.

D.40681. Zoarial fragment, now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Locality and collection as for D.28947.

D.40682. Zoarial fragment, now unattached. Horizon, locality and collection as for D.28934.

D.40683–87. Five zoarial fragments, now unattached, the latter with ovicelled zooecia. Horizon, locality and collection as for D.28972.

D.40688–95. Eight zoarial fragments, now unattached. Horizon, locality and collection as for D.40682.

D.40696. Zoarial fragment, now unattached. Horizon, locality and collection as for D.40688.

D.40697. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula*. Section 113 of Gaster, cliff, Rottingdean to Lower Bannings, east of Saltdean Cottages, Sussex.

D.40698–99. Two zoarial fragments, now unattached. Senonian, zone and subzone of *Marsupites*. Section 75 of Gaster, cliff and foreshore east of Brighton at Black Rock, Sussex.

D.40700. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Section 114 of Gaster, cliff, Lower Bannings to Old Nore Point, near Brighton, Sussex.

D.40701. Zoarial fragment, now unattached. Horizon, locality and collection as for D.40700.

D.40702. Zoarial fragment now unattached. Horizon, locality and collection as for D.40697.

D.40703, D.40704–05. Zoarial fragment encrusting a piece of echinoid test and two zoarial fragments now unattached. Horizon, locality and collection as for D.40682.

D.40706. Zoarial fragment, now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 87 of Gaster, near cart-track to Newlands Barn, north end of East Hill, Rottingdean, Sussex.

D.40797–802. Six zoarial fragments, now unattached. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 10 of Gaster, pit adjoining Charman Dean Lane, on east side, and west of Pit 9 of Gaster, north of Broadwater and Worthing, Sussex.

D.40803–54. Fifty-two zoarial fragments, now unattached. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 12 of Gaster, large pit north of Pumping Station, top of Waterworks Lane, north-east of " The Warren ", north of Broadwater and Worthing, Sussex.

D.40855. Zoarial fragment, now unattached, partly embedded in small block of chalk. Senonian, zone of *G.* [*A.*] *quadrata*. Pit south of Beech Copse, Michelgrove, Sussex.

D.40856. Zoarial fragment, now unattached. Horizon, locality and collection as for D.28968.

D.40857–59, D.40930. Four zoarial fragments, now unattached. Senonian, zone of *G.* [*A.*] *quadrata*. Large pit about one-third of a mile south-south-east of Salvington Windmill, and north of Hill Cottages, Salvington, Sussex.

D.40946. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.40706.

D.40947. Zoarial fragment encrusting three pieces of echinoid test. Horizon, locality and collection as for D.40697.

D.40948. *Pelmatopora* cf. *gregoryi*. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.40697.

D.40949. Small zoarial fragment encrusting part of a cheilostome polyzoan. Horizon, locality and collection as for D.28934.

D.40950–51. *Pelmatopora* cf. *gregoryi*. Two zoarial fragments, now unattached. Horizon, locality and collection as for D.28934.

D.40952–53. *Pelmatopora* cf. *gregoryi*. Two zoarial fragments, now unattached. Horizon, locality and collection as for D.40697.

D.40955–67. Thirteen zoarial fragments, now unattached. Horizon, locality and collection as for D.28934.

D.40968–71. Four zoarial fragments, now unattached. Horizon, locality and collection as for D.28953.

D.40972. Zoarial fragment, now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 25 of Gaster, small roadside pit, just over half a mile south of Burpham, one-third of a mile north of South Woodleighs Pit (26 of Gaster), west-north-west of Warningcamp Hill, north-east of Arundel, Sussex.

D.40974. Incomplete erect, cylindrical zoarium. Horizon, locality and collection as for D.28934.

D.40975. Zoarial fragment, now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 5 of Gaster, locality and collection as for D.28963, see above.

D.40976. Zoarial fragment, now unattached. Horizon, locality and collection as for D.28953.

D.40977–81. Five zoarial fragments, now unattached. Horizon, locality and collection as for D.28947.

D.40982–90. Nine zoarial fragments, now unattached. Horizon, locality and collection as for D.28934.

D.40991–92, D.40993. Two zoarial fragments encrusting pieces of echinoid test, and a zoarial fragment, now unattached. Horizon, locality and collection as for D.28953.

D.40994–41018. Twenty-five zoarial fragments, now unattached. Horizon, locality and collection as for D.28934.

D.41019. Zoarial fragment, now unattached, some zooecia with sealed orifices. Horizon, locality and collection as for D.28934.

D.41020–29. Ten zoarial fragments, now unattached. Horizon, locality and collection as for D.28934.

D.41030–35. Six zoarial fragments, now unattached. Horizon, locality and collection as for D.28953.

D.41041. Zoarial fragment with some ovicelled zooecia, fragment now unattached. Horizon, locality and collection as for D.28934.

D.41043. *Pelmatopora* cf. *gregoryi*. Worn, large zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Section Z of Gaster, Swansbourn Lake Section, Arundel Park, Arun Gap, Sussex.

D.41044. Zoarial fragment, now unattached, very well preserved. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Locality and collection as for D.28963.

D.41045–46. Two zoarial fragments encrusting the same piece of echinoid test. Horizon, locality and collection as for D.40700.

D.41058. Very well preserved, erect, cylindrical zoarial fragment. Horizon, locality and collection as for D.28934.

D.41059. Well-preserved, large, zoarial fragment, now unattached. Horizon, locality and collection as for D.28953.

D.41060–61. Two zoarial fragments encrusting pieces of echinoid test. Horizon, locality and collection as for D.40697.

D.41062. Well-preserved zoarial fragment, now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Section 142 of Gaster, cliff, Old Nore Point to Inn on west side of entrance to Newhaven Harbour, Sussex.

D.41063. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for D.40697.

OTHER MATERIAL

S.M., B.36120. Lectotype—see above.

21. ***Pelmatopora somptingensis*** Lang

(Pl. 14, figs. 5, 6; Pl. 15, figs. 1, 2, 4; Text-figs. 95–99)

1906 *non Cribrilina gregoryi* Brydone, pp. 290, 300, text-fig. 13.
1913 *Cribrilina gregoryi* Brydone: Lang, p. 171 [*partim*—D.23963 only].
1916*a* *Pelmatopora somptingensis* Lang, pp. 103, 106.
1916*a* *Pelmatopora danktonensis* Lang, pp. 103, 106.
1919*d* *Pelmatopora somptingensis* Lang: Lang, pp. 214, 221, 224, text-figs. 57–58.
1919*d* *Pelmatopora danktonensis* Lang: Lang, pp. 215, 221.
1921 *Pelmatopora somptingensis* Lang: Lang, p. xci.
1921 *Pelmatopora danktonensis* Lang: Lang, p. xcii.
1922 *Pelmatopora somptingensis* Lang: Lang, p. 300, text-figs. 73*c*, 94.
1922 *Pelmatopora danktonensis* Lang: Lang, p. 309, text-figs. 73*e*, 96.
1924 *Pelmatopora somptingensis* Lang: Gaster, second of three tables opp. p. 110.
1930 *Pelmatopora danktonensis* Lang: Gaster, table opp. p. 340.

HOLOTYPE. D.28762. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *G.* [*A.*] *quadrata*. Sompting, Sussex. T. H. Withers Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with a pair of laterally flattened, distally expanded, enlarged distal oral spines which have a marked tendency to bifurcate.

DESCRIPTION. *Zoarium* unilaminar, encrusting, or now unattached. *Adult zooecia* elliptical or with almost parallel sides. *Orifice* wider than high, distally and laterally rounded, with a straight proximal margin, the outline much indented by the pair of enlarged distal oral spines. *Oral spines* 4, only the distal pair visible; they are much enlarged, laterally flattened, expanded distally, and have a marked tendency to bifurcate—the " outer horn " of the bifurcation often considerably produced towards the proximal-lateral margins of the orifice. The unmodified lateral oral spines are always concealed by secondary tissue round the margin of the orifice. *Frontal shield* of variable length, apparently flat. *Costae* 14–22 (often 16 or 17), well spaced and well defined, each with a larger outer and a smaller inner *pelma*, and often with a *pelmatidium* close to the mid-line. *Lateral costal fusions* occur at the levels of the pelmata, producing two paired rows of *intercostal spaces*. *Apertural bar* wide, low, sloping down distally to the orifice and with obscure pelmata. *Gymnocyst* concealed by plentiful *interzooecial secondary tissue* which fills the interzooecial valleys. Slight, shallow, irregularly elongate depressions occur in the surface of the interzooecial tissue.

Avicularia large, prominent, distally pointed and directed, regularly paired, one near each proximal-lateral corner of the orifice. The form of the avicularian aperture is variable; it may be slit-like, as in D.39361, widely oval but pointed distally, oval proximally and slit-like distally, as in D.39343, or very occasionally, with a *transverse bar* dividing the aperture into an approximately semicircular proximal portion and a triangular *rostrum*. The various avicularian apertures are often raised on robust stalks of variable length (Text-fig. 98).

Ovicells apparently endozooecial. In ovicelled zooecia, the space between the enlarged distal oral spines is infilled with secondary calcareous tissue producing a prominent, somewhat pointed distal oral shield.

Young zooecia smaller than the adult and less complex. Very young zooecia close to the ancestrula have typically well-arched frontal shields with somewhat flattened median areas. The enlarged distal oral spines are more peg-like rather than expanded as in the adult zooecia. Costae 10–12, even in the smallest zooecia, each with a pelma about half-way along its length. It is difficult to determine whether pelmatidia are present—possibly they occur very close to the mid-line. Apertural bar relatively very wide. The gymnocyst is possibly revealed in small, deep lacunae in the plentiful interzooecial secondary tissue. Avicularia are apparently already paired in the very young stages of zoarial growth, one on either side of each zooecium, but they are not definitely placed near the proximal-lateral corners of each orifice.

Less young zooecia are larger than those immediately surrounding the ancestrula, but otherwise generally similar, the avicularia tend to be more definitely paired near the proximal-lateral corners of the orifice, and the enlarged distal oral spines are larger with a definite tendency to expand at their distal ends.



95



96



97

FIGS. 95–97. *Pelmatopora somptingensis* Lang.

(95) D.28747. Paratype. (*a*) Young zooecium and four worn avicularia. (*b*) Adult zooecium and four worn avicularia; the enlarged distal oral spines are very worn.

(96) D.28762. Holotype. Adult zooecium with well-preserved distal oral spines and one worn and three unworn avicularia. The two proximal avicularia belong to adjacent zooecia.

(97) Orifices of seven adult zooecia of several zoaria. The variation in form of the enlarged distal oral spines is due largely to different degrees of wear. (a^{1-2}) D.28762, holotype, front and oblique side views; (*b, f*) D. 39346; (*c, g*) D. 39343; (*d*) D.28762, holotype; (*e*) D.39338.



98



99

FIGS. 98–99. *Pelmatopora somptingensis* Lang.

(98) Unworn avicularia. (*a, b*) D.39361 ; (*c, d*) D. 39343 ; (*e, f*) D. 28770, paratype.

(99) D.28770. Paratype. (*a*) Front view and (*b*) side view of parts of two adult, ovicelled zooecia with shelf-like distal oral shields produced by the infilling of the space between the distal oral spines with secondary calcareous tissue.

MEASUREMENTS. *Adult zooecia* Lz = 1·00 to 1·15 mm.
lz = 0·45 to 0·55 mm.
hr = 0·18 to 0·21 mm.
lr = 0·21 to 0·24 mm.
Avicularia Lz = 0·15 to 0·20 mm.
lz = 0·07 to 0·09 mm.
Young zooecia
(Outer edge of circum-ancestrular area)
Lz = 0·65 to 0·78 mm.
lz = 0·39 to 0·45 mm.
(Very close to the ancestrula)
Lz = 0·50 to 0·54 mm.
lz = 0·33 to 0·35 mm.

REMARKS. The holotype and all available specimens identified by Lang as *P. somptingensis* have been re-examined. *P. danktonensis* Lang is synonymous with *P. somptingensis* Lang. The holotype of *P. danktonensis*, D.23963, has all the features typical of the worn examples of *P. somptingensis*. The range in number of costae in D.23963 is 13–19; the enlarged distal oral spines, where preserved, are lengthened distally and are relatively broad, the position and form of the avicularia compares closely with those in *P. somptingensis*, and the stratigraphical distribution is well within the range of *P. somptingensis*.

In *P. somptingensis* the avicularian aperture varies in form as described above. It is significant that the avicularia, where unworn, are raised on robust stalks so that the apertures slope upwards distally and are raised above the general level of the zoarium. They are thus particularly exposed to wear, and may often be reduced to the apparently oval form which the avicularia of most slightly abraded specimens appear to have, and which is almost certainly produced by abrasion wearing down the hollow, oval avicularian stalks (e.g., the worn avicularia shown in Text-fig. 95*a*, *b*).

The enlarged distal oral spines of *P. somptingensis* may show a considerable variety of shape and size, largely dependent on the amount of abrasion they have undergone. Text-fig. 97*a–g* illustrates a series of orifices from several zoaria: the variations in form are due almost entirely to different degrees of wear, but the enlarged distal oral spines all tend to be distally extended and bifurcated, contrasting with the more narrowly elongate and more pointed enlarged distal oral spines of *P. gregoryi*, and with the smaller, peg-like enlarged distal oral spines of *P. marsupitorum*.

The description, given above, of very young zooecia and zooecia at the periphery of the circum-ancestrular area is based primarily on D.28747. Although Lang (1922 : 301) has described its ancestrula, it is not possible to determine its form definitely. The specimen is well preserved, particularly in the circum-ancestrular area, but the smallest visible zooecium is considerably damaged and further obscured by two somewhat larger zooecia which stand up above the general level of the zoarium.

D.29064 is an extremely large, somewhat worn, encrusting zoarium of *P. somptingensis* with at least one abnormal zooecium produced, apparently, by the coalescence of the buds of two adjacent rows of zooecia. It is similar in general form to that figured from D.39301—a specimen of *P. gregoryi.*

A large number of specimens have been assigned to *P. somptingensis*. The species is particularly abundant in the material collected by Gaster from the zone of *G.* [*A.*] *quadrata.*

In paratype D.28770, and in D.39343, ovicelled zooecia with prominent distal oral shields are present. The variations shown by D.39343 in the number of costae in individual zooecia, in the shape of the zooecia, and in the form of the enlarged distal oral spines, are fairly typical of the species as a whole. Well-preserved oral spines and avicularia occur particularly in the following specimens: D.39331, D.39335, D.39338, D.39346, D.39349, D.39350, D.39361, D.39372, D.39377, D.39407, D.39409, and D.39413. The enlarged distal oral spines of D.39377 have an extremely wide bifurcated form which approaches that typical of the distal oral spines of *P. palmata.*

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula* (upper part) into the zone of *G.* [*A.*] *quadrata* (especially abundant in the latter zone).

SPECIMENS. D.28762. Holotype—see above.

D.28101–09, D.28747–61, D.28763–75. Thirty-seven paratypes. Zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 7 of Gaster, Upton Lane (Lambley's Lane), north-west of Sompting, Sussex. [The same locality, horizon and collection as the holotype.]

D.28776–86, D.28788–94, D.28797, D.28979–85. Twenty-five paratypes. Zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 9 of Gaster, eastern pit in lane on eastern side of Charman Dean, north of Broadwater, Sussex. T. H. Withers Collection [except D.28979–85, C. T. A. Gaster Collection].

D.28798–801, D.28988–92. Nine paratypes. Zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 10 of Gaster, western pit in lane on eastern side of Charman Dean, north of Broadwater, Sussex. D.28798–801, T. H. Withers Collection; D.28988–92, C. T. A. Gaster Collection.

D.4338, D.29062. Two zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* East Harham, south of Salisbury. W. Gamble Collection.

D.29056–58. Three zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 8 of Gaster, about half-way between Upton Lane (Lambley's Lane), Sompting, and the lane east of Charman Dean, north of Broadwater, Sussex.

D.39331. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 12 of Gaster, large pit north of Pumping Station, top of Waterworks Lane, north-east of "The Warren", north of Broadwater and Worthing, Sussex.

D.39332–37, D.39465. Seven zoarial fragments. Horizon, locality and collection as for D.28798.

D.39338, D.39342–43, D.39345–47, D.39349–50, D.39353–54, D.39358, D.39361–62, D.39364–65, D.39367, D.39409, D.39466, D.40054–56. Twenty-one zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 15 of Gaster, large pit about one-third of a mile south-east of Salvington Windmill, and north of Hill Cottages, Salvington, Sussex.

D.23963. Holotype of *P. danktonensis* Lang. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula.* Pit 5 of Gaster, pit in Dankton Lane, north of Sompting Church, north-east of Worthing, Sussex. W. D. Lang Collection.

D.29862. Zoarial fragment labelled by Lang "*P. danktonensis*". Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta.* Cliffs, west of Broadwater and Castle Hill, Newhaven, Sussex.

D.29877–78, D.29864. Three zoarial fragments, the last labelled by Lang "*P. danktonensis*". Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula* (upper part). Pit 2 of Gaster, pit by reservoir, near Hill Barn, North Lancing, Sussex.

D.29863. Zoarial fragment labelled by Lang "*P. danktonensis*". Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *depressula.* Pit 4 of Gaster, pit west of Boundstone Lane, south of Lancing Ring, north-east of Worthing, Sussex.

D.39372, D.39374–75, D.39377, D.39380–81*a*, *b*, D.39384–85, D.39467–68. Fragments of eleven zoaria. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 20 of Gaster, lower pit on Patching Hill, north of St. John's Church, Patching, Sussex.

D.39391, D.39469. Two zoarial fragments encrusting pieces of echinoid test. Horizon, locality and collection as for D.28101.

D.39394, D.39470–72. Four zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 44 of Gaster, by Mile Oak Road, about one-third of a mile north-east of Windmill Inn, Southwick, one furlong north of Upper Shoreham Road, Sussex.

D.39395–96, D.40129, D.40131. Four zoarial fragments. Horizon, locality and collection as for D.29056.

D.39397, D.39473–75. Four zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 18 of Gaster, Old Lime Kiln, north-west corner of Munery's Copse, west of Pit 17 of Gaster, and east end of Clapham Wood, Sussex.

D.39402, D.39476–78, D.40067–73, D.40087–97, D.40109–10, D.40111–24, D.40132. Thirty-nine zoarial fragments. Horizon, locality and collection as for D.28776.

D.39404, D.39479, D.40050–53. Six zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 26 of Gaster, South Woodleighs Pit, east of gravel pit, north-east of Woodleighs Lodge, about half a mile west of pit 24 of Gaster, Warningcamp, north-east of Arundel, Sussex.

D.39407. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata.* From material labelled "Pits 17, 18, 19, 24 and 32" of Gaster [all in the zone of *G.* [*A.*] *quadrata*].

D.39411. Zoarial fragment with some ovicelled zooecia. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 35 of Gaster, small pit, north of Rake Mead Plantation, Church Hill Shaw, about half a mile north-west of Roger's Farm, and west of South Park, just above 400 ft. contour, Findon, Sussex.

D.39413. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. From material labelled " Pits 8, 11, 15, 16, 17, and 23 " of Gaster. [All in the zone of *G.* [*A.*] *quadrata*.]

D.40074–78, D.40104, D.40081–86, D.40079–80. Fourteen zoarial fragments. Horizon locality and collection as for D.28101.

D.40106–07. Two zoarial fragments. Horizon, locality and collection as for D.28101.

D.40098–103, D.40105, D.40125–28, D.40130, D.39450–53. Sixteen zoarial fragments. Horizon, locality and collection as for D.28798. D.39450–53, T. H. Withers Collection.

D.40108. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 23 of Gaster, near Angmering Park Cottages, Angmering, Sussex.

D.39435–49. Fifteen zoarial fragments. Horizon and locality as for D.28101. T. H. Withers Collection.

D.29064. Zoarial fragment. Senonian, zone uncertain, very probably zone of *G.* [*A.*] *quadrata*. South of England. F. H. Butler Collection.

D.39454–63. Ten zoarial fragments. Horizon, locality and collection as for D.29877.

D.39464. Zoarial fragment. Horizon, locality and collection as for D.28101.

D.29876. Zoarial fragment. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit by roadside between Burpham and Arundel, and west-north-west of Warningcamp Hill, Sussex.

D.40718–19. Two zoarial fragments encrusting pieces of echinoid test. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Locality as for D.29877.

D.40720. Zoarial fragment, now unattached. Horizon and locality as for D.28101.

D.40721. Zoarial fragment, now unattached. Horizon and locality as for D.28776.

D.40748–56. Nine zoarial fragments, now unattached. Horizon, locality and collection as for D.28798.

D.40757–85. Twenty-nine zoarial fragments, now unattached. Horizon, locality and collection as for D.29056.

D.40786–88. Three zoarial fragments now unattached. Horizon, locality and collection as for D.28798.

D.40789–96. Eight zoarial fragments, now unattached. Horizon, locality and collection as for D.39331.

D.40954. *Pelmatopora* cf. *somptingensis*. Zoarial fragment, now unattached. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta*. Pit 25 of Gaster, small roadside pit, just over half a mile south of Burpham, one-third of a mile north of Pit 26 of Gaster, west-north-west of Warningcamp Hill, north-east of Arundel, Sussex.

22. *Pelmatopora palmata* Lang

(Pl. 15, figs. 3, 5, 6; Text-figs. 100–102)

1916*a* *Pelmatopora palmata* Lang, pp. 103, 107.
1916*a* *Pelmatopora damicornis* Lang, pp. 103, 107.
1916*a* *Pelmatopora* (*Cribrilina*) *gregoryi* (Brydone): Lang, p. 106 [*partim*—D.8009, D.28537, D.21169 only].
1919*d* *Pelmatopora palmata* Lang: Lang, pp. 199, 214, 221, text-figs. 12–14, 59, 60.
1919*d* *Pelmatopora damicornis* Lang: Lang, pp. 215, 221.
1921 *Pelmatopora palmata* Lang: Lang, p. xci.
1921 *Pelmatopora damicornis* Lang: Lang, p. xci.
1922 *Pelmatopora palmata* Lang: Lang, p. 306, pl. 6, fig. 7; text-figs. 73*d*, 96.
1922 *Pelmatopora damicornis* Lang: Lang, p. 307.
1922 *Pelmatopora gregoryi* (Brydone): Lang, p. 304 [*partim*—D.8009, D.28537, D.21169 only].
1924 *Pelmatopora palmata* Lang: Gaster, second of three tables opp. p. 110.
1930 *Pelmatopora palmata* Lang: Gaster, table opp. p. 340.

HOLOTYPE. D.8010. Zoarial fragment, now unattached, Senonian, zone of *G.* [*A.*] *quadrata*. Winchester, Hampshire. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Pelmatopora* with a pair of enlarged distal oral spines which are very broad distally and distinctly bilobed.

DESCRIPTION. *Zoarium* uni- or bilaminar, encrusting or now unattached. *Zooecia* elliptical, the sides tending to be parallel. *Orifice* wider than high, distally and laterally rounded, with a straight proximal margin, its outline very much indented by the pair of enlarged distal oral spines. *Oral spines* 4 only the distal pair visible; these are much enlarged, distally expanded and distinctly bilobed. The unmodified pair of lateral oral spines is always concealed by secondary tissue accumulated round the margin of the orifice. *Frontal shield* of variable length, apparently fairly flat. *Costae* 16–20 (often 18 or 19), well defined, each with a larger outer and a smaller inner *pelma*, and usually with a *pelmatidium* near the mid-line. *Lateral costal fusions* arise at the levels of the pelmata producing two paired rows of *intercostal spaces*. *Apertural* bar wide, low, with obscure pelmata; the apertural bar slopes down slightly towards the proximal edge of the orifice. *Gymnocyst* entirely concealed by *interzooecial secondary tissue* which fills the interzooecial valleys and which contains irregularly elongate shallow depressions.

Avicularia prominent, large, distally pointed and directed, regularly paired, one near each proximal-lateral corner of the orifice. Well-preserved avicularia lack transverse bars, the aperture is semicircular proximally with a distally pointed *rostrum*, the apertures of many avicularia are raised on stout stalks above the general level of the zoarium. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 1·00 to 1·25 mm.
lz = 0·42 to 0·50 mm.
hr = 0·18 mm.
lr = 0·21 mm.
Avicularia Lz = 0·15 mm.
lz = 0·06 mm.



100



102



101

FIGS. 100–102. *Pelmatopora palmata* Lang.

(100) D.8010. Holotype. Adult zooecium and four worn avicularia; those near the proximal end belong to adjacent zooecia. The distal oral spines are typically flat and bilobed, that on the right is worn obliquely.

(101) D.8010. Holotype. Orifices of two adult zooecia. (*a*) markedly flat, bilobed distal oral spines, that on the right is worn; (*b*) both distal oral spines are worn obliquely.

(102) D.8010. Holotype. Avicularia. (*a*) Unworn, (*b*) worn—the oval form is a cross-section of the avicularian stalk.

REMARKS. Lang (1922 : 305) interpreted *P. gregoryi* (Brydone) from Brydone's 1906 description and figure. Unfortunately these were not sufficiently clear to show the nature of the enlarged distal oral spines, and Lang supposed *P. gregoryi* to be similar to *P. damicornis* in these respects. He maintained the three species (*P. gregoryi, P. damicornis* and *P. palmata*) as separate solely on supposed differences in zoarial growth habit. However, it has been shown that the enlarged distal oral spines of the lectotype of *P. gregoryi*, S.M., B.36120, as well as other zooecial characters, are different from those of *P. palmata*. The latter is thus retained as a separate species, but is regarded as synonymous with *P. damicornis*. The supposed differences in zoarial growth habit of *P. palmata* and *P. damicornis* are very uncertain, and the two species are otherwise morphologically alike.

D.8009, D.28537 and D.21169, labelled by Lang " *Pelmatopora* [*Cribrilina*] *gregoryi* (Brydone) ", are also referred to *P. palmata* Lang. D.8009 and D.28537 are somewhat worn, but appear to have distally expanded, bilobed distal oral spines, and are thus placed in *P. palmata* with a query. D.21169 is better preserved and is placed in *P. palmata* with certainty. D.20204, holotype of *P. damicornis* Lang, is well preserved and may be definitely assigned to *P. palmata*.

It may be significant that there is relatively little material which can definitely be placed in *P. palmata*. This is possibly explained by difficulties imposed by poor preservation, particularly by wear of the enlarged distal oral spines. It has, however, been noted that worn bilobed oral spines differ in form from worn elongate oral spines such as occur in *P. gregoryi*. Where wear has reduced oral spines it is thus possible to differentiate between bilobed and elongate types, but the identification must also rest largely on other zooecial characters, such as zooecial size, range in number of costae, type and amount of interzooecial tissue and form and distribution of avicularia.

D.39355 and D.39356 have especially well-preserved enlarged distal oral spines typical of *P. palmata*.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta* to zone of *G.* [*A.*] *quadrata*.

SPECIMENS. D.8010. Holotype—see above. [The specimen is labelled " Idiotype of *P. gregoryi* (Brydone) ".]

D.28971. Paratype. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 8 of Gaster, between Upton Lane (Lambley's Lane), Sompting, and lane east of Charman Dean, north of Broadwater, Worthing, Sussex.

D.20204. Holotype of *P. damicornis* Lang. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata*. Seaford, Sussex. Collection unspecified.

?D.8009. Large fragment of a worn zoarium. Labelled by Lang " *Pelmatopora gregoryi* (Brydone). Idiotype ". Senonian, zone of *G.* [*A.*] *quadrata*. Sussex Coast. R. M. Brydone Collection.

?D.28537. Three fragments of the same zoarium. Labelled as for D.8009. Senonian, horizon unspecified, probably zone of *G.* [*A.*] *quadrata*. Southern England. F. H. Butler Collection.

D.21169. Zoarial fragment. Labelled as for D.8009. Senonian, zone of *G.* [*A.*]

quadrata. Southampton Waterworks Pit, Otterbourne, south of Winchester, Hampshire. L. Treacher Collection.

D.28865. Zoarial fragment. Senonian, zone of *O. pilula*, subzone of *E. scutata* var. *cincta.* Pit on East Hill, Rottingdean, east of Brighton, Sussex.

D.39339, D.39355–56. Three zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 15 of Gaster, large pit about one-third of a mile south-south-east of Salvington Windmill, and north of Hill Cottages, Salvington, Sussex.

D.39370, D.39378. Two zoarial fragments. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 20 of Gaster, lower pit on Patching Hill, north of St. John's Church, just above the 200 ft. contour, Patching, Sussex.

D.39410. Zoarial fragment. Senonian, zone of *G.* [*A.*] *quadrata.* Pit 35 of Gaster, small pit north of Rake Mead Plantation, Church Hill Shaw, about half a mile north-west of Rogers Farm, and west of South Park, just above the 400 ft. contour, Findon, Sussex.

PELMATOPORA sp.

(Pl. 15, fig. 7)

SPECIMEN. D.39330. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *B. mucronata.* Pit 105 of Rowe. White Pit Lane, Newport, Isle of Wight.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* elongate-elliptical with somewhat parallel sides. *Orifice* wider than high, distally and laterally rounded, with a straight proximal margin, its outline indented by two worn, enlarged distal oral spines. Only the distal pair of *oral spines* is visible, and only their distinctly rounded bases are preserved ; one enlarged oral spine is apparently hollow. Unmodified lateral oral spine-bases, if present, are concealed by secondary tissue. *Frontal shield* of variable length. *Costae* 20–24 (often 22), well defined, each with a fairly prominent *pelma* near its outer end, a second smaller pelma about half-way along its length, and a *pelmatidium,* often obscure, near the mid-line. *Lateral costal fusions* arise at the level of the pelmata producing two paired rows of *intercostal spaces* ; an additional irregularly developed row of small intercostal spaces may occur very close to the mid-line. The outermost intercostal spaces are slot-like. *Apertural bar* wide, flat, sloping down towards the proximal margin of the orifice. *Gymnocyst* entirely concealed by *interzooecial secondary tissue,* which fills the interzooecial valleys and which contains irregular, elongate, shallow *lacunae.*

Avicularia either oval or elongate and distally pointed. In the specimen, all avicularia are worn. There is, however, a strong tendency for the avicularia to be arranged in pairs, one near each proximal-lateral corner of the orifice. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 1·24 to 1·50 mm.
lz = 0·42 to 0·50 mm.
hr = 0·15 to 0·18 mm.
lr = 0·18 to 0·21 mm.
Avicularia Not measurable.

REMARKS. Lang has stated that *Pelmatopora* does not range above the zone of *G.* [*A.*] *quadrata*. However, D.39330 was collected from Pit 105 of Rowe, which was placed by him (1908 : 280) in the zone of *B. mucronata* " beyond doubt ". Recent collecting neither confirms nor disproves Rowe's suggested horizon for his Pit 105.

Some features of D.39330 are unlike any seen in the other known species of *Pelmatopora* ; the large size of the zooecia and the large number of costae are particularly unusual. One zooecium with 28 costae is distorted proximally and is regarded as abnormal, the usual range in the specimen being 20–24. The specimen obviously belongs to the group of *Pelmatopora* species which possess enlarged distal oral spines, and it is very probable that its enlarged oral spines are not of the expanded bilobed type but rather of the elongate, outwardly curving form, with rounded bases, such as are developed in *P. gregoryi*.

Although the specimen is unlike any known species of *Pelmatopora*, it is not advisable to erect a new species on a single somewhat worn specimen, and diagnosis must depend on better preserved material. The specimen is important as the only definite *Pelmatopora* from the zone of *B. mucronata*.

Although Voigt (1925*b* ; 1930) has assigned species from the zone of *B. mucronata* and higher horizons to *Pelmatopora*, these may not be correctly placed in that genus. An exceptional occurrence of *Pelmatopora* in the Danian is noted in the remarks on *P. daniensis* Voigt (p. 202).

SPECIES OF UNCERTAIN SYSTEMATIC POSITION PREVIOUSLY ASSIGNED TO *PELMATOPORA*

(1) ***Pelmatopora incerta*** Voigt

1925*b* *Pelmatopora incerta* Voigt, pp. 99, 100, pl. 3, figs. 9, 10 ; text-fig. 1.

1930 *Pelmatopora incerta* Voigt : Voigt, pp. 515, 561, pl. 31, fig. 15 [part of the same zoarium figured by Voigt, 1925*b*, pl. 3, fig. 9].

1939 *Pelmatopora incerta* Voigt : Voigt, p. 93, text-fig. 4*a* [copy of Voigt, 1925*b*, text-fig. 1].

DESCRIPTION. [Based on Voigt's descriptions and figures.] *Zoarium* unilaminar, encrusting. *Zooecia* about two-thirds as wide as long. *Orifice* about as long as wide, distally and laterally rounded, with a straight proximal margin, and with proximal-lateral " slits ". *Oral spine-bases* 4, *frontal shield* flatly arched. *Costae* about 20, closely spaced, " fused " only in the mid-line, each with a number of *pelmatidia* (the outermost and the innermost pelmatidia appear to be better developed than the intervening ones). Individual costae taper towards their inner ends and have slightly curved margins (? = the slightly crenulate outlines shown in Voigt's text-figure). *Lateral costal fusions* possibly present. The costae meet and fuse in the mid-line, producing a slight median ridge, or, as in at least one zooecium figured by Voigt, there is a flat median area. *Gymnocyst* well exposed, especially proximally to the frontal shield. *Interzooecial secondary tissue* apparently absent.

Avicularia apparently rare, elongate, oval (their form is not apparent from the photographs). *Ovicells* not seen.

REMARKS. Voigt has suggested that the species belongs to the group of " primitive ", less complex *Pelmatopora* species in which the costae are fused only in the mid-line. Although this is apparently so in several of the zooecia figured by Voigt, each costa also has a row of pelmata or pelmatidia—a condition found in the later, more complex species of *Pelmatopora*. Also, in the earlier forms of *Pelmatopora*, the costae are characterized by single, large pelmata developed near the mid-line of the frontal shield, or surrounding the median area of fusion when this is present. In *incerta* the larger pelmata or pelmatidia are distinctly at the extreme outer ends of the costae. The avicularia, described by Voigt as rare and elongate oval, are not discernible in the photographs. The stylized text-figure shows a single avicularium which is directed proximally and constricted laterally so that it is partly divided into a smaller semicircular portion and a distal area which is elongate-oval. This type of avicularium is unknown in other species of *Pelmatopora*.

In the absence of actual material it is undesirable to attempt to decide the definite systematic position of a species known only from inadequate photographs and descriptions. However, *incerta* is very probably not a *Pelmatopora*.

STRATIGRAPHICAL DISTRIBUTION. " Ob. Senongeschiebe ", Köthen.

(2) ***Pelmatopara grandiporosa*** Voigt

1930 *Pelmatopora grandiporosa* Voigt, pp. 514, 561, pl. 31, figs. 13, 14.

DESCRIPTION. [Based on Voigt's description and figures.] *Zoarium* unilaminar, encrusting. *Zooecia* about twice as long as wide. *Orifice* large, distally and laterally rounded, with a straight proximal margin. Number and form of *oral spines* unknown. *Frontal shield* apparently fairly flat, but curving down steeply at the margins to meet the gymnocyst. *Costae* about 10, very wide, very irregularly arranged, each with a large, oval *pelma* at the outer end. *Interzooecial secondary tissue* probably abundant.

Avicularia not distinct in the photographs, but, according to Voigt, many small avicularia occur between the zooecia.

REMARKS. From the photographs it is not possible to place this species in *Pelmatopora*. Voigt has suggested that the irregular arrangement of the few,very thick costae distinguishes the species from all known species of *Pelmatopora*, but this irregular arrangement of the costae, their coarseness, and the numerous interzooecial avicularia which Voigt mentioned, all contrast with the generally settled more complex pattern of *Pelmatopora* at the horizon involved. The placing of the species as a *Pelmatopora* is thus very uncertain, and, in the absence of actual material, it may not be described with certainty beyond " cribrimorph of the Pelmatoporidae ".

STRATIGRAPHICAL DISTRIBUTION. Granulatensenon. Gt. Bülten.

(3) ***Pelmatopora*** sp.

1930 *Pelmatopora* sp., Voigt, pp. 514, 561, pl. 31, fig. 12.

REMARKS. Voigt described the zoarium as " free ", bilaminar, with zooecia of Lz = 0·75 to 0·80 mm., and with frontal shield with 10–13 costae, he could not identify it with any of Lang's species of *Pelmatopora*.

The photograph of the specimen is small, but apparently shows at least one lateral costal fusion arising between the inner ends of the costae near a narrow median area.

From the description and photograph alone it seems very uncertain that the specimen should be placed in *Pelmatopora*.

STRATIGRAPHICAL DISTRIBUTION. "Granulatensenon". Gt. Bülten.

(4) *Pelmatopora daniensis* (Voigt)

1925*b* *Pelmatopora daniensis* Voigt, pp. 99, 100, pl. 3, figs. 4–6; text-fig. 2.
1930 *Pelmatopora daniensis* Voigt: Voigt, pp. 514, 561, pl. 31, fig. 11.

DESCRIPTION. (Based on Voigt's figures.) *Zoarium* apparently unilaminar, erect. *Zooecia* elongate-oval. *Orifice* wider than high, evenly rounded distally and laterally, proximal margin straight. *Oral spine-bases* 4, slightly enlarged, unmodified. *Frontal-shield* fairly well arched. *Costae* 20–24, each with 4 or 5 *lateral costal fusions* and *pelmatidia*; *outermost intercostal* spaces distinct. *Apertural bar* narrow, form obscure. *Gymnocyst* hardly visible; no *interzooecial secondary tissue*.

Avicularia proximally directed, large, elongate-oval, somewhat pointed, a prominent *transverse bar* divides a small semicircular proximal portion from a longer, bluntly triangular *rostrum*. Avicularia one-quarter to one-third as long as the zooecia they accompany, irregularly distributed, apparently comparatively rare. *Ovicells* not seen.

MEASUREMENTS. (According to Voigt, 1925*b*, p. 101.)
Zooecia Lz = 0·75 to 1·05 mm.
lz = 0·36 to 0·44 mm.
Avicularia Lz = 0·21 to 0·24 mm.
lz = 0·08 to 0·12 mm.

REMARKS. The shape of the orifice, the nature of the frontal shield, and particularly the form of the avicularia, distinguish this species from other species of *Pelmatopora*, but the characters are reminiscent in some ways of *Castanopora*.

It is difficult to be certain that the specimens which Voigt (1925: 99, 100, pl. 3, figs. 4–6) assigned to *Pelmatopora daniensis* should be placed definitely in that genus. However, Mr. Ole Berthelsen has very kindly shown me a single specimen from the Upper Danian of Herfølge, Seeland, which differs in some respects from Voigt's description and figures, but which is undoubtedly a *Pelmatopora*. The occurrence of this genus at this horizon is exceptional.

Voigt himself compared his species (1925*b*: 101) with *Repteschareila radiata* d'Orbigny (1852, pl. 716, figs. 4–6), commenting "Eine gewisse Ähnlichkeit mit unserer Form zeigt *Repteschareila radiata* d'Orb., ist jedoch von ihr durch das inkrustierende Zoarium und durch weniger zahlreiche (16–18) aus etwa je 7–8 Poren gebildete Punktreihen verschieden". *Repteschareila radiata* d'Orbigny was assigned by Lang (1922: 187) to *Rhiniopora*. As Voigt stated, it is an encrusting form with fewer costae and slightly fewer intercostal spaces than *daniensis*. It is also possible that *Repteschareila radiata* lacks avicularia.

The horizon of d'Orbigny's material is Campanian and Santonian of Meudon and

Saintes, France, whilst Voigt's specimen is of Danian age from Cöthen. On morphological and stratigraphical features the two are maintained as separate.

Stratigraphical distribution. Danian.

Genus ***CASTANOPORA*** Lang

1851 *Cellepora* (? *Dermatopora*) : von Hagenow, p. 99, pl. 10, fig. 19.
1851 *Escharipora* d'Orbigny, pls. 603, 684–687, 700, 703 [*partim*].
?1852 *Reptescharella* d'Orbigny, pls. 604, 715–716 [*partim*].
1852 *Reptescharipora* d'Orbigny, pls. 719, 720 [*partim*].
1852 *Escharipora* : d'Orbigny, p. 221 [*partim*].
?1853 *Reptescharella* : d'Orbigny, p. 464 [*partim*].
?1854 *Reptescharella* : d'Orbigny, p. 1097 [*partim*].
1854 *Escharipora* : d'Orbigny, p. 1097 [*partim*].
?1854 *Reptescharipora* : d'Orbingy, p. 1098 [*partim*].
1865 *Escharipora* : Ubaghs, p. 51.
1887 *Cribrilina* : Marsson, p. 96 [*partim*].
1894 *Membranipora* : Gregory, p. 62.
1896 *Membranipora* : Gregory, p. 212.
?1900*b* *Reptescharipora* : Canu, p. 457 [*partim*].
?1900*b* *Cribrilina* : Canu, p. 445 [*partim*].
1906 *Cribrilina* : Brydone, pp. 296–298 [*partim*].
1907 *Cribrilina* : Levinsen, pp. 155–156.
1913 *Cribrilina* : Brydone, pp. 436–438 [*partim*].
1916*a* *Rhiniopora* Lang, pp. 93, 96–97.
1916*a* *Castanopora* Lang, pp. 93–96.
1917 *Rhiniopora* : Lang, p. 172.
1919*c* *Castanopora* : Lang, p. 105, footnote.
1922 *Rhiniopora* : Lang, p. 180.
1922 *Castanopora* : Lang, p. 202.
1923 *Rhiniopora* : Waters, pp. 548, 551.
1923 *Castanopora* : Waters, pp. 548, 551, 558–560.
1925 *Cribrilina* : Levinsen, p. 375 [*partim*].
?1925 *Barroisina* : Levinsen, p. 384 [*partim*].
1925*c* *Cribrilina* : Voigt, p. 176 [*partim—C. asperula* only].
1927 *Cribrilina* : Voigt, p. 102 [*partim*].
?1927 *Barroisina* : Voigt, p. 102 [*partim*].
1927 *Rhiniopora* : Canu & Bassler, p. 37.
1927 *Castanopora* : Canu & Bassler, p. 25.
1929 *Castanopora* : Lang, pp. 438, 439.
1929 *Rhiniopora* : Lang, pp. 438, 439.
1930 *Rhiniopora* : Voigt, pp. 510, 511, 551, 561.
1930 *Castanopora* : Voigt, pp. 512, 513, 551, 561.
1935 *Castanopora* : Bassler, pp. 31, 65.
1935 *Rhiniopora* : Bassler, pp. 31, 190.
1953 *Castanopora* : Bassler, p. G192.
1953 *Rhiniopora* : Bassler, p. G193.
1953 *Rhiniopora* : Illies, p. 76.
1958 *Rhiniopora* : Butler & Cheetham, p. 1154.

Type species. (By original designation.) *Castanopora castanea* Lang, 1916*a* : 95. Senonian, " zone of *B. mucronata* ", probably boreal Maastrichtian, Rügen.

Emended diagnosis. Pelmatoporinae with numerous costae and numerous pelmata or pelmatidia and lateral costal fusions ; no secondary orifice ; no tertiary front wall, avicularia large, variously directed, of one, two or three kinds, or occasionally none.

REMARKS. Lang established the genera *Castanopora* and *Rhiniopora*, which I consider synonymous, to distinguish between the larger forms of Castanoporinae [here synonymous with the Pelmatoporinae] which lack secondary orifices and tertiary front walls. *Castanopora* was briefly diagnosed (Lang, 1916*a* : 93) as having " avicularia of two kinds, (1) directed distally and (2) directed proximally (the latter kind may be very rare or absent) ", while *Rhiniopora* was distinguished by its avicularia which " (if present) [are] of one kind only, proximally directed ".

In his subsequent expansion of these definitions (1922 : 181, 203) Lang diagnosed the two genera as follows : *Rhiniopora*—" Large Castanoporinae with numerous costae ; with no secondary [= tertiary] front-wall and no secondary aperture ; aviculoecia dimorphic, monomorphic, or none ; when dimorphic, some aviculoecia have long rostra, others pointed (but comparatively short) rostra, the former variously directed, the latter (? always) proximally directed ; when monomorphic, the aviculoecia are always of the latter type ". *Castanopora*—" Large Castanoporinae with numerous costae, no tertiary front-wall and no secondary aperture ; paired, distally directed aviculoecia, having apertures with elongate pointed rostra, are present, and some species have proximally-directed aviculoecia also, with short rostra ".

Lang thus distinguished between *Rhiniopora* and *Castanopora* solely on the form, orientation and kinds of avicularia present. He summarized his general conclusions (1922 : 156) and stated " In both genera the early species possess both long and short aviculoecia, but in *Rhiniopora* they are all variously and on the whole proximally directed, while in *Castanopora* two of the longer aviculoecia, situated more-or-less symmetrically one on each side of each aperture, are distally directed. *Rhiniopora* soon loses the longer aviculoecia and gradually tends to lose the shorter ones as well, while *Castanopora* retains the distally-directed ones and likewise gradually loses the aviculoecia with shorter apertures ".

Waters (1923 : 548) commented on the close similarity of some Cretaceous cribrimorphs with Recent European cribriline genera and stated "—in fact, it almost seems as if some of the Cretaceous forms might be considered varieties of our recent species, though rather larger ; and for the zoologist, the genera *Castanopora* and *Rhiniopora*, separated on slight grounds, are the most interesting (even if the names are dropped) as being represented in the living fauna ". Later (p. 551) Waters states "—the avicularia are given an important place and, like the other characters, must be carefully studied, but we may well remember that in recent Cheilostomatous forms there are not only genera, but species, which sometimes have avicularia, sometimes none. Sometimes, when Lang speaks of dimorphic avicularia, there seems no difference except size ... Judging from recent forms it would seem natural to say that avicularian characters have been used for the fossil genera where they have not more than specific value, even if that, and in recent species there is very great variation ".

Re-examination of all the available material assigned by Lang to *Rhiniopora* and *Castanopora* makes it difficult to confirm his separation of the genera. Variations in the form, position and orientation of the avicularia are insufficient for such a purpose. But these variations, in conjunction with other characters, are here used to distinguish species. Also, several of the original species which Lang placed in *Rhiniopora* appear to be synonymous, and the same applies to several species originally placed in *Castanopora*.

In the one species which has three kinds of avicularia there are a constantly paired, distally directed and pointed form and two unpaired proximally directed forms, the one sporadic and the other constantly placed just distal to each orifice. In species with two types of avicularia both types are not present in all zooecia; the most frequently occurring form is usually paired, distally directed and pointed, or occasionally spatulate, the less frequent form is unpaired, rounded, probably due to wear, or pointed, proximally directed and sporadic. Species with only one kind of avicularium also occur—avicularia of this type are frequently paired, pointed and either all distally or all proximally directed; or occasionally single, almost rounded and sporadic. Very occasionally avicularia may be absent.

I retain the name *Castanopora*, rather than *Rhiniopora*, for the revised genus, and thus the type-species is *Castanopora castanea* Lang.

STRATIGRAPHICAL DISTRIBUTION. Senonian—zone of *Marsupites* to Danian. Canu & Bassler (1933 : 53, 54) assigned two species from the Vincentown Limesand to *Rhiniopora*, thus extending the upward range of the genus into the Palaeocene.[1]

KEY TO THE SPECIES OF *CASTANOPORA*

- (I) Zoarium encrusting or erect, uni- or bilaminar.
  - (A) *Castanopora* with three types of avicularia . . . 1. ***C. armata*** sp. nov.
  - (B) *Castanopora* with two types of avicularia.
    - (1) 28–34 costae; avicularia distally directed, unpaired, very large, rounded, widely spatulate; a more frequent, proximally directed, small, rounded form also present . . . . . . 2. ***C. labiata*** (Levinsen)
    - (2) 16–23 costae.
      - (*a*) 4 (less often 3) lateral costal fusions on each costa; avicularia distally directed, frequently paired, very elongate, sharply pointed; a proximally directed, shorter, pointed form also present 3. ***C. retrorsa*** Lang
      - (*b*) 4-8 (often 6–7 lateral costal fusions on each costa).
        - (i) 4 oral spines only; orifice wider than high; avicularia distally directed; frequently paired, elongate, pointed; an occasional probably proximally directed, rounded sporadic form also present 4. ***C. castanea*** Lang
        - (ii) 4 (occasionally 5) oral spines; orifice slightly wider than high or as wide as high; avicularia distally directed, elongate, pointed, constantly paired near orifice; a less frequent, proximally directed, elongate pointed, sporadic form also present 5. ***C. dibleyi*** (Brydone)
        - (iii) 5 or 6 oral spines; avicularia possibly proximally directed, paired near orifice; an occasional, proximally directed, elongate, pointed, sporadic form also present . . . 6. ***C. guascoi*** (Ubaghs)

---

[1] The age of the Vincentown Limesand has been established as Palaeocene by (Loeblich & Tappan, 1957).

(C) *Castanopora* with one type of avicularium.
  (1) Zooecia small, Lz = 0·26 to 0·34 mm., lz = 0·17 to 0·27 mm., hr = 0·04 to 0·07 mm., lr = 0·06 to 0·09 mm. . 7. ***C. spooneri*** (Butler & Cheetham)
  (2) Zooecia larger, Lz more than 0·75 mm., lz more than 0·36 mm., hr more than 0·15 mm., lr more than 0·15 mm.
    (*a*) Orifice wider than high.
      (i) 5–6 oral spines ; 26–30 costae, each with about 9 lateral costal fusions ; avicularia distally directed, paired near orifice, elongate, pointed . . . . . . 8. ***C. multicostata*** (Voigt)
      (ii) 6 oral spines only ; costae with 8 lateral costal fusions ; 25–30 costae ; avicularia rounded, fairly frequent, sporadic 9. ***C. jurassica*** (Gregory)
    (*b*) Orifice higher than wide or height approximately the same as width.
      (i) Avicularia all distally directed, elongate, pointed ; 24–27 costae, each with 6–8 (often 7) lateral costal fusions ; 5 (sometimes 4) oral spines . . . . 10. ***C. glandulosa*** Lang
      (ii) Avicularia all proximally directed.
        (α) Avicularia with sharply pointed, elongate rostra 11. ***C. aviculosa*** (Lang)
        (β) Avicularia rounded, small or relatively large, occasional, sporadic, never paired ; 14–30 (often about 20) costae ; 4–8 (often 5–7) lateral costal fusions ; 4–5 oral spines 12. ***C. magnifica*** (d'Orbigny)

(D) *Castanopora* with avicularia ; 13–15 costae, each with 6–7 lateral costal fusions ; gymnocyst widely exposed . . . . . . 13. ***C. faujasi*** (Hagenow)

(II) Zoarium cylindro-quadrate, erect ; zooecia of the four adjacent rows alternate ; 20–25 costae . . . . 14. ***C. voigti*** sp. nov.

Cellepora perforata Quenstedt and *Reptescharella radiata* d'Orbigny were both referred to *Rhiniopora* by Lang (1922 : 183, 187). But as their systematic position is very uncertain, they have been omitted from the above key. A discussion of the status of these species is on p. 237.

Rhiniopora hispida Lang is also omitted from the above key ; the holotype, the only specimen, is very badly damaged and worn, making generic and specific diagnosis impossible. The species is discussed on p. 237.

Repteschoripora ornata d'Orbigny, referred to *Castanopora* by Lang (1922 : 213), is also omitted from the above key. The holotype is so poorly preserved that it is not possible to place the species with any certainty at all. The species is discussed on p. 238.

1. *Castanopora armata*[1] sp. nov.

(Pl. 16, fig. 1 ; Text-fig. 103)

HOLOTYPE. D.31717, an encrusting zoarial fragment of about ten zooecia. Maastrichtian, zone of *Belemnella lanceolata* (*O. lunata* Chalk). Southern Bluff, Trimingham, Norfolk. T. H. Withers Collection.

DIAGNOSIS. *Castanopora* with orifice as wide as high ; oral spine-bases 4 ; costae 18–22, each with 4–6 (often 5) lateral costal fusions and pelmatidia ; avicularia of three kinds : one form constantly paired on either side of the orifice, distally directed and elongate ; a second form very prominent, proximally directed, sporadic and variable in position ; a third form also proximally directed, but constantly placed just distally to the orifice.

---

[1] Lat., *armare*, " to arm ", referring to the numerous avicularia.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* slightly elongate-oval to broadly oval, very closely placed. *Orifice* about as wide as high, evenly rounded distally and laterally and without proximal-lateral constrictions, the proximal margin straight. *Oral spine-bases* 4, in two pairs placed distally and laterally respectively, all of about the same thickness. *Frontal shield* almost flat, its margins descending sharply at the inner ends of the somewhat slot-like *outer intercostal spaces*. *Costae* 18–22, each with 4–6 (usually 5) *lateral costal fusions* and an equivalent number of pelmatidia. *Apertural bar* only slightly raised, triangular, the apex directed proximally with a very slight median ridge. *Gymnocyst* almost concealed by the closeness of the zooecia and the development of the avicularia : no *interzooecial secondary tissue* present.



FIG. 103. *Castanopora armata* sp. nov. D.31717. Holotype. Adult zooecium with five avicularia and part of orifice of proximally adjacent zooecium ; *a*, avicularium of form *a* ; *b*, avicularium of form *b* ; *c* avicularium of form *c*.

Avicularia of three kinds : (*a*) a distally directed form, constantly paired one on each side of the orifice, each with a straight or slightly proximally curved *transverse bar*, separating a large, semicircular proximal portion from an elongate, spatulate *rostrum*, the distal end of which is narrower and almost parallel-sided—these avicularia may be directed distally or may tend to embrace the lateral-distal margins of the orifice and point obiquely inwards ; (*b*) proximally directed, sporadic, variable in position, and raised on a very stout stalk with a slightly curved *transverse bar* dividing a lunate or semicircular proximal portion from a pointed *rostrum* with inwardly curving margins ; (*c*) proximally directed, very similar to (*b*) but regularly placed just distally to each orifice. Several broken avicularia of this last type look superficially like broken ovicells. They may be recognized in the broken state by the fact that they do not obscure or interrupt the development of the distal pair of oral spines. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·72 to 0·75 mm.
lz = 0·45 to 0·56 mm.
hr = 0·14 to 0·16 mm.
lr = 0·16 to 0·17 mm.
Avicularia:
Type (*a*) Lz = 0·24 mm.
lz = 0·095 mm.
Type (*b*) Lz = 0·24 mm.
lz = 0·12 mm.
Type (*c*) Not measurable.

REMARKS. D.31717 differs from other species in possessing three types of avicularia. The most unusual type present is the regularly placed, proximally directed avicularium lying immediately distally to each orifice. The occurrence of frequent, proximally directed avicularia, taking into account both types, is unusual at this high stratigraphic level. Lang in fact suggested that the various species of *Castanopora* tend, in successively higher horizons, to have fewer and fewer proximally directed avicularia.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian, zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk).

SPECIMEN. Holotype—see above.

2. *Castanopora labiata* (Levinsen)

1907 *Cribrilina labiata* Levinsen, pp. 155, 156, 158, pl. opp. p. 160, figs. 1, 1*a*–*f*.
1916*a* *Rhiniopora labiata* (Levinsen) Lang, p. 96.
1922 *Rhiniopora labiata* (Levinsen) : Lang, p. 186.
1925 *Cribrilina valida* Levinsen, p. 376, pl. 5, figs. 58*a*–*d*—(copy of Levinsen's 1907 figures).
1927 *Cribrilina valida* Levinsen : Voigt, p. 102.
1930 *Cribrilina valida* Levinsen : Voigt, p. 551.

LECTOTYPE. (Chosen by Lang, 1916*a* : 96) Specimen figured by Levinsen, 1907, pl. opp. p. 160, fig. 1. Lower Maastrichtian, zone of *Belemnella lanceolata*. Møen and Aalborg, Denmark.

EMENDED DIAGNOSIS. *Castanopora* with 28–34 costae ; avicularia of two kinds : one form distally directed, very large, vicarious, rare, spatulate, with a straight transverse bar, the other form more frequent, proximally and obliquely directed, small, oval and slightly pointed proximally, with small lateral constrictions.

DESCRIPTION. (Based on Levinsen's description and figure.) *Zoarium* apparently erect, unilaminar. *Zooecia* large, broadly oval. *Orifice* evenly rounded distally and laterally, the sides tending to converge at the straight proximal margin, height of orifice about one-quarter the length of the zooecium. *Oral spines* unknown. *Frontal shield* flat. *Costae* 28–34, each with at least 4 (possibly 5 or 6) *lateral costal fusions*. *Pelmatidia* presumably present, number unknown. *Apertural bar* fairly wide, slightly curved proximally, with a slight median furrow. *Gymnocyst* probably concealed by *interzooecial secondary tissue* with triangular *lacunae*.

Avicularia rare, of two forms : (*a*) distally directed, large, vicarious, sporadic, broadly spatulate, divided by a *transverse bar* into a small, semicircular proximal portion and a very large, distally rounded *rostrum* ; and (*b*) more frequent, proximally directed, unpaired, small [about one half the size of form (*a*)], rounded or slightly pointed proximally, with slight lateral constrictions dividing a semicircular proximal portion from a larger oval-triangular *rostrum*. *Ovicells* not seen.

REMARKS. Levinsen's MSS. and figures were published posthumously in 1925 and edited by Nielsen & Mortensen, who stated (p. 285) that " the first seven plates represent the figures found together with the manuscript ".

On pl. 5, fig. 58*a–d*, appears a copy of Levinsen's 1907 figures of *Cribrilina labiata*. Here (1925), however, the species is called *Cribrilina valida* sp. nov. and it is described as such on p. 376 and 377. It is probable that Levinsen himself, having omitted a description of *Cribrilina labiata* in 1907, regarded the name as invalid, and fully described the form as *Cribrilina valida* intending to re-use the original 1907 figure. But the name *labiata* is valid, and *valida* is a synonym.

Levinsen's description (1925) supersedes that of Lang (1922 : 186) which was based only on Levinsen's 1907 figure. Levinsen described the zoarial habit, but gave no measurements. He stated that there were " no marks of spines "—this may imply that the margins of the orifices were worn obliterating the oral spines.

Lang (1922 : 186) referred the species to *Rhiniopora*. He regarded the marked dimorphism of the avicularia as a primitive feature suggesting affinities with his genus *Castanopora*, which here includes *Rhiniopora*.

The horizon of the species was given by Levinsen (1925) as Senonian, zone of *Belemnitella mucronata*.

STRATIGRAPHICAL DISTRIBUTION. Lower Maastrichtian, zone of *Belemnella lanceolata*.

SPECIMENS. None in the collection.

3. *Castanopora retrorsa* Lang

(Pl. 16, fig. 2 ; Text-fig. 104)

1909 *Cribrilina dibleyi* Brydone : White, pp. 34, 39 [*fide* Lang] (*non Cribrilina dibleyi* Brydone, 1906, p. 297, text-fig. 8).

1916*a* *Castanopora retrorsa* Lang, pp. 94, 95.

1922 *Castanopora retrorsa* Lang : Lang, p. 206, pl. 4, fig. 9 ; text-fig. 65.

HOLOTYPE. D.21170. Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *Marsupites*, subzone of *Uintacrinus*, *Uintacrinus* Band. North-west of Newland's Barn, Odiham, Hampshire. L. Treacher Collection.

EMENDED DIAGNOSIS. *Castanopora* with orifice wider than high ; oral spine-bases at least 4 ; costae 21–23, each with 4 (occasionally 3) lateral costal fusions and pelmatidia; avicularia of two kinds: a distally directed, frequently paired form, very long and sharply pointed, and a proximally directed, shorter, pointed form.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* relatively large, closely placed, elongate oval. *Orifice* wider than high, evenly rounded distally and laterally, the sides tending to diverge at the straight proximal margin. *Oral spine-bases* at least 4. *Frontal shield* evenly and well arched, possibly flattening slightly in the median area of fusion. *Costae* 21–23, fairly coarse, each with 4 (occasionally 3) *lateral costal fusions* and an equivalent number of *pelmatidia*. The *outer intercostal spaces* are long and slot-like. *Apertural bar* with a small, but distinct, median vertical projection formed by the slightly upturned distal ends of its component costae. *Gymnocyst* concealed by the closeness of the zooecia and the development of avicularia and ovicells.



FIG. 104. *Castanopora retrorsa* Lang, D.21170. Holotype. Parts of two adult zooecia (frontal shields not shown) five avicularia and two broken and worn ovicells (*ov.*). *a*, avicularia of form *a*; *b*, avicularia of form *b*.

Avicularia of two kinds: either (*a*) distally directed, usually paired one on each side of the orifice or sometimes unpaired near the orifice, divided by a *transverse bar*, which is distally concave, into a narrow, elongate, acute *rostrum* and a short, somewhat rounded, proximal portion; or (*b*) shorter, proximally directed, usually paired, one on each side of the orifice, or sometimes unpaired near the orifice, divided by a straight *transverse bar* into a somewhat triangular, pointed *rostrum* and a semicircular proximal portion: these avicularia may sometimes be directed proximally outwards. *Ovicells* hyperstomial, large, prominent, high, globular or slightly narrowed, smooth, resting on the frontal shield of the distally adjacent zooecium. The distal pair of oral spines may flank the opening of the ovicell.

MEASUREMENTS. *Zooecia* Lz = 0·84 to 1·06 mm.
lz = 0·50 to 0·59 mm.
hr = 0·15 to 0·18 mm.
lr = 0·18 to 0·21 mm.
Avicularia :
Type (*a*) Lz = 0·39 to 0·42 mm.
lz = 0·12 mm.
Type (*b*) Lz = 0·21 mm.
lz = 0·09 mm.

REMARKS. Lang (1922 : 208) regarded *C. retrorsa* as the most primitive species of *Castanopora.* Stratigraphically it is the earliest, well separated from the main group in the upper part of the zone of *B. mucronata* and in the *O. lunata* Chalk.

The very narrow, distally directed avicularia are particularly distinctive, as are the frequent proximally directed forms. Lang (1922, text-fig. 65) indicated four avicularia—a distally directed pair near the orifice, and a proximally directed pair near the proximal end of the zooecium, which are shown arising from the proximal-lateral areas of the gymnocyst. I have not seen in D.21170 any zooecium which has four definitely associated avicularia : there is either a distally directed pair near the orifice or these are replaced by a proximally directed pair, or one or other of the avicularian forms may occur singly near the orifice.

The widespread development of ovicells in D.21170 tends to obscure the oral spine-bases, which are, however, probably always 4 in number.

Lang regarded *C. retrorsa* as a form radical to the other species of *Castanopora.* Certainly the frequent apparent " replacement " of one form of avicularia by another seems to be a more primitive character. Some of the later species may possess proximally directed avicularia, but they always possess distally directed avicularia of fairly constant shape and position.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *Marsupites,* subzone of *Uintacrinus, Uintacrinus* Band.

SPECIMEN. Holotype—see above.

4. ***Castanopora castanea*** Lang

(Pl. 16, figs. 3–5 ; Text-fig. 105)

1916*a* *Castanopora castanea* Lang, p. 95.
1916*a* *Castanopora nucifera* Lang, p. 95.
1916*a* *Castanopora juglans* Lang, p. 95.
1922 *Castanopora nucifera* Lang : Lang, p. 214, pl. 4, fig. 11 ; text-fig. 67.
1922 *Castanopora juglans* Lang : Lang, p. 215, pl. 5, fig. 1 ; text-fig. 68.
1922 *Castanopora castanea* Lang : Lang, p. 217, pl. 5, fig. 2 ; text-fig. 69.
1923 *Castanopora castanea* Lang : Waters, pp. 548, 558, 560.
1923 *Castanopora juglans* Lang : Waters, pp. 558, 560.
1929 *Castanopora castanea* Lang : Lang, p. 438, text-fig. 39 (24).

1930 *Castanopora castanea* Lang : Voigt, pp. 513, 561, pl. 30, fig. 17.
1935 *Castanopora castanea* Lang : Bassler, p. 65.
1953 *Castanopora castanea* Lang : Bassler, p. G192, text-fig. 144, 21.



Fig. 105. *Castanopora castanea* Lang, D.16654. Holotype. Adult ovicelled zooecia, showing five distally directed, generally worn avicularia (*a'*) and one sporadic, large, rounded avicularium (*a*). *fs*, broken edge of frontal shield of adjacent zooecium ; *o*, worn orifices of adjacent zooecia ; *ov*, broken and worn ovicells ; *x*, break in the surface—not an avicularium.

Holotype. D.16654. Zoarial fragment encrusting a piece of echinoid test. Horizon given as Senonian, zone of *B. mucronata*—probably boreal Maastrichtian. Rügen. Agnes Laur Collection.

Emended diagnosis. *Castanopora* with orifice wider than high ; 4 oral spine-bases ; 16–23 costae, each with 4–8 (often 6–7) lateral costal fusions and pelmatidia ; avicularia of two kinds—a common, frequently paired, distally directed and pointed form with elongate, triangular rostrum, and an occasional, sporadic form which is large, rounded and probably proximally directed.

Description. *Zoarium* encrusting, unilaminar. *Adult zooecia* closely placed, relatively large, broadly-oval to somewhat elongate-oval. *Orifice* generally somewhat wider than high, not very large, evenly rounded distally and laterally and without proximal-lateral constrictions, the proximal margin straight, usually shorter than the greatest width of the orifice. *Oral spine-bases* 4, of medium size, the lateral pair only slightly thicker than the distal pair. *Frontal shield* very flatly arched or slightly but evenly arched, its margin descending sharply at the inner ends of the *outermost intercostal* spaces which are more slot-like than rounded. *Costae* 16–23, each with 4–8 (often 6–7) *lateral costal fusions* and an equivalent number of *pelmatidia*. *Apertural bar* of variable shape; either of medium thickness, straight and not very prominent, or fairly wide, triangular, thickened, and slightly expanded proximally in the mid-line: a very slight median ridge may be present, or the median part of the apertural bar may distinctly protrude. None of the apertural bars appears to be furrowed in the mid-line. *Gymnocyst* almost totally concealed due to the closeness of the zooecia and the development of the avicularia: no *interzooecial secondary tissue* present.

Avicularia of two kinds: (*a*) distally directed, elongate, with a long, triangular, pointed *rostrum*, divided by a straight or slightly proximally curved *transverse bar* from the small, semicircular proximal portion. This type is most frequently developed, commonly paired, one on either side of each orifice and placed near the distal-lateral margins, or more occasionally occurring singly on one side or the other of the orifice. (*b*) A second type occurs very occasionally: it is proximally directed, large, rounded, sporadic, and found more commonly in the earlier stages of zoarial growth.

Ovicells hyperstomial, smooth, globular, of medium size but prominent. The opening of the ovicell may be flanked by the distal pair of oral spines.

Young zooecia possess 4 oral spine-bases, the frontal shield is more evenly and markedly arched than that of the adult zooecia but still possesses a relatively flat median area of fusion. Costae 14 or 15, each with ?4 lateral costal fusions and ?an equivalent number of pelmatidia. Apertural bar much less prominent than in the adult zooecia. Definitely paired, distally directed avicularia are scarce. One large, rounded, proximally directed avicularium of the second type described above occurs between two zooecia of the second ring round the ancestrula in specimen N.C.M.76–937 (27/1).

Measurements. *Adult zooecia* Lz = 0·78 to 1·00 mm.
lz = 0·50 to 0·75 mm.
hr = 0·12 to 0·15 mm.
lr = 0·15 to 0·18 mm.
Avicularia:
Type (*a*) Lz = 0·27 to 0·39 mm.
lz = 0·12 mm.
Type (*b*) Lz = 0·21 mm.
lz = 0·18 mm.
Young zooecium Lz = 0·57 mm.
lz = 0·39 mm.
hr = 0·15 mm.
hr = 0·15 mm.

REMARKS. *Castanopora juglans* Lang and *C. nucifera* Lang are here regarded as synonymous with *C. castanea* Lang.

All the available specimens previously assigned to *C. castanea*, *C. juglans* and *C. nucifera* have been re-examined. Three specimens had been assigned to *C. castanea*, and *C. nucifera* was represented only by the holotype (D.15600). Though the latter has a more delicate structure than the specimens of *C. castanea* and has a slightly arched frontal shield, these differences are probably due to somewhat different growth conditions. But in both species the size and shape of the zooecia are closely similar; there are 4 oral spine-bases; the orifice tends to be wider than high; there are 16–23 costae, and often 6 or 7 lateral costal fusions on each costa (in D.15600 there may be 8 lateral costal fusions present between some costae); the distally directed avicularia are similar in position and type; and both species also have very occasional, rounded, sporadic, proximally directed avicularia. The horizon of both species is Senonian, zone of *B. mucronata*.

C. juglans is represented in the collection by the holotype (D.15608) and one other specimen (D.30110). According to Lang (1922 : 216) "*C. juglans* differs from *C. nucifera* ... only in the shape of the aperture [orifice] and in the width of the apertural bar. It is of interest as being, in both respects, a connecting link between *C. nucifera* and *C. castanea*".

The shape of the orifice of the specimens assigned to *C. juglans* differs from that of *C. castanea* only very slightly and the same relationship between height and width is maintained. Beyond using the values of hr and lr, which appear to be relatively constant for a given species, it is very difficult to apply a fine division of shapes of orifices as a diagnostic feature. In *C. castanea*, *C. juglans* and *C. nucifera*, hr is almost always less than lr but the shape of the orifice may vary slightly even within a single zoarium. I do not regard slight widening or thickening of the apertural bar as of diagnostic value. A marked median ridge or process is a definite feature which can be used, in conjunction with other zooecial characters, for specific diagnosis, but slight variations in the width or thickness of the apertural bar occur frequently in different zooecia of the same zoarium. In the material examined and here assigned to *C. castanea*, such variations within single zoarial fragments were frequently found. Lang himself regarded *C. juglans* as a linkage form between *C. castanea* and *C. nucifera*: the three belong to a single species. The holotype of *C. castanea* is well preserved: in one part two zooecia coalesce and share a single orifice.

Voigt (1930 : 513, pl. 30, fig. 17) briefly noted and figured a specimen which he assigned to *C. castanea* Lang. The horizon is again Upper *B. mucronata* Zone of Rügen. The measurements given and the range in number of costae agree with the material which has here been revised.

Specimens N.C.M.76.937(27/1–4) are parts of a previously undescribed zoarium also placed in *C. castanea*; much of the early growth stages of the zoarium is fairly well preserved and the description and measurements of the young zooecia of the species are based on parts of this zoarium.

STRATIGRAPHICAL DISTRIBUTION. Senonian, upper part of zone of *B. mucronata*, and Maastrichtian, zone of *Belemnella lanceloata* (*O. lunata* Chalk), and boreal Maastrichtian of Rügen.

SPECIMENS. D.16654. Holotype—see above.

D.14990. Zoarial fragment encrusting a piece of echinoid test. Specimen badly damaged and worn, labelled " *C. castanea* Lang ", but the identification is here questioned. Horizon given as Senonian, zone of *B. mucronata*, Rügen, but probably boreal Maastrichtian. Agnes Laur Collection.

D.15347. Zoarial fragment encrusting a piece of echinoid test. Labelled " Senonian, zone of *B. mucronata*, Rügen ", probably Maastrichtian. Agnes Laur Collecton.

D.15600. Zoarial fragment encrusting a piece of echinoid test. Holotype of *C. nucifera* Lang. Maastrichtian, zone of *Belemnella lanceolata* (*O. lunata* Chalk). Trimingham, Norfolk. A. C. Savin Collection.

D.15608. Zoarial fragment encrusting a piece of echinoid test. Holotype of *C. juglans* Lang. Horizon, locality and collection as for D.15600.

D.30110. Zoarial fragment labelled " *C. juglans* Lang " encrusting a fragment of an *Ostrea lunata*. Horizon, locality and collection as for D.15600.

OTHER MATERIAL. N.C.M.76.937(27/1–4). Parts of an extensive zoarium with young zooecia, encrusting four fragments of the same echinoid test. Senonian, zone of *B. mucronata*. Weybourne Mill, Weybourne, Norfolk. R. M. Brydone Collection.

5. ***Castanopora dibleyi*** (Brydone)

(Pl. 16, figs. 6–8 ; Pl. 17, figs. 1, 2 ; Text-figs. 106, 107)

1906 *Cribrilina dibleyi* Brydone, p. 297, text-fig. 8.
1909 *non Cribrilina dibleyi* Brydone : White, pp. 34, 39.
1913 *Cribrilina dibleyi* Brydone : Brydone, p. 437, pl. 14, fig. 9.
1916*a* *Castanopora dibleyi* (Brydone) Lang, pp. 94, 95.
1921 *Castanopora dibleyi* (Brydone) : Lang, p. xci.
1922 *Castanopora dibleyi* (Brydone) : Lang, p. 209, pl. 14, fig. 10 ; text-fig. 66*a*, *b*.
1923 *Castanopora dibleyi* (Brydone) : Waters, pp. 548, 559, 560.
1929 *Castanopora dibleyi* (Brydone) : Lang, p. 438, text-figs. 39 (22, 23).
1930 *Castanopora dibleyi* (Brydone) : Voigt, pp. 512, 543, 561, pl. 30, fig. 18.
?1930 *Castanopora* sp. Voigt, pp. 512, 561, pl. 30, fig. 20.

LECTOTYPE. (Chosen by Lang, 1922 : 210.) S.M., B.36115. A very large fragment of zoarium, with some young zooecia, encrusting a piece of echinoid test. Two zooecia of this specimen were figured by Brydone (1906 : 297, text-fig. 8) as *Cribrilina dibleyi*, and chosen by Lang as lectotype of *Castanopora dibleyi*. Maastrichtian, zone of *Belemnella lanceolata* (*O. lunata* Chalk). Trimingham, Norfolk. R. M. Brydone Collection.

EMENDED DIAGNOSIS. *Castanopora* with orifice as wide as, or slightly wider than, high ; oral spine-bases usually 4 (occasionally 5) ; costae 16–20, each with 5, 6 or 7 lateral costal fusions and pelmatidia ; avicularia of two kinds : a constantly paired, distally directed, elongate, pointed form near the orifice, and a proximally directed, less frequent, sporadic, pointed form.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Adult zooecia* relatively small, broadly oval, some almost circular in outline ; very closely placed. *Orifice* slightly wider than high or of equal height and width, straight sided but rounded distally, straight posteriorly; there are no proximal-lateral constrictions. *Oral spine-bases* commonly 4, the lateral pair usually thicker than the distal pair, a fifth small, slender, median, distal oral spine may be present. *Frontal shield* broad, often almost circular in outline, very flatly arched, not sharply descending at the margin. *Costae* 16–20, each with 5, 6 or 7 *lateral costal fusions* and an equivalent number of *pelmatidia*, each of which may be surrounded by a small rim producing a minute boss-like process opposite each lateral costal fusion: this is well seen in the lectotype. *Apertural bar* not prominent, somewhat swollen at either end; a very slight median ridge may be present. *Gymnocyst* only very occasionally exposed where the closeness of the zooecia and the development of avicularia have not obscured it.

Avicularia of two kinds: (*a*) a constantly paired form placed one on either side of and close to the orifice, distally directed, elongate and pointed, divided by a slightly curved *transverse bar* into a semicircular proximal portion and a very elongate, narrow, triangular or spatulate *rostrum*; (*b*) a proximally directed, infrequent, sporadic form, similar to type (*a*) but probably slightly shorter.

Ovicells hyperstomial, prominent, smooth, globular, slightly laterally compressed near the front, often with a broad median ridge. The lateral pair of oral spines may flank the entrance to the ovicell.

Young zooecia smaller and relatively less broad than the adult zooecia. Orifices as wide as, or slightly wider than, high. Oral spine-bases 4 or 5, the latter number apparently occurring more frequently in the younger zooecia. Frontal shield more roundly arched than in the adult. Costae 11–15, each with 4 or 5 lateral costal fusions. The apertural bar of the ancestrula of D.8003 has a slight median, proximal bend, but the zooecia in the first ring have straight apertural bars. Gymnocyst not visible in the circum-ancestrular area. Avicularia not paired on the ancestrula. The typically paired form which is distally directed on either side of the orifice occurs in the first ring of zooecia. The rostra of these avicularia tend to be directed outwards (in the adult the paired avicularia near the orifice are more parallel to its mid-line). Between two zooecia of the first ring round the ancestrula of D.8003 is a proximally directed, slightly pointed, avicularium, which is larger than the avicularia of type (*a*) in the same ring of zooecia. Proximally directed avicularia possibly occur more frequently in the circum-ancestrular area than in the adult parts of the zoarium, but, as in S.M., B.36267, they may be quite abundant locally in ovicelled parts of the zoarium.

Measurements. *Adult zooecia* Lz = 0·66 to 1·00 mm.
lz = 0·54 to 0·67 mm.
hr = 0·12 to 0·19 mm.
lr = 0·12 to 0·19 mm.
Young zooecia
(of S.M., B.36267) Lz = 0·50 to 0·55 mm.
lz = 0·32 to 0·40 mm.
hr = 0·10 to 0·11 mm.
lr = 0·12 to 0·13 mm.

Ancestrula
(of D.8003) Lz = 0·42 mm.
lz = 0·24 mm.
hr = 0·095 mm.
lr = 0·090 mm.

Avicularia
(in adult part of zoarium) :

$$\text{Type } (a)\begin{cases}\text{Lz} = 0{\cdot}21 \text{ to } 0{\cdot}43 \text{ mm.}\\ \text{lz} = 0{\cdot}065 \text{ to } 0{\cdot}14 \text{ mm.}\end{cases}$$

$$\text{Type } (b)\begin{cases}\text{Lz} = 0{\cdot}27 \text{ to } 0{\cdot}30 \text{ mm.}\\ \text{lz} = 0{\cdot}10 \text{ to } 0{\cdot}19 \text{ mm.}\end{cases}$$

(Associated with the ancestrula of D.8003)

$$\text{Type } (a)\begin{cases}\text{Lz} = 0{\cdot}085 \text{ mm.}\\ \text{lz} = 0{\cdot}050 \text{ mm.}\end{cases}$$

(Associated with the first ring of zooecia of D.8003)

$$\text{Type } (a)\begin{cases}\text{Lz} = 0{\cdot}125 \text{ mm.}\\ \text{lz} = 0{\cdot}060 \text{ mm.}\end{cases}$$

$$\text{Type } (b)\begin{cases}\text{Lz} = 0{\cdot}180 \text{ mm.}\\ \text{lz} = 0{\cdot}155 \text{ mm.}\end{cases}$$

REMARKS. All the available specimens of *C. dibleyi* (Brydone) in the British Museum (Natural History) have been re-examined, as well as those figured by Brydone (1906 : 297, text-fig. 8—the lectotype S.M., B.36115 ; 1913 : 437, pl. 14, fig. 9—S.M., B.36267). D.8003 and S.M., B.36267 are of particular interest : the former is labelled as a metatype and has a well-preserved ancestrula and circum-ancestrular area ; the latter also possesses young zooecia close to the ancestrula. The description of the young zooecia is based largely on these specimens. In D.8004 well-preserved proximally directed avicularia are present in addition to some very young zooecia. D.15599 also has a preserved ancestrula and numerous broken ovicells. D.20609 is a very worn zoarial fragment encrusting a piece of *Inoceramus* shell—fairly young zooecia are present, but the preservation is poor and the identification as *Castanopora dibleyi* is here queried. All the specimens are from the Maastrichtian, zone of *Belemnella lanceolata* (*O. lunata* Chalk) of Trimingham, Norfolk, except for D.20609 which is from the Senonian, zone of *B. mucronata*, also from Norfolk. Apart from *C. retrorsa* apparently no other species of *Castanopora* occur below the zone of *B. mucronata*.

Voigt (1930 : 512, pl. 30, fig. 18) briefly described a specimen as *C. dibleyi* (Brydone). The measurements and the number of costae agree with those given above. The reference to " comparatively short and blunt " avicularia apparently refers to worn examples. The figure shows no elongate pointed avicularia, those shown all appearing to be somewhat worn. It is probable that the specimen referred to *Castanopora* sp. by Voigt (1930 : 512, 561, pl. 30, fig. 20) is also *C. dibleyi* (Brydone).

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian, zone of *Belemnella lanceolata* (*O. lunata* Chalk) and Senonian, zone of *B. mucronata* in England, and boreal Maastrichtian of Rügen.



106



107

FIGS. 106–107. *Castanopora dibleyi* (Brydone).

(106) D.8003—metatype and D.8004. Orifices and avicularia of three adult zooecia (frontal shields not shown). Avicularia of two kinds: *a*, paired, distally pointed and directed; *b*, occasional, sporadic, interzooecial.

(107) S.M., B. 36267. Figured Brydone (1913, pl. 14, fig. 9) as *Cribrilina dibleyi*. Orifice of adult ovicelled zooecium with four avicularia. Proximally directed avicularia of type *b* occur more frequently than is usual in other specimens of *Castanopora dibleyi* examined here. Possibly type *b* avicularia develop particularly in association with ovicelled zooecia, especially where development of the ovicell has apparently disturbed the normal development of the distally directed avicularia of type *a*.

SPECIMENS. D.8003. Zoarial fragment with ancestrula, encrusting a piece of echinoid test. Labelled " metatype specimen ". Horizon, locality and collection as for the lectotype.

D.8004. Zoarial fragment encrusting a piece of echinoid test. Horizon, locality and collection as for the lectotype.

D.15599. Zoarial fragment encrusting a piece of echinoid test. Horizon and locality as for the lectotype. A. C. Savin Collection.

D.20609. Part of a worn zoarium encrusting a fragment of *Inoceramus* shell. Labelled by Lang " *C. dibleyi* (Brydone) ", but the determination is doubtful. Senonian, zone of *B. mucronata*. Norwich, Norfolk. T. G. Bayfield Collection.

OTHER MATERIAL. S.M., B.36115. Lectotype—see above.

S.M., B.36267. Zoarial fragment with ovicelled and very young zooecia encrusting a piece of echinoid test. [Part of this specimen was figured by Brydone (1913 : 437, pl. 14, fig. 9) as *Cribrilina dibleyi*]. Horizon, locality and collection as for the lectotype.

6. ***Castanopora guascoi*** (Ubaghs)

1865	*Escharipora guascoi* Ubaghs, p. 51, pl. 2, fig. 3*a–c*.
1913	*Cribrilina guascoi* (Ubaghs) Brydone, p. 437.
1916*a*	*Castanopora guascoi* (Ubaghs) Lang, pp. 95, 96.
1922	*Castanopora guascoi* (Ubaghs) : Lang, p. 221.
1930	*Castanopora guascoi* (Ubaghs) : Voigt, pp. 512, 561, pl. 31, fig. 1.

LECTOTYPE. (Chosen by Lang, 1922 : 221.) The specimen figured by Ubaghs, 1865, pl. 2, fig. 3*b*. Maastrichtian, Valkenburg (Fauquemont), east of Maastricht, Limbourg, Holland.

EMENDED DIAGNOSIS. *Castanopora* with orifice wider than high ; oral spine-bases probably 5 or 6 ; costae 18–20 ; gymnocyst well exposed ; avicularia of two kinds, a distal-lateral pair, apparently proximally directed, near each orifice, and an occasional, unpaired, sporadic, proximally directed form.

DESCRIPTION. (Based on Ubaghs' and Voigt's figures.) *Zoarium* encrusting or erect, uni- or bilaminar. *Zooecia* relatively broad, elliptical. *Orifice* wider than high, evenly rounded distally and laterally, the sides tending to converge at the straight proximal border. *Oral spine-bases* fairly large, of uncertain number, definitely more than 4, probably 5 or 6. *Frontal shield* flat, descending sharply at the outer ends of the costae. *Costae* 18–20. Number of *lateral costal fusions* and *pelmatidia* unknown, but probably few. *Apertural bar* fairly prominent, swollen at each end, somewhat rounded, not very thick, straight, with no marked median-proximal widening. *Gymnocyst* well exposed, much more so than in other species of *Castanopora*, visible laterally and proximally to each frontal shield ; no *interzooecial secondary tissue*.

Avicularia of two kinds : (*a*) a distal-lateral pair near each orifice, each with a straight *transverse bar*, orientation uncertain, possibly proximally directed ; (*b*) an occasional, sporadic form, proximally directed, elongate and pointed, divided by a straight *transverse bar* into a small semicircular proximal portion and an elongate, triangular *rostrum*. *Ovicells* not seen.

MEASUREMENTS. (According to Voigt, 1930 : 512.)
Zooecia Lz = About 1·0 mm.
lz = 0·60 mm.

REMARKS. The emended diagnosis and description of *C. guascoi* are based on a consideration of Ubaghs' original description and figure and Voigt's description and figure of a specimen from Maastricht.

Voigt (1930 : 512) mentioned the zoarial habit, size of zooecia, and number of costae, and his figure (pl. 31, fig. 1) also shows sporadic avicularia and more than four oral spine-bases. The nature of the paired avicularia is not determinable from the photograph.

Ubaghs (1865 : 51) in his original description referred to a " distal bulge hiding the ovicell ", suggesting that the ovicell, as with other species of *Castanopora*, is hyperstomial. He also mentioned that " above the orifice, almost at the foot of the following cell [zooecium] there is found, on either side of the aforementioned orifice, a depression in which there lies an angular, non-marginate, lateral pore " [translation]. This could refer to worn, paired avicularia, and similar worn avicularia appear to be present in the specimen figured by Voigt.

Ubaghs compared his original specimen with *Escharipora pretiosa* d'Orbigny [=*E. magnifica* d'Orbigny] which Lang (1922 : 212) also referred to *Castanopora*. Ubaghs stated that *E. guascoi* " has several similarities with *E. pretiosa* d'Orbigny, but may be distinguished from it by the lateral pores, which in d'Orbigny's species are elongate, slit-like and marginate. Also the cells [zooecia] of d'Orbigny's species are more elongate and show more rows of pores [intercostal spaces] piercing the cell surface [frontal shield] . . . Apart from this, our species (*E. guascoi*) shows, between the rows of cells, " blister-like ", elongate slits, which are rounded above [distally] and " turned down " below [proximally]. I am of the opinion that these should be regarded as avicularia . . . " [translation]. In his figure of *E. guascoi* Ubaghs showed this second type of proximally directed sporadic avicularium ; a similar avicularium is shown in the photograph of Voigt's specimen from Maastricht.

Until more specimens are available the description of *Castanopora guascoi* must rest on the interpretation of Ubaghs' and Voigt's figures. One of the most important features—the nature of the paired avicularia near the orifice—is still not determinable ; apart from this most of the other diagnostic characters are reasonably clear and the described specimens are retained as a separate species of *Castanopora*.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian.

SPECIMENS. None in the collection.

7. **Castanopora spooneri** (Butler & Cheetham)

1958 *Rhiniopora spooneri* Butler & Cheetham, p. 1154, text-figs. 1, 2.

HOLOTYPE. Louisiana State University Geology Museum 1390. " . . . from a chalk bed exposed at Rayburn's dome, Bienville Parish, Louisiana . . . equivalent to the Upper Cretaceous Saratoga Chalk of Arkansas . . . ".

PARATYPE. L.S.U. Geology Museum 1391. Horizon and locality as for the holotype.

EMENDED DIAGNOSIS [Based on Butler & Cheetham's diagnosis and detailed description and figures (1958 : 1154–1156).] *Castanopora* with small zooecia ; orifice of equal height and width or wider than high ; oral spine-bases 5 ; frontal shield with 14–20, usually 16–18 costae, each with 4–6 inconspicuous pelmatidia with 4 or 5 lateral costal fusions ; apertural bar curved, with a small median process ; gymnocyst concealed by the closely spaced zooecia ; avicularia interzooecial, of one kind only, distally pointed and directed without transverse bars or condyles ; ovicells hyperstomial, prominent, large, globular or somewhat elongate with a median, longitudinal ridge.

MEASUREMENTS. (According to Butler & Cheetham, 1958 : 1156.)
Zooecia Lz = 0·26 to 0·34 mm.
lz = 0·17 to 0·27 mm.
hr = 0·04 to 0·07 mm.
lr = 0·06 to 0·09 mm.
Avicularia Lz = 0·11 to 0·15 mm.
lz = 0·11 to 0·15 mm.

REMARKS. The holotype and paratype specimens of *Rhiniopora* [=*Castanopora*] *spooneri* encrust the inner surfaces of valves of *Ostrea vesicularis* and *Inoceramus*.

The most distinctive features of the species are the small dimensions of the adult zooecia and the character and size of the avicularia. These features, combined with the range in number of the costae, pelmatidia and lateral costal fusions, clearly distinguish the species.

STRATIGRAPHICAL DISTRIBUTION. Upper Cretaceous, equivalent of the Saratoga Chalk of Arkansas, U.S.A. The Saratoga Chalk was placed in the lower part of the Maastrichtian by Stephenson *et. al.* (1942, Chart 9, opp. p. 448).

SPECIMENS. None in the collection.

8. ***Castanopora multicostata*** Voigt

1930 *Castanopora multicostata* Voigt, pp. 512, 561, pl. 30, fig. 19.

LECTOTYPE. Specimen figured by Voigt, 1930, pl. 30, fig. 19. Horizon given as " Senonian, upper part of zone of *B. mucronata*. Rügen "—probably boreal Maastrichtian.

DIAGNOSIS. *Castanopora* with orifice generally wider than high ; oral spines 5–6 ; 26–30 costae, each with about 9 lateral costal fusions ; avicularia of possibly only one type, distally directed, paired near the orifice, pointed, elongate.

DESCRIPTION. Voigt described the species as follows : " *Zoarium* encrusting. *Zooecia* about 0·9 mm. long and 0·55 to 0·60 mm. wide, elliptical. *Gymnocyst* very slightly developed. *Frontal shield* formed of about 30 very closely placed *costae*, between which there are, on average, 9 *lateral costal fusions*. *Orifice* 0·15 mm. high, 0·15 to 0·20 mm. wide, with 5–6 *oral spines* (=' normal ' orifice of Lang).

" *Avicularia* mainly paired, beside orifice, between 0·25 and 0·45 mm. long, distally directed, elongate, pointed " [translation].

To these characters it is possible to add the following from Voigt's figure : *Zoarium* unilaminar. *Zooecia* not so closely placed as in other species of *Castanopora*. *Orifice* generally wider than high, relatively small, evenly rounded distally and laterally, proximal margin straight. The lateral pair of *oral spine-bases* tend to be larger than the others. *Frontal shield* flatly arched ; *costae* 26–30. *Apertural bar* fairly thick and prominent, slightly swollen at each end, bluntly triangular, with the apex proximally directed. *Gymnocyst* exposed laterally and proximally to each frontal shield.

Avicularia apparently of one type only, divided by a straight *transverse bar* into a small semicircular proximal portion and a narrowly elongate, acute, triangular *rostrum* ; placed near the distal-lateral margins of the orifice. *Ovicells* not described or figured.

Measurements (*vide* Voigt).

Zooecia Lz = About 0·90 mm.
lz = 0·55 to 0·60 mm.
hr = 0·15 mm.
lr = 0·15 to 0·20 mm.
Avicularia Lz = 0·25 to 0·45 mm.
lz not given.

Remarks. In his observations on *C. multicostata* Voigt stated (1930 : 513), " The figure (pl. 30, fig. 19) is interesting in that two ' lobes ' of the same colony coalesce and that in the critical zone (where the two growing branches of the zoarium meet) the zooecia appear to be deformed in several ways. In parts, several zooecia have coalesced and others have been constricted and have not developed an orifice, so that they appear as ' rosettes '. This situation is, of course, only possible because of the intercommunication of the individuals. From several *Castanopora* species from the Rügen chalk, described by Lang, this species is distinguished by its large number of costae, and, at the same time, the small size of the zooecia. The nearest species appears to be *Castanopora magnifica* (d'Orb.) which is 1·14 to 1·26 mm. long and which possesses 12 oral spines—as far as can be seen from d'Orbigny's drawing. Apart from this, the French species is bilamellar " [translation].

The large number of costae and lateral costal fusions and the occurrence of 5 or 6 oral spines are distinctive features of *C. multicostata* Voigt. The generally small size of the zooecia, and of the orifices, may also be diagnostic. Only a relatively small area of zoarium is shown in Voigt's figure, and no sporadic, more rounded and proximally directed avicularia, such as usually occur in other species of *Castanopora* from Rügen, can be seen.

Stratigraphical distribution. Given as " Senonian, upper part of zone *B. mucronata* ", but probably boreal Maastrichtian.

Specimens. None in the collection.

9. *Castanopora jurassica* (Gregory)

(Pl. 17, figs. 3–5 ; Text-figs. 108, 109)

1894 *Membranipora jurassica* Gregory, p. 62, text-fig. 1.
1896 *Membranipora jurassica* Gregory : Gregory, p. 212, text-fig. 21.
1897 *Membranipora jurassica* Gregory : Woodward, pp. 63, 64, text-fig. 107.
1907 *Membranipora jurassica* Gregory : Bather, pp. 120, 122, text-fig. 65.
1916*a* *Rhiniopora jurassica* (Gregory) Lang, p. 96.
1916*a* *Rhiniopora scabra* Lang, pp. 96, 97.
1922 *Rhiniopora scabra* Lang : Lang, p. 196, pl. 4, fig. 7 ; text-fig. 62.
1922 *Rhiniopora jurassica* (Gregory) : Lang, p. 197.
1930 *Rhiniopora scabra* Lang : Voigt, p. 551.

HOLOTYPE. D.180, large fragment of worn and damaged zoarium partly embedded in matrix. Very probably Maastrichtian. Maastricht. Old Collection.

EMENDED DIAGNOSIS. *Castanopora* with orifice wider than high ; zooecia up to 1·45 mm. long ; oral spine-bases 6 ; costae 25–30, each with 8 lateral costal fusions and pelmatidia ; avicularia sporadic, frequent.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* oval, large. *Orifice* wider than high, evenly rounded distally, with straight sides converging somewhat at the straight proximal margin. *Oral spine-bases* 6, occurring in two groups of three on the distal lateral margins of the orifice or as a distal and two lateral-distal pairs, the most proximal pair largest. *Frontal shield* flat, descending quite sharply at the inner ends of the *outermost intercostal spaces*, which are more often slot-like than rounded. *Costae* 25–30, each with 8 *lateral costal fusions* and *pelmatidia*. *Apertural bar* prominent, not very wide, straight, with a small, median ridge on the frontal shield. *Gymnocyst* exposed proximally, and to a less extent laterally, to the frontal shield : no *interzooecial secondary tissue.*

Avicularia fairly frequent, sporadic, probably proximally directed, unpaired, placed near the distal-lateral margins of the orifice, relatively large, oval or slightly pointed proximally.

Ovicells probably hyperstomial, apparently globular, prominent, and overlapping on to the distally adjacent zooecium : seen only in the broken state.

MEASUREMENTS. *Zooecia* Lz = 1·18 to 1·45 mm.
lz = 0·66 to 0·81 mm.
hr = 0·21 mm.
lr = 0·24 to 0·33 mm.
Avicularia Lz = 0·27 to 0·36 mm.
lz = 0·21 mm.

REMARKS. Gregory (1894 : 62) incorrectly assigned D.180 to the genus *Membranipora* and to the Jurassic (to the Calcaire à polypiers, Bathonian, of Ranville, Normandy, France). Lang (1916*a* : 96 and 1922 : 197) recognized its cribrimorph structure and its very probable Maastrichtian age and referred it to his genus *Rhiniopora.*

Both *Rhiniopora jurassica* (Gregory) and *R. scabra* Lang are synonymous and belong to *Castanopora.* It is unfortunate that the misleading specific name *jurassica* has to be retained for a Cretaceous species but, as it has priority of publication, no other course is open.

In the holotype (D.14207) of *Rhiniopora scabra* Lang only one complete zooecium is preserved. The dimensions of this and surrounding broken zooecia are the same as those of the two available specimens (D.180 and D.3313) of *Castanopora jurassica* (Gregory), and the number of costae in the complete zooecium of D.14207 is 25, each with 8 lateral costal fusions and pelmatidia, compared with 25–30 costae each with 8 lateral costal fusions and pelmatidia in D.180 and D.3313. The measurements given by Lang for *R. scabra* and *R. jurassica* were overestimated, and, in the latter species, avicularia are fairly frequent rather than "few". Lang stated "the preservation, both of the type specimen [D.180] and of D.3313 is but poor; the detailed characters, therefore, are uncertain". However, careful examination of



FIGS. 108, 109. *Castanopora jurassica* (Gregory).

(108) D.180. Holotype. Adult zooecium and parts of five adjacent, worn zooecia and worn, slightly pointed, probably proximally directed avicularium.

(109) D.14207. Holotype of *Rhiniopora scabra* Lang. Adult zooecium and parts of six adjacent worn zooecia.

the more complete zooecia in these specimens reveals most of the diagnostic characters.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian.

SPECIMENS. D.180. Holotype—see above.

D.3313. Large fragment of worn and damaged zoarium partly embedded in matrix. Maastrichtian. Maastricht, Holland. Old Collection.

D.14207. Holotype of *Rhiniopora scabra* Lang. Worn zoarial fragment. Horizon given as " Senonian, zone of *B. mucronata*, Rügen ", but probably boreal Maastrichtian. Agnes Laur Collection.

10. ***Castanopora glandulosa*** Lang

(Pl. 18, fig. 1 ; Text-fig. 110)

1916*a* *Castanopora glandulosa* Lang, pp. 95, 96.
1922 *Castanopora glandulosa* Lang : Lang, p. 219, pl. 5, fig. 3 ; text-fig. 70,
1923 *Castanopora glandulosa* Lang : Waters, pp. 548, 558, 560.
1930 *Castanopora glandulosa* Lang : Voigt, p. 551.

HOLOTYPE. D.15009, zoarial fragment. Horizon given as " Senonian, zone of *B. mucronata*, Rugen "—probably boreal Maastrichtian. Agnes Laur Collection.

EMENDED DIAGNOSIS. *Castanopora* with very flatly arched frontal shield ; orifice generally of equal height and width ; oral spine-bases 5 (occasionally 4) ; costae 24–27, each with 6–8 (often 7) lateral costal fusions and pelmatidia ; avicularia frequently paired, probably all distally directed, elongate and pointed.

DESCRIPTION. *Zoarium* encrusting or erect, unilaminar. *Zooecia* elongate-oval. *Orifice* generally as wide as high, relatively large, evenly rounded distally and laterally and without proximal-lateral constrictions, the proximal margin straight. *Oral spine-bases* 5 (occasionally 4), all relatively small, but the lateral pair tend to be larger, a fifth distal median oral spine is sometimes present. *Frontal shield* evenly, but flatly, arched. *Costae* 24–27, each with 6–8 (often 7) *lateral costal fusions* and an equivalent number of *pelmatidia*. *Apertural bar* flat, relatively wide with a slight thickening at the ends, triangular, the apex pointing proximally ; no median ridge or furrow. *Gymnocyst* only slightly exposed, mainly proximally to the frontal shield, but largely concealed owing to the closeness of the zooecia : no interzooecial secondary tissue present.

Avicularia of one kind : frequently, but not always, paired, one on each side of the orifice near the distal-lateral margins. There are apparently no proximally directed avicularia. In the one specimen available, all the avicularia are broken, but they were very probably of the same type as in *C. castanea*. It is possible to see the broken proximal ends of some avicularia ; one of these possesses constrictions marking the position of a broken *transverse bar*. All the evidence suggests that all the avicularia are distally directed. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·93 to 1·12 mm.
lz = 0·50 to 0·64 mm.
hr = 0·18 mm.
lr = 0·18 mm.
Avicularia Not measurable.

REMARKS. Only the holotype, a zoarial fragment, is available. In his diagnosis of this species Lang (1922 : 219) suggested that occasionally proximally directed avicularia may be present : I cannot see any indications of these. The zooecial length was slightly overestimated as 1·25 to 1·50 mm., although the generally larger size of this species is still apparent from the measurements given above. The range in number of costae (24–27) is generally above that of other species. The zoarium may have been encrusting. Lang stated that the number of oral spines is 4, and in the ensuing description he referred to 5 : both numbers occur as described, but the latter arrangement is more frequent.



FIG. 110. *Castanopora glandulosa* Lang, D.15009. Holotype. Distal half of adult zooecium with five oral spines and paired, distally pointed and directed avicularia : that on the left is worn, that on the right is broken at its distal end.

STRATIGRAPHICAL DISTRIBUTION. Probably boreal Maastrichtian.

SPECIMEN. D.15009. Holotype—see above.

11. *Castanopora aviculosa* (Lang)

(Pl. 19, fig. 3 : Text-fig. 111).

1916*a* *Rhiniopora aviculosa* Lang, p. 96.
1922 *Rhiniopora aviculosa* Lang : Lang, p. 184, text-fig. 58.
1929 *Rhiniopora aviculosa* Lang : Lang, p. 438, text-fig. 39 (19).
1930 *Rhiniopora aviculosa* Lang : Voigt, p. 551.

HOLOTYPE. In the Collection of Mons. F. Canu. Photograph in the British Museum (Natural History). Maastrichtian ; Maastricht, Limbourg, Holland.

EMENDED DIAGNOSIS. *Castanopora* with 24–26 costae ; avicularia proximally directed, paired, placed distal-laterally, elongate with sharply pointed rostra.

Description. *Zoarium* unilaminar, encrusting or erect. *Zooecia* broadly oval. *Orifice* evenly rounded distally and laterally, proximal margin straight, with distinct, pointed, proximal-lateral constrictions. *Oral spine-bases* of uncertain number, but at least 4, somewhat enlarged. *Frontal shield* flatly arched, variable in length. *Costae* 24–26, each generally with 6, occasionally 7 *lateral costal fusions*. *Outermost intercostal spaces* slot-like and narrow. *Apertural bar* not very wide, form obscure, but probably triangular. *Gymnocyst* not clearly visible, largely obscured by the avicularia; the zooecia are, however, well spaced: little or no *interzooecial secondary tissue*.

Avicularia of one form: all proximally directed, paired near the distal-lateral margins of the orifice; other occasional, similar, sporadic forms may occur: a *transverse bar* divides a small, semicircular, proximal portion from a large, elongate, sharply pointed, triangular *rostrum*. *Ovicells* not seen.



Fig. 111. *Castanopora aviculosa* Lang. Drawing from the photograph reproduced in Pl. 17, fig. 6. Scale approximate.

Measurements. (According to Lang, 1922 : 184.)
Zooecia Lz = 1·25 to 1·50 mm.
(No other measurements available.)

Remarks. Lang's figure and description of *Rhiniopora aviculosa* (1922 : 184, text-fig. 58), based on photographs of the holotype, are misleading. These photographs are not sufficiently clear to show the exact number of oral spine-bases. In MS. notes on the photographs Lang refers to 6 oral spines as present but remarks "sometimes five". He referred also to dimorphic avicularia "consisting of those with a very long and sharp rostrum, and those whose rostrum is comparatively short and not so sharp; the former are numerous, sporadically distributed, variously, but generally proximally, directed; the latter are more numerous, sporadically distributed, and probably always proximally directed". All the definitely pointed, elongate avicularia visible in the photographs are proximally directed; the broken and worn bases of this type are the "comparatively short and not so sharp" avicularia of Lang. The avicularia are distinctly paired, with only occasional, sporadic forms.

Nowhere are condyles visible in the broken avicularian apertures. A transverse bar is visible in one complete, pointed avicularium, and the lateral constrictions visible in other elongate avicularia are the broken remains of such transverse bars.

Rhiniopora aviculosa Lang is here regarded as a species of *Castanopora* characterized by proximally directed and pointed avicularia throughout the zoarium.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian.

SPECIMENS. None in the collections—only photographs of part of the holotype zoarium.

12. ***Castanopora magnifica*** (d'Orbigny)

(Pl. 18, figs. 2–7; Pl. 19, figs. 1, 2, 4–6; Pl. 20, fig. 1; Text-figs. 112–116)

1851 *Escharipora magnifica* d'Orbigny, pl. 686, figs. 1–5.
1852 *Escharipora pretiosa* d'Orbigny, pp. 227, 235; 1854, p. 1097.
1860 *Escharipora pretiosa* d'Orbigny: Coquand, p. 181.
1865 *Escharipora pretiosa* d'Orbigny: Ubaghs, p. 51.
1887 *Cribilina asperula* Marsson, pp. 97, 109, pl. 10, fig. 8.
1890 *Cribrilina asperula* Marsson: Osswald, p. 108.
1895 *Cribillina* [*sic*] *asperula* Marsson: Deecke, p. 79.
1895 *Cribrillina* [*sic*] *asperula* Marsson: Deecke, p. 87.
1900*b* *Cribrilina asperula* Marsson: Canu, p. 445.
1900*b* *Cribrilina magnifica* (d'Orbigny): Canu, p. 447.
1913 *Cribrilina cacus* Brydone, p. 437, pl. 14, figs. 6–8.
1916*a* *Castanopora magnifica* (d'Orbigny) Lang, pp. 94, 95.
1916*a* *Rhiniopora aspera* Lang, pp. 96, 97.
1916*a* *Rhiniopora asperula* (Marsson) Lang, pp. 96, 97.
1916*a* *Rhiniopora cacus* (Brydone) Lang, pp. 96, 97.
1916*a* *Rhiniopora horrida* Lang, pp. 96, 97.
1917 *Rhiniopora aspera* Lang: Lang, p. 172.
1922 *Rhiniopora aspera* Lang: Lang, p. 118, pl. 4, fig. 4; text-fig. 59.
1922 *Rhiniopora asperula* (Marsson): Lang, p. 190, pl. 4, fig. 5; text-fig. 60.
1922 *Rhiniopora cacus* (Brydone): Lang, p. 193.
1922 *Rhiniopora horrida* Lang: Lang, p. 194, pl. 4, fig. 6; text-fig. 61.
1922 *Castanopora magnifica* (d'Orbigny): Lang, p. 212.
1925 *Cribrilina asperula* Marsson: Levinsen, p. 376.
1925*c* *Cribrilina asperula* Marsson: Voigt, p. 176.
1929 *Rhiniopora aspera* Lang: Brydone, p. 13.
1929 *Rhiniopora asperula* (Marsson): Lang, p. 438, text-fig. 39 (20).
1929 *Rhiniopora horrida* Lang: Lang, p. 438, text-fig. 39 (21).
1930 *Rhiniopora cacus* (Brydone): Voigt, pp. 511, 543, 561, pl. 31, fig. 3.
1930 *Rhiniopora aspera* Lang: Voigt, p. 511.
1930 *Rhiniopora asperula* (Marsson): Voigt, pp. 511, 543, 561.
1930 *Rhiniopora horrida* Lang: Voigt, pp. 511, 561, pl. 31, fig. 5.
1935 *Rhiniopora aspera* Lang: Bassler, p. 190.
?1940 *Cribrilina magnifica* (d'Orbigny): Gillard, p. 15.
1953 *Rhiniopora aspera* Lang: Bassler, p. G193, text-fig. 144, 25.
1953 *Rhiniopora aspera* Lang: Illies, pp. 76, 91, 94.
1953 *Rhiniopora asperula* (Marsson): Illies, p. 97.
1953 *Rhiniopora cacus* (Brydone): Illies, pp. 76–101, pl. 15, figs. 1–6, pl. 16, figs. 1–2.

Lectotype. (Chosen by Lang, 1922 : 212.) The specimen figured by d'Orbigny, 1851, pl. 686, fig. 2. Maastrichtian of Sainte-Colombe (Manche), and Royan (Charente Inférieure). d'Orbigny Collection.

Emended diagnosis. *Castanopora* with relatively large orifice ; oral spine-bases 4 or 5 ; costae 14–30 (often about 20), each with 4–8 (often 5–7) lateral costal fusions and pelmatidia ; avicularia oblique, proximally directed, occasional, sporadic, somewhat variable in size, rounded.

Description. *Zoarium* encrusting or erect, unilaminar. *Zooecia* oval, large. *Orifice* may be wider than high, or of equal height and width, relatively large, evenly rounded distally, sides approximately straight and slightly converging proximally, proximal margin straight, slight proximal-lateral constrictions may be present. *Oral spine-bases* generally 4, less frequently 5, the lateral pair more thickened than the distal pair, a fifth, slender, median, distal oral spine may be present. Where only 4 oral spines are present there is seldom enough space between the distal pair for another single oral spine to develop. *Frontal shield* generally flatly arched. *Costae* of very variable number (from 14 to 30 according to Illies, 1953). In the specimens examined the range was found to be from 14 to 24, taking account of adult individuals only. Each costa generally bears from 4 to 8 (commonly 5–7) *lateral costal fusions* and *pelmatidia*, but there are fewer on the shorter costae close to the apertural bar. *Outermost intercostal spaces* more rounded than slot-like. *Apertural bar* prominent, not very wide, but thickened, especially near the outer ends ; a median ridge is present on the apertural bar and it tends to be prolonged on the frontal shield as an irregular, slight, median ridge producing a narrow median area of fusion. *Gymnocyst* more prominently exposed proximally to the frontal shield than on its lateral margins, not concealed by *interzooecial secondary tissue*. *Communication pores* are seen as two or three shallow, circular depressions in the lateral walls of some zooecia, and as single, larger depressions in the proximal and distal walls, the centre of each depression being perforated by a small, round pore.

Avicularia of one kind : of variable size and occurrence, approximately twice as long as wide, prominent, fairly frequent, proximally directed, oblique, divided by a *transverse bar* into a somewhat smaller proximal portion and a larger, rounded or slightly pointed *rostrum*. Their appearance alters considerably with the degree of wear.

Ovicells rare, hyperstomial, prominent, large, globular, overlapping on to the proximal part of the frontal shield of the distally adjacent zooecium. Each bears two ridges, which converge distally at the mid-line from the proximal-lateral corners. The lateral pair of oral spines may flank the opening of the ovicell.

Measurements. *Zooecia* Lz = 0·75 to 1·48 mm.
lz = 0·45 to 1·20 mm.
hr = 0·15 to 0·33 mm.
lr = 0·15 to 0·30 mm.
Avicularia Lz = 0·17 to 0·25 mm.
lz = 0·09 to 0·18 mm.

The highest values of length and breadth given by Illies (1953) are Lz = 1·80 mm.



112



114



Scale as for Fig. 112



116



115

and lz = 1·7 mm. These unusually high values possibly include measurement of abnormally long and wide zooecia which occur occasionally in this species.

REMARKS. Voigt (1930 : 511) regarded a free specimen, which he assigned to Marsson's species *asperula*, as synonymous with *Rhiniopora aspera* Lang. Marsson (1887 : 97) originally included both free and encrusting forms in his species *Cribrilina asperula.*

Lang (1922 : 192) suggested that Marsson's encrusting specimens were *Rhiniopora aspera* which was distinguished by Lang from *R. asperula* by its supposedly different growth habit, its slightly greater number of costae, and the occurrence in *R. aspera* of " generally five but occasionally only four " oral spine-bases compared with " 4 considerably thickened " oral spine-bases in *R. asperula.* In the present re-examination of the specimens named *R. aspera* by Lang, it was found that 4 oral spine-bases, rather than 5, are more frequently present. The range in number of costae was given by Lang as 18–20 for *R. aspera* and 20–24 for *R. asperula.*

Only one specimen (D.14176) labelled by Lang " *R. asperula* (Marsson) " is available : 4 oral spine-bases are present and the range in number of costae is 19–22. In the re-examined specimens labelled " *R. aspera* " the range in number of costae is 14–20. Thus the differences between the species are very slight and it is difficult to separate them.

Recently (1953), a statistical variation-study of 40 zoaria of a species placed in *Rhiniopora* [*R. cacus* (Brydone)], from the Maastrichtian of Hemmoor, Germany, has been carried out by Illies. This careful analysis of characters which can be studied quantitatively (those which can be measured or counted) has shown in particular that the number of costae in each frontal shield varies widely, both between individuals of the same colony and individuals of separate zoaria. If anything, the length and breadth measurements of zooecia show the least variation.

Illies (1953 : 99) summarized her work as follows : " 40 colonies of the bryozoan *Rhiniopora cacus* (Brydone) from the chalk of the Maastrichtian of Hemmoor (Niederelbe) were investigated by variation-statistical methods.

" In the first section, the variation of quantitative characters within a single

---

FIGS. 112–116. *Castanopora magnifica* (d'Orbigny).

(112) D.31630. A very well-preserved single adult zooecium. Distal and lateral communication pores are well displayed.

(113) D.15620. Holotype of *Rhiniopora aspera* Lang. Distal end of adult zooecium with five oral spines and large oval avicularium with narrow transverse bar.

(114) D.14176. Labelled by Lang " *Rhiniopora asperula* (Marsson) ". Adult zooecium and parts of three distally adjacent zooecia and one worn, oval avicularium. The gymnocyst is typically widely exposed.

(115) N.C.M.76.937 (25/1). Distal end of adult zooecium with five oral spines and worn oval avicularium.

(116) D. 15591*d*. One of five fragments of the same zoarium labelled by Lang " *Rhiniopora cacus* (Brydone) ". Adult zooecium, parts of four adjacent zooecia and two worn, interzooecial avicularia.

colony are examined. From this, it can be shown that a number of characters in the central area of the zoarium vary differently from the characters in the younger (outer) zones. Whereas length-breadth measurements increase in value and lateral costal fusions, costae etc. increase in number in the first four growth-rings, they remain more or less constant in the outer rings.

" The comparison of several zoaria—in the second section—based on the outer rings only, gave wider limits to the variations than has previously been acknowledged. The number of costae varies greatly, so that their taxonomic importance must now be regarded as less than that of the length and breadth measurements—which are seen to be of the greatest value " [translation].

In a preceding section on the limits of variation of the species, Illies listed the extremes of length-breadth measurements of zooecia and the following characters :

Lateral costal fusions	4–8, mainly 5–7.
Form of orifice	Normal, subcircular or super-normal[1] (the last form being found in all colonies).
Oral spines	Some colonies always with 5, others with 4–5.
Apertural bar	Possesses lateral " swellings " of varying strength in different colonies.
Gymnocyst	Varies in breadth.
Avicularia	Vary from 5 to 66 per 100 zooecia.
Median area of fusion	May be " rib-like ", flattened, or not particularly distinct.
Length-breadth ratio	Varies considerably . . . indicating irregular growth.
Ovicells	Rare, those found, all developed to the same degree.

[1] Lang, 1921, p. xlvii, and p. 34 here.

Except for the statements on avicularia and ovicells this revision is confirmed by an examination of the material available in the British Museum (Natural History) and other collections mentioned.

Illies (p. 97) referred to Voigt's (1930) suggestion of re-combining *Rhiniopora aspera* Lang with *R. asperula* (Marsson). In her opinion " such a synonymy is only justified if an encrusting form of *asperula* is also found with, apart from the same measurements, less than twenty costae and with thin oral spines. The statistical investigation of variation in *R. cacus* (Brydone) has shown, however, that the variation of definitive characters is far greater than was previously recognised in the literature, so that it may be presumed that in a similar investigation of *R. aspera* and *R. asperula* the present limits are like-wise found to be too narrow and a merging of *aspera*—and even *cacus*—to *asperula* may be necessary " [translation].

It is apparent that for *R. cacus* the results gained from more usual combined qualitative and quantitative investigations are very similar to those achieved by a statistical study of those characters which may be measured or counted.

In the material examined here Illies' revision of the characters of *R. cacus* is confirmed, and this species is regarded as synonymous with *Rhiniopora aspera* Lang, in its turn a growth form of *Rhiniopora asperula* (Marsson). *R. horrida* is, as stated by Lang (1922 : 194), like *R. cacus*, except for its growth habit, and is also, therefore, included in the synonymy.

But *R. asperula is itself a junior synonym of Castanopora magnifica* (d'Orbigny). Mons. E. Buge has kindly provided a photograph (Pl. 20, fig. 1) [by Mons. P. Mémin of the Serv. Photo. Géol., E.N.S.] of the holotype (?) of d'Orbigny's species as well as the following details : " Costules : 24–26, 4 épines orales, aviculaires oraux à bec dirigé vers la partie proximale de la zoécie ". The characters given, as well as those shown in the photograph, clearly justify the synonymy of the species.

Apparent growth habit is not here regarded as of diagnostic value. Accepting this, the ranges in number of costae, lateral costal fusions and the pelmatidia and oral spines of these species (i.e. *Castanopora magnifica*, *Rhiniopora asperula*, *R. aspera*, *R. cacus*, *R. horrida*) all overlap, or are similar, and are contained within the limits established by Illies on statistical evidence for *R. cacus*. Qualitative characters, such as degree of arching of the frontal shield, shape and arrangement of avicularia, and width of exposed gymnocyst, all appear to vary within the same relatively wide limits, both in zooecia of the same colony and between zooecia of separate colonies.

The following specimens (labelled by Lang as *Rhiniopora aspera*) have been re-examined ; they include the holotype and 5 paratypes of this species and are here assigned to *Castanopora magnifica* (d'Orbigny).

In D.15620 the better preserved zooecia were measured, Lz ranges from 0·78 to 0·90 mm., lz from 0·45 to 0·50 mm. ; the range in number of costae varies from 14 to 16 and the number of lateral costal fusions varies from 4 to 6, oral spine-bases generally number 4, infrequently 5 may be present. In D.14153, D.14210, D.15383, D.16675 and D.16677—all from Rügen—the range in number of costae is 15–20. D.14210 and D.16677 have well exposed gymnocysts. In these specimens only 4 oral spine-bases are developed on each zooecium ; compared with those of D.15620 these are all slightly thickened. D.15302 and D.15324, also from Rügen, are very badly damaged and worn.

D.31630 is a well-preserved zooecium not included in Lang's 1922 Catalogue. Seventeen costae are present, each with 5 lateral costal fusions and pelmatidia ; there are 4 oral spine-bases. The lateral, proximal and distal walls are completely exposed displaying communication pores. The dimensions of the zooecium are : Lz = 1·15 mm., lz = 0·66 mm., hr = 0·24 mm., lr = 0·21 mm.

The holotype of *Rhiniopora horrida* Lang, D.14171, possesses only four whole zooecia, each with four oral spine-bases, the lateral pair in each case being thicker than the distal pair, a condition common in the other specimens here referred to *Castanopora magnifica* (d'Orbigny). All the oral spine-bases in D.14171 tend to be slender, and the zooecia are generally less robust than in the encrusting forms, probably reflecting the growth habit (cf. p. 22 for remarks on the growth habit of *Steganoporella buski* Harmer). The range in number of costae in D.14171 is 19–21 with 5–6 lateral costal fusions on each costa. D.15020 is much broken and very worn. The zooecia are over 1·00 mm. long. It is not possible to determine the specific characters, and its placing, by Lang, in this species is very doubtful.

D.15591 *a–e*, labelled by Lang as *R. cacus* (Brydone), is an imperfectly preserved, fragmented zoarium encrusting five pieces of echinoid test. In fragment *b* three abnormal zooecia are present, two being very wide and one unusually long. The measurement of zooecia such as these may account for the somewhat high values recorded by Illies for *R. cacus*. Several zooecia in fragment *a* possess broken ovicells.

Poorly preserved avicularia are present in D.15591, D.14167 and D.16676—the last two are specimens from Rügen also labelled by Lang as *R. cacus*.

Four specimens from the Norwich Castle Museum collections—N.C.M.2194 (3), 76.937 (25/1), 76.937 (26/1) and 112.22 (1/1)—belong to *Castanopora magnifica* (d'Orbigny). The range in number of costae is 16–21, with 4, occasionally 5, oral spine-bases and from 4 to 6 lateral costal fusions and pelmatidia on each costa. N.C.M.76.937 (26/1) has less thickened oral spine-bases than is usual.

A very well preserved zoarial fragment, D.40673, encrusting a piece of echinoid test, from a recently exposed outcrop of the *O. lunata* Chalk at Mundesley, Norfolk, is also assigned to *Castanopora magnifica* (d'Orbigny).

In addition, the three zoarial fragments figured by Brydone (1913 : 437, pl. 14, figs. 6–8) as *Cribrilina cacus* have been re-examined. S.M., B.36264, that shown in fig. 6, was placed by Lang (1922 : 188) in *Rhiniopora aspera*. S.M., B.36265 (fig. 7 of Brydone) was chosen by Lang (p. 193) as lectotype of *Rhiniopora cacus* (Brydone), to which species Lang also assigned S.M., B.36266 (fig. 8 of Brydone). These specimens all belong to *Castanopora magnifica* (d'Orbigny) ; their dimensions and other details are as follows :

Zooecia Lz = 1·18 to 1·45 mm.
lz = 0·68 to 1·20 mm.
hr = 0·25 to 0·31 mm.
lr = 0·22 to 0·30 mm.
Costae 20–24.
Lateral costal fusions 5–8.
Oral spine-bases 4 or 5.
Avicularia Lz = 0·17 to 0·25 mm.
lz = 0·10 to 0·18 mm.

It is evident that the specimens are generally larger than most of the material which is here assigned to *C. magnifica* (d'Orbigny) ; they all, however, fall well within the limits of variation established by Illies (1953) and confirmed here by qualitative as well as quantitative study.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *Belemnitella mucronata*, and Maastrichtian, zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk).

SPECIMENS. D.15620. Zoarial fragment. Holotype of *Rhiniopora aspera* Lang. Maastrichtian, zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk). Trimingham, Norfolk. A. C. Savin Collection.

D.14153, D.14210, D.15383, D.16675, D.16677. Five paratypes of *Rhiniopora aspera* Lang. Zoarial fragments encrusting pieces of echinoid test. Horizon given as " Senonian, zone of *B. mucronata* ", but probably boreal Maastrichtian. Rügen. Agnes Laur Collection.

D.14171. Holotype of *Rhiniopora horrida* Lang. Small zoarial fragment. Horizon locality and collection as for D.14153.

D.14176. Zoarial fragment labelled by Lang " *Rhiniopora asperula* (Marsson) ". Horizon, locality and collection as for D.14153.

D.15020. Very worn and damaged zoarial fragment labelled by Lang " *Rhiniopora horrida* ". Horizon, locality and collection as for D.14153.

D.15302, D.15324. Broken and worn fragments of two zoaria labelled " *Rhiniopora aspera* Lang ". Horizon, locality and collection as for D.14153.

D.15591*a–e*. Five fragments of the same zoarium encrusting pieces of echinoid test. Horizon, locality and collection as for D.15620. The specimen was labelled by Lang " *Rhiniopora cacus* (Brydone) ".

D.14167, D.16676. Two small zoarial fragments labelled by Lang " *Rhiniopora cacus* (Brydone) ". Horizon, locality and collection as for D.14153.

D.31630. A well-preserved single zooecium. Senonian, zone of *B. mucronata*. Norwich, Norfolk. T. H. Withers Collection.

D.31707. Small zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *B. mucronata*. Pit a quarter mile south-south-west of Coltishall, Norfolk. T. H. Withers Collection.

D.40673. Zoarial fragment encrusting a piece of echinoid test. Maastrichtian, zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk). Exposed on foreshore, 1954, 100 yards south of the Kiln Camping Site, Mundesley, Norfolk. D. R. Howlett Collection.

OTHER MATERIAL. N.C.M.76.937 (25/1). Incomplete zoarium encrusting a whole test of *Cardiaster ananchytis* Leske. Senonian, zone of *B. mucronata*. Catton, near Norwich, Norfolk. R. M. Brydone Collection.

N.C.M.2194(3). Incomplete zoarium encrusting a whole echinoid test labelled " *Epiaster gibba* S. Woodward ". Senonian, zone of *B. mucronata*. Collection unknown.

N.C.M.76.937(26/1). Zoarial fragment encrusting a whole test of *Cardiaster ananchytis* Leske. Senonian, zone of *B. mucronata*. Harford, near Norwich, Norfolk. R. M. Brydone Collection.

N.C.M.112.22(1/1). Incomplete zoarium encrusting a whole echinoid test labelled " *Galerites vulgaris* Lam.". Senonian, zone of *B. mucronata*. Thorpe-next-Norwich, Norfolk. Collection unknown.

S.M., B.36264–66. Three well-preserved zoarial fragments encrusting pieces of echinoid test. Maastrichtian, zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk). Trimingham, Norfolk. R. M. Brydone Collection.

S.M., B.36264—part of this specimen was figured by Brydone (1913 : 437, pl. 14, fig. 6) as *Cribrilina cacus*. Lang (1922 : 188) tentatively assigned the specimen to *Rhiniopora aspera*.

S.M., B.36265—part of this specimen was figured by Brydone (1913 : 437, pl. 14, fig. 7) as *Cribrilina cacus*. Lang (1922 : 193) chose it as lectotype of *Rhiniopora cacus* (Brydone).

S.M., B.36266—part of this specimen was figured by Brydone (1913 : 437, pl. 14, fig. 8) as *Cribrilina cacus*. Lang (1922 : 193) assigned it to *Rhiniopora cacus*.

13. *Castanopora faujasi* (von Hagenow)

1851 *Cellepora* (? *Dermatopora*) *faujasi* von Hagenow, p. 99, pl. 10, fig. 19*a*, *b* [the question-mark is omitted on p. 99].

1930 *Rhiniopora faujasi* (von Hagenow) Voigt, p. 511, pl. 31, fig. 2.

EMENDED DIAGNOSIS. *Castanopora* with orifice wider than high ; Lz about 1·00 mm., lz about 0·60 mm. ; costae 13–15, each with 6–7 lateral costal fusions ; gymnocyst widely exposed ; apparently no avicularia.

REMARKS. Von Hagenow's figure of this species (1851, pl. 10, fig. 19*b*) shows only broken zooecia lacking frontal shields. But the shape of the broken apertures strongly suggests a cribrimorph, with well-spaced zooecia which had wide orifices. Von Hagenow described the zoarium as a series of " quincuncially arranged cells with transversely oval, almost round, apertures, each with a semicircular emargination " [translation]. This " emargination " is undoubtedly the orifice lacking its proximal border, and it is thus an apparent continuity with the worn aperture of the zooecium.

Voigt (1930 : 511) assigned a specimen to the species ; complete frontal shields are present and it is clearly a *Castanopora*. Apart from the characters mentioned in the above diagnosis it is not possible to interpret other details from the figure or from Voigt's note on the specimen.

STRATIGRAPHICAL DISTRIBUTION. Horizon given as " Senonian, upper part of zone of *B. mucronata* ", but very probably Maastrichtian (Maastricht, Holland).

SPECIMENS. None in the collection.

14. *Castanopora voigti*[1] sp. nov.

1930 *Rhiniopora* cf. *hispida* Lang : Voigt, pp. 511, 561, pl. 31, fig. 6.

HOLOTYPE. The specimen figured by Voigt, 1930, pl. 31, fig. 6. Horizon given as " Senonian, upper part of zone of *B. mucronata*, Møen ", but since shown to be Maastrichtian, zone of *Belemnella lanceolata* (see below).

DIAGNOSIS. *Castanopora* with erect, cylindro-quadrate zoarium ; zooecia of the 4 adjacent rows alternate ; orifice higher than wide ; costae 20–25.

DESCRIPTION. (Based on Voigt's figure.) *Zoarium* erect, cylindro-quadrate, composed of 4 adjacent rows of alternating zooecia. *Zooecia* oval or rectangular. *Orifice* higher than wide, evenly rounded distally and laterally, proximal margin straight. *Oral spine-bases* of uncertain number, at least two pairs present. *Frontal shield* oval, very flatly arched, descending sharply near the outer ends of the costae. *Costae* 20–25, each with 4, possibly at least 5 *lateral costal fusions*. *Apertural bar* not very wide, slightly curved proximally. *Gymnocyst* well exposed, especially proximally to the frontal shield. *Avicularia* and *ovicells* not shown in the figure.

MEASUREMENTS. (According to Voigt.)
Zooecia Lz = 0·75 to 0·80 up to 1·00 mm.

REMARKS. Voigt (1930 : 511, pl. 31, fig. 6) briefly described a specimen from the same horizon as the holotype of *Rhiniopora hispida* Lang, but he only compared it with that species in view of the uncertainty of the characters given by Lang. *R. hispida* was founded on an extremely worn and badly damaged specimen (D.14996), in which all the frontal shields are missing except for a fragment of one. The characters given by Lang are largely hypothetical as he himself indicated by placing most of them in square brackets, a usage which he reserved for conjectural and uncertain statements. The name *Rhiniopora hispida* must therefore be discarded since the species cannot be satisfactorily defined.

---

[1] After Prof. Ehrhard Voigt.

C. voigti is distinguished from all other known species of *Castanopora* by its cylindro-quadrate zoarium. Further material may show the presence of avicularia and the number of oral spine-bases.

Voigt (in a letter dated 16th June, 1952) has indicated that the chalk of Møen, the locality of the holotype, is of Maastrichtian age belonging to the zone of *Belemnella lanceolata*. Lang (1921 : lxxxiv) had regarded the Møen chalk as Danian, and Voigt (1930 : 511) had originally placed it as Senonian, uppermost part of the zone of *Belemnitella mucronata*.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian, zone of *Belemnella lanceolata*.

SPECIMENS. None in the collection.

SPECIES OF UNCERTAIN SYSTEMATIC POSITION PREVIOUSLY ASSIGNED TO *RHINIOPORA* AND *CASTANOPORA*

(1) ***Cellepora perforata*** Quenstedt

1879 *Cellepora perforata* Quenstedt, p. 312, pl. 154, fig. 37.

REMARKS. Lang (1922 : 183) referred this species to *Rhiniopora*, but from Quenstedt's original description and figure it is possible to deduce only that the specimen was an encrusting cribrimorph with hyperstomial ovicells. It cannot be assigned with any certainty at all to *Castanopora* or any other genus, even though its horizon is recorded as Maastrichtian of Maastricht.

(2) ***Reptescharella radiata*** d'Orbigny

1852 *Reptescharella radiata* d'Orbigny, pl. 716, figs. 4–6.
1853 *Reptescharella radiata* d'Orbigny : d'Orbigny, p. 468.

REMARKS. Lang placed this species in *Rhiniopora*. In the original description of *Reptescharella radiata* d'Orbigny (1853 : 468) referred to the encrusting habit of the species and to the presence of 16 costae, and the orifice was described as " ovale, transverse " suggesting that lr exceeds hr in value. Further interpretation of the species must rest on d'Orbigny's figures (1852, pl. 716, figs. 4–6), which, however, are highly stylized ; no avicularia are shown, and the zooecia are closely placed with 16–18 costae each with 5–7 lateral costal fusions. The orifices are about twice as wide as high.

Without information on the nature of the oral spines, the avicularia and other details, it is not possible to state with certainty that the species belongs to *Castanopora*. The recorded horizon, Campanian and Santonian, is lower than for most species of *Castanopora*.

(3) ***Rhiniopora hispida*** Lang

1916*a* *Rhiniopora hispida* Lang, pp. 96, 97.

This species has been discussed on p. 236 under the remarks on *Castanopora voigti* sp. nov.

(4) ***Reptescharipora ornata*** d'Orbigny

1852 *Reptescharipora ornata* d'Orbigny, pl. 720, figs. 6–8.
1853 *Reptescharipora ornata* d'Orbigny : d'Orbigny, p. 494.
1854 *Reptescharipora ornata* d'Orbigny : d'Orbigny, p. 1098.

HOLOTYPE. d'Orbigny Collection, No. 8012, Laboratoire de Paléontologie, Muséum National d'Histoire naturelle, Paris. Senonian. Vendôme, (Loir et Cher), France.

REMARKS. Lang (1916*a* : 94 ; 1922 : 213) placed this species in *Castanopora*. Dr. E. Buge has kindly provided me with a photograph (by Mons. P. Mémin of the Serv. Photo. Géol. E.N.S.) of the holotype and details of its preservation. In a letter (8th December, 1954) he stated, " Costules nombreuses, mais presque indiscernables sur le spécimen. Epines non distinctes (probablement détruites). Aviculaires oraux (?) érodés, non définissables ".

The recorded horizon, Senonian [Santonian] of Vendôme, Loir et Cher, is again lower than for most species of *Castanopora* and it is most uncertain that the species belongs to that genus.

Genus ***UBAGHSIA*** Jullien

1826 *Celepora* : Goldfuss, pp. 26, 248 [*partim*].
1828 *Cellepora* : Morren, p. 34 [*partim*].
1832 *Cellepora* : Dumont, p. 360 [*partim*].
1834 *Cellepora* : von Klöden, pp. 264, 341 [*partim*].
1836 *Discopora* : Edwards, p. 248 [*partim*].
1840 *non Eschardoides* : Roemer, p. 14.
1846 *Cellepora* : Geinitz, p. 609 [*partim*].
1848 *Cellepora* : Bronn, p. 253 [*partim*].
1848 *Discopora* : Bronn, pp. 255, 431 [*partim*].
1849 *Discopora* : Bronn, p. 130 [*partim*].
1849 *Cellepora* : Geinitz, p. 248 [*partim*].
1850 *Escharina* : d'Orbigny, p. 262 [*partim*].
?1851 *Cellepora* (*Discopora*) : von Hagenow, p. 96 [*partim*].
1851 *Cellepora* (*Dermatopora*) : von Hagenow, p. 98 [*partim*].
1851 *Cellepora* (*Dermatopora*) : Bronn & Roemer, p. 102 [*partim*].
?1852 *Reptescharella* d'Orbigny, pls. 604, 715–716 [*partim*].
?1852 *Semiescharipora* d'Orbigny, pls. 717–719 [*partim*].
?1852 *Reptecharipora* d'Orbigny, pls. 719–720 [*partim*].
?1853 *Reptescharella* : d'Orbigny, p. 464 [*partim*].
?1853 *Semiescharipora* : d'Orbigny, p. 479 [*partim*].
?1853 *Reptescharipora* : d'Orbigny, p. 489 [*partim*].
?1854 *Semiescharipora* : d'Orbigny, p. 1097 [*partim*].
?1854 *Reptescharipora* : d'Orbigny, p. 1098 [*partim*].
?1857 *Reptescharella* : Pictet, p. 110 [*partim*].
?1857 *Semiescharipora* : Pictet, p. 112 [*partim*].

?1860 *Semiescharipora* : Coquand, p. 183.
1862 *Cellepora* : Goldfuss, p. 24 [*partim*].
1865 *Steginopora* : Ubaghs, p. 55.
1879 *Cellepora* : Quenstedt, p. 303 [*partim*].
1879 *Semiescharipora* : Ubaghs, p. 217 [*partim*].
?1879 *Lepralia* : Ubaghs, p. 221 [*partim*].
1881 *Semiescharipora* : Mourlon, p. 116 [*partim*].
?1881 *Lepralia* : Mourlon, p. 119 [*partim*].
1885 *Semiescharipora* : Vine, pp. 116, 156 [*partim*].
1885 *Cellepora* : Vine, p. 161 [*partim*].
1885 *Cellepora* (*Dermatopora*) : Vine, p. 164 [*partim*].
?1885 *Cellepora* (*Discopora*) : Vine, p. 164 [*partim*].
1886 *Steginopora* : Jullien, pp. 612, 614 [*partim—non S. irregularis* d'Orbigny, ?*non S. aculeata* d'Orbigny, *non S. pulchella* d'Orbigny, ?*non S. ornata* d'Orbigny].
1886 *Ubaghsia* Jullien, pp. 610, 616 [*partim—U. reticulata* only].
1887 *Cribrilina* : Marsson, pp. 96, 109 [*partim*].
1889 *Semiesharipora* : Ubaghs, p. 52.
1900*b* *Cribrilina* : Canu, p. 445 [*partim*].
1900*b* *Cribrilina* (*Costula*) : Canu, p. 450.
1900*b* *Steginopora* (*Ubaghsia*) : Canu, p. 455 [*partim—S.* (*Ubaghsia*) *reticulata* only].
1900*b* *Semiescharipora* : Canu, p. 457 [*partim*].
1900*b* *Reptescharipora* : Canu, p. 457 [*partim*].
1910 *Cribrilina* : Canu, p. 63 [*partim*].
1916*a* *Ubaghsia* : Lang, 93, 99.
1916*a* *Batrachopora* Lang, pp. 101, 110 [*partim*—?*B. ovalis* (d'Orbigny), ?*B. convexa* (d'Orbigny), ?[*B.*] *signata* (von Hagenow), *non B. royanensis* Lang].
?1916*a* *Cellepora* (*Discopora*) : Lang, p. 111.
1919*c* *Batrachopora* : Lang, p. 105.
1919*d* *Batrachopora* : Lang, pp. 192, 197, 199, 203, 208, 217, 223 [*partim*—?*B. ovalis* (d'Orbigny), ?*B. convexa* (d'Orbigny), ?[*B.*] *signata* (von Hagenow), *non B. royanensis* Lang].
?1919*d* *Cellepora* (*Discopora*) : Lang, p. 205.
1920 *Steginopora* : Canu & Bassler, p. 282, fig. 81 S.
1921 *Batrachopora* : Lang, pp. lvii, xc, xci [*partim*].
1921 *Ubaghsia* : Lang, pp. lxvi, xc, xci, xciii.
1922 *Ubaghsia* : Lang, p. 225.
1922 *Batrachopora* : Lang, p.357 [*partim*—?*B. ovalis* (d'Orbigny), ?*B. convexa* (d'Orbigny), ?[*B.*] *signata* (von Hagenow), *non B. royanensis* Lang].
1925 *Cribrilina* : Levinsen, p. 375 [*partim—C. brachiata* only].
1925*b* *non Batrachopora* : Voigt, pp. 99, 103.
1925*c* ?*Batrachopora* : Voigt, p. 175.
1927 *Cribrilina* : Voigt, p. 102 [*partim*].
1927 *Ubaghsia* : Canu & Bassler, p. 41.
1927 *Batrachopora* : Canu & Bassler, p. 55.
1929 *Steginopora* : Canu & Bassler, p. 243 [*partim—S. ocellata* only].
1929 *Ubaghsia* : Canu & Bassler, p. 244.
1929 *Batrachopora* : Lang, p. 439.
1929 *Ubaghsia* : Lang, p. 439.
1930 *Ubaghsia* : Voigt, pp. 513, 515, 538, 561.
1930 *Batrachopora* : Voigt, pp. 515, 561 [*partim—non B. ultima* Voigt].
1935 *Batrachopora* : Bassler, pp. 31, 55.
1935 *Ubaghsia* : Bassler, pp. 31, 222.
1939 *non Batrachopora* : Voigt, p. 93.
?1940 *Cribrilina* : Gillard, p. 15 [*partim*].

1949 *non Batrachopora* [*Escharoides*]: Voigt, pp. 41, 43.
1953 *Batrachopora*: Bassler, p. G192.
1953 *Ubaghsia*: Bassler, p. G193.

TYPE SPECIES. *Steginopora reticulata* Ubaghs, 1865 : 55. Maastrichtian. Ober Bryozoenschichte (horizon *f* of Ubaghs). Valkenburg (Fauquemont), east of Maastricht, or Geulem, north-east of Maastricht, Holland.

EMENDED DIAGNOSIS. Pelmatoporinae with relatively large zooecia; branched or unbranched distal oral spines; proximal and distal oral shields present, the latter may be less developed; the former composed of paired, large, proximal-lateral oral avicularia, raised on stout stalks, united above the secondary orifice and often fused on their distal sides with the distal-lateral oral spines, the distal oral shield less prominent, formed by the enlarged, sometimes branched, often fused, distal oral spines and the upgrowth of secondary calcareous tissue on the rim of the secondary orifice; apertural bar not involved in the proximal oral shield; each costa with two or more pelmata; secondary tissue forming a tertiary front wall, those of adjacent zooecia usually united (the tertiary front wall is frequently destroyed during fossilization); interzooecial avicularia may occur.

REMARKS. *Ubaghsia* Jullien and *Batrachopora* Lang were regarded by Lang as distinct genera which he placed in separate subfamilies of the Pelmatoporidae, the former in the Castanoporinae (1922 : 154) and the latter in the Pelmatoporinae (1922 : 235). The two subfamilies are here merged: see remarks on the subfamily Pelmatoporinae.

Neither pelmata nor pelmatidia were indicated in the original figures or descriptions of *Ubaghsia* given by Ubaghs (1865) and Jullien (1886), but other characters, which were clearly shown, are very closely comparable with those of *Batrachopora* Lang.

The diagnoses of *Ubaghsia* and *Batrachopora* given by Lang (1916*a* : 93, 101) and later slightly modified (1922 : 225, 358) show that he distinguished between the genera on the basis that the former possessed pelmatidia and branched distal oral spines, while the latter had pelmata and unbranched distal oral spines. The distinction between pelmata and pelmatidia is not a valid one, nor should the branching of oral spines be considered of generic importance. Further, it is very doubtful whether *Ubaghsia reticulata* has branched spines (cf. Jullien, 1886, pl. 18, fig. 4; and Voigt, 1930, pl. 31, fig. 19). Thus *Ubaghsia* and *Batrachopora* are synonymous, as Voigt (1930 : 515) has already suggested might be the case.

In the present revision, all the available material, previously assigned to *Ubaghsia* and *Batrachopora*, has been re-examined, and two new species of *Ubaghsia* are described.

Where specimens have been available, the species have been distinguished on general zooecial and avicularian characters rather than on the form of the oral shields which are mainly of generic significance. The form of the proximal oral shield is very often much modified by wear, but the original presence or absence of this large secondary structure can usually be determined even in well-worn specimens.

Lang (1922 : 358), in his remarks on *Batrachopora*, was concerned principally with the evolutionary relationships of the species which he recognized as belonging to the genus. Several of these species, and others later assigned to *Batrachopora* by Voigt, are of very uncertain systematic position (see p. 264); they are not accepted here as species of *Ubaghsia*.

Jullien (1886 : 614, pls. 17–20) described and figured several species which he assigned to *Steginopora* d'Orbigny, and erected the genus *Ubaghsia* to include *Steginopora reticulata* Ubaghs and a new species, *Ubaghsia arcifera*. Lang (1916*a* : 99, 100; 1922 : 225) regarded all these species as belonging to *Ubaghsia*, with the exception of *U. arcifera* which he placed in *Phrynopora* (Lang, 1916*a* : 97, 98; 1922 : 201). If the species of uncertain systematic position, previously assigned to *Batrachopora*, are excluded from *Ubaghsia*, and if Jullien's species are taken into account, much of Lang's suggested evolution of the genera must also be questioned.

It is possible that *Ubaghsia* Jullien is synonymous with *Disteginopora* d'Orbigny and *Steginopora* d'Orbigny, which apparently differ from one another only in their zoarial growth-habit. The tertiary front wall of *Ubaghsia ocellata* and *U. demorgani* appears to be somewhat similar to that found in *Steginopora ornata* d'Orbigny and *Disteginopora horrida* d'Orbigny. However, it is difficult to interpret species from d'Orbigny's stylized figures and inadequate descriptions. Until the holotypes, or new topotype material, of d'Orbigny's species can be examined it is better to retain *Ubaghsia* Jullien as a separate genus.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zones of *M. cor-anguinum*, *G.* [*A.*] *quadrata*, *B. mucronata* and Maastrichtian.

KEY TO THE SPECIES OF *UBAGHSIA* JULLIEN

- (I) *Ubaghsia* with orifice much smaller than the frontal shield.
  - (A) Oral spines unbranched.
    - (1) Avicularia of one kind only—a large oral pair.
      - (*a*) 8–10 costae, each with 3 lateral costal fusions, Lz = 0·70 to 0·80 mm., lz = 0·50 to 0·57 mm., oral avicularia fused above secondary orifice to give a complete proximal oral shield. . 1. ***U. ranunculus*** (Lang)
      - (*b*) 18–22 costae; proximal oral shield apparently incomplete; frontal shield distinctly pointed proximally . . . 2. ***U. aurita*** (Lang)
    - (2) Avicularia of two kinds: (*a*) a large oral pair, (*b*) an interzooecial or vicarious, paired or unpaired, small form.
      - (*a*) Shape of orifice variable: wider than high or of equal height and width.
        - (i) Zooecia larger, 9–10 costae, each with 2 lateral costal fusions; interzooecial avicularia small, rounded, unpaired . 3. ***U. gasteri*** sp. nov.
        - (ii) Zooecia shorter, 7–10 costae, each with 2, more often 3, lateral costal fusions; interzooecial avicularia larger, pointed . 4. ***U. langi*** sp. nov.
      - (*b*) Orifice always wider than high.
        - (i) 7 costae, each with 2, occasionally 3, lateral costal fusions; zooecia over 1·00 mm. long . . . . . . 5. ***U. crassa*** (Lang)
        - (ii) 10–14 costae, each with 3 or 4 lateral costal fusions.
          - (α) Zooecia over 1·00 mm. long . . . . 6. ***U. coaxans*** (Lang)
          - (β) Zooecia about 0·73 mm. long . . 7. ***U. perforata*** (Marsson)
  - (B) Oral spines branched.
    - (1) Distal oral spines bifurcate only; tertiary front wall with regular lacunae and a longitudinal, narrow, median depression proximal to each orifice 8. ***U. ocellata*** (Jullien)

(2) Distal oral spines each with 3–6 branches.
    (*a*) Orifice often as wide as high ; tertiary front wall thick and very irregular 9. ***U. meudonensis*** (Jullien)
    (*b*) Orifice wider than high ; transversely elongate-oval or oblong ; tertiary front wall thin, with regularly arranged lacunae and a longitudinal, narrow, median depression proximal to each secondary orifice 10. ***U. demorgani*** (Jullien)

(II) *Ubaghsia* with orifice nearly as large as, or larger than the frontal shield.
    (A) Orifice wider than high (hr = 0·20 to 0·25 mm., lr = 0·26 to 0·35 mm.) ; distal oral spines unbranched, often fused together at base . 11. ***U. reticulata*** (Ubaghs)
    (B) Orifice very large, of equal height and width (hr = 0·40 to 0·42 mm., lr = 0·40 to 0·45 mm.) . . . . 12. ***U. ornata*** (Goldfuss)

The following species, previously assigned to *Batrachopora* Lang, are of uncertain systematic position, and are therefore omitted from the above key. They are discussed on pp. 264–266.

(1) *Cellepora* (*Discopora*) *signata* von Hagenow.
(2) *Semiescharipora ovalis* d'Orbigny.
(3) *Reptescharipora convexa* d'Orbigny.
(4) *Batrachopora royanensis* Lang.
(5) *Batrachopora ultima* Voigt.
(6) *Escharoides peltata* Roemer.

1. *Ubaghsia ranunculus* (Lang)

(Pl. 20, fig. 2 ; Text-fig. 117)

1916*a* *Batrachopora ranunculus* Lang, pp. 110, 111.
1919*d* *Batrachopora ranunculus* Lang : Lang, pp. 203, 206, 208, 219, 221, text-figs. 33–35, 81.
1921 *Batrachopora ranunculus* Lang : Lang, pp. lvii, xci.
1922 *Batrachopora ranunculus* Lang : Lang, p. 362, pl. 8, fig. 1 ; text-fig. 114.
1935 *Batrachopora ranunculus* Lang : Bassler, p. 55.
1953 *Batrachopora ranunculus* Lang : Bassler, p. G192, text-fig. 144, 14.

HOLOTYPE. D.23388. Small zoarial fragment encrusting part of a *Cidaris* plate. Labelled "*Batrachopora ranunculus* Lang. Upper Senonian, Rügen"—probably boreal Maastrichtian. Agnes Laur Collection.

EMENDED DIAGNOSIS. *Ubaghsia* with zooecia 0·70 to 0·85 mm. long, 0·50 to 0·57 mm. wide ; secondary orifice wider than high (hr = 0·13 to 0·15 mm., lr = 0·20 mm.), much smaller than the frontal shield ; costae 8–10, each with 3 lateral costal fusions and pelmata ; avicularia of one kind only, a relatively large lateral oral pair fused above the secondary orifice to form a prominent proximal shield.

DESCRIPTION. *Zoarium* encrusting, unilaminar. *Zooecia* broadly oval. *Secondary orifice* wider than high, much smaller than the frontal shield, distally rounded, sides and proximal margin straight ; *distal oral shield* indistinct ; *proximal oral shield* prominent, formed by the paired, large, lateral oral avicularia which are raised on stout stalks and connected above the orifice by a bar of calcareous tissue. *Oral spines* probably 4, the distal pair are somewhat enlarged, but not lengthened, and are immersed in the secondary tissue accumulated on the distal rim of the orifice. The distal-lateral pair of oral spines are very probably fused on to the distal sides of the lateral oral avicularia and thus form part of the proximal oral shield. *Frontal shield* flat, except at the margins, broadly oval. *Costae* 8–10, fairly narrow, slightly tapering inwards. Near the outer end of each costa is a *pelma,* proximally to this the costa slopes down steeply to meet the gymnocyst; inwards from this pelma each costa is level and carries two smaller *pelmata.* Other pelmata or pelmatidia have not been seen. Somewhat arcuate *lateral costal fusions* arise at the levels of the pelmata producing fairly regular, somewhat elongate *intercostal spaces. Apertural bar* not markedly thickened, but generally wider than the largest costae, flat and straight, with at least two large primary pelmata near either end.



Fig. 117. *Ubaghsia ranunculus* (Lang), D.23388. Holotype. Adult zooecium with complete proximal oral shield.

Gymnocyst smooth, well-exposed proximally to the frontal shield, but often obscured laterally by *interzooecial secondary tissue* which is fairly abundant, in places filling the interzooecial valleys and containing irregularly oval *lacunae. Tertiary front wall* not seen, very probably never formed.

Avicularia apparently of one kind only, a large, prominent, lateral oral pair occurring one on either side of, and very close to, the proximal-lateral margins of the secondary orifice. They are raised on stout, much thickened stalks and are joined above the orifice by a bar of calcareous tissue forming a complete proximal oral shield. *Interzooecial avicularia* not seen, very probably absent. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·70 to 0·85 mm.
lz = 0·50 to 0·57 mm.
hr = 0·13 to 0·15 mm.
lr = 0·20 mm.

REMARKS. Only one specimen, D.23388, is available. Lang (1922 : 364) has stated that " occasional, sporadically-distributed aviculoecia [avicularia] appear in the interoecial [interzooecial] secondary tissue ", but they have not been seen. He probably interpreted oval lacunae as avicularia. Nor have " quaternary pelmatidia " been observed. Lang suggested that this species might have been derived from a form resembling *Ubaghsia perforata* (Marsson), but this is very uncertain.

STRATIGRAPHICAL DISTRIBUTION. Probably Maastrichtian.

SPECIMEN. D.23388. Holotype—see above.

2. *Ubaghsia aurita* (Lang)

(Pl. 20, fig. 3)

1916*a* *Batrachopora aurita* Lang, p. 111.
1919*d* *Batrachopora aurita* Lang : Lang, pp. 205, 206.
1922 *Batrachopora aurita* Lang : Lang, p. 376, text-fig. 119.

EMENDED DIAGNOSIS. *Ubaghsia* with zooecia more than twice as long as wide [according to Lang, Lz = about 1·10 mm., lz = about 0·49 mm.] ; secondary orifice wider than high or of equal height and width, much smaller than the frontal shield ; oral spines probably 4 ; frontal shield distinctly pointed proximally ; costae very numerous, 18–22 ; avicularia apparently of one kind only, a large lateral oral pair.

DESCRIPTION. [Based on a photograph of the holotype.] *Zoarium* unilaminar, encrusting [according to Lang]. *Zooecia* fairly closely placed, noticeably elongate. *Secondary orifice* wider than high, or of equal height and width, rounded distally with straight sides which tend to converge somewhat proximally, and a straight proximal margin. *Oral spines* probably 4. It is possible to distinguish, from the photograph, a distal pair of thickened, but not lengthened, oral spines ; distal-lateral oral spines, if present, are probably fused on to the distal sides of the lateral oral avicularia, forming, with these, the *proximal oral shield*. *Distal oral shield* indistinct. *Frontal shield* tapered proximally. *Costae* very numerous, 18–22, closely placed. Number of *pelmata* and *lateral costal fusions* not determinable. *Apertural bar* apparently fairly wide, straight, with at least two large pelmata, one at each end. *Gymnocyst* exposed proximally to the frontal shield, apparently not obscured by *interzooecial secondary tissue*.

Avicularia apparently of one kind only, a prominent, large, lateral oral pair. It cannot be determined, from the photograph, whether or not the lateral oral avicularia unite above the orifice to produce a complete proximal oral shield. *Ovicells* not seen.

MEASUREMENTS. [According to Lang, 1922 : 376.]
Zooecia Lz = About 1·10 mm.
lz = About 0·49 mm.

REMARKS. On the evidence obtainable from the photograph of the specimen (in Mons. F. Canu's Collection) Lang placed it as a species of *Batrachopora* [=*Ubaghsia*.]. This is apparently correct, judging from the general form of the secondary orifice and its proximal shield. The figure given by Lang (1922 : 375, text-fig. 119) is, however, largely hypothetical. The photograph is not sufficiently clear to show the precise shape of the oral avicularia, and certainly not clear enough to allow any accurate assessment of the number of pelmata or lateral costal fusions.

If the specimen is correctly placed as a species of *Ubaghsia* it is clearly distinguishable from other species of the genus by its unusually high number of costae, 18–22, and its single kind of avicularia.

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian.

SPECIMEN. Only a photograph of the holotype in the F. Canu Collection. Royan, south of Rochefort, Charente-Inférieure, France.

3. *Ubaghsia gasteri*[1] sp. nov.

(Pl. 20, figs. 4, 7 ; Text-figs. 118–120)

HOLOTYPE. D.40546. Unattached zoarial fragment. Senonian, lower part of zone of *G.* [*A.*] *quadrata*. Pit 21 of Gaster, northernmost of two pits on Patching Hill, Patching, Sussex.

PARATYPES. D.40547. Unattached zoarial fragment. Horizon, locality and collection as for the holotype.

D.40548. Unattached zoarial fragment. Senonian, lower part of zone of *G.* [*A.*] *quadrata*. Pit 15*a* of Gaster, small pit by roadside, south-east of Salvington Windmill, north-west of large pit (No. 15 of Gaster), and south of Pit 14 of Gaster, Salvington, Sussex.

DIAGNOSIS. *Ubaghsia* with closely placed zooecia (Lz = 0·80 to 0·93 mm., lz = 0·50 to 0·63 mm.) ; secondary orifice relatively large (hr = 0·19 to 0·25 mm., lr = 0·20 to 0·25 mm.), of variable shape, usually a little wider than high or of equal height and width, but occasionally a little higher than wide, smaller than the frontal shield ; distal oral shield present, proximal oral shield prominent ; oral spines 4 ; frontal shield flat, not descending steeply to meet the gymnocyst ; costae 9 or 10 (very occasionally more), each with two lateral costal fusions and pelmata ; interzooecial secondary tissue abundant ; avicularia of two kinds : (*a*) a large, prominent, proximal-lateral oral pair, (*b*) an occasional small, rounded, interzooecial form.

DESCRIPTION. *Zoarium* probably encrusting (but all specimens fragmentary and now unattached), unilaminar. *Zooecia* broadly oval, closely placed. *Secondary orifice* large, but much smaller than the frontal shield, usually a little wider than high, or of equal height and width, but occasionally slightly higher than wide. The shape of the orifice varies from almost square to rounded distally, with straight sides and a straight proximal margin. Proximal-lateral constrictions are visible in some orifices, particularly those with worn rims. *Distal oral shield* formed by the distal oral spines and secondary tissue ; *proximal oral shield* very prominent, formed mainly by the proximal-lateral oral avicularia with the distal-lateral pair of oral spines apparently fused on to the distal sides of the avicularia. *Oral spines* 4, enlarged, thickened, partly embedded in secondary tissue, hollow and cylindrical, arranged as a distal and a distal-lateral pair—the latter may be fused on to the distal sides of the lateral oral avicularia. *Frontal shield* flat : the edges may be partly

---

[1] After C. T. A. Gaster.



118



120



119

FIGS. 118–120. *Ubaghsia gasteri* sp. nov.

(118) D.40548. Paratype. Adult zooecium with worn, but very prominent, proximal-lateral oral avicularia fused to distal-lateral pair of oral spines. A small oval avicularium of type *b* occurs near the right edge of the frontal shield.

(119) D. 40546. Holotype. (*a*) Worn orifice of adult zooecium with proximal-lateral constrictions in the primary orifice. (*b*) Orifice of adult zooecium showing the square form produced by proximal oral shield. (*c, d, e*) Proximal, inner and outer views respectively of one side of a broken proximal oral shield showing the proximal-lateral oral avicularia fused with the distal-lateral oral spines.

(120) D.40546. Holotype. Orifice of adult zooecium with worn proximal-lateral oral avicularia and worn oral spines.

concealed by expansions of the interzooecial secondary tissue. *Costae* 9 or 10, very occasionally 13, tapering inwards, wide at their outer ends, where they descend slightly to meet the gymnocyst. Each costa carries a large, prominently rimmed *pelma* near its outer end and one, occasionally two, other pelmata nearer to the inner end. *Lateral costal fusions* arise at the levels of the pelmata, producing irregular intercostal spaces. *Apertural bar* straight, thickened, particularly at each end, and bearing two large pelmata and a pair of smaller pelmata between them nearer to the mid-line, which may be slightly furrowed. *Gymnocyst* almost entirely concealed by the close placing of the zooecia, which may result in the frontal shields of adjacent zooecia appearing contiguous, and by abundant *interzooecial secondary tissue* which generally fills the interzooecial valleys and in places expands on to the frontal shields. The interzooecial secondary tissue is marked by slight, irregular ridges surrounding shallow *lacunae*.

Avicularia of two kinds : (*a*) a very prominent proximal-lateral form, well raised on stout stalks with secondarily thickened walls, paired, one on each side of, and very close to, the orifice, (*b*) an occasional, interzooecial form which is very small, and distally pointed and directed. Avicularia of type (*a*) probably did not unite above the secondary orifice to form a complete proximal oral shield. Each of the lateral oral avicularia may be fused on its distal side to one of the distal-lateral pair of oral spines. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 0·80 to 0·93 mm.
lz = 0·50 to 0·63 mm.
hr = 0·19 to 0·25 mm.
lr = 0·20 to 0·25 mm.
Avicularia Not measurable.

REMARKS. *U. gasteri* is distinguished from other species of *Ubaghsia* by the close placing of its zooecia, the four thickened, often visibly separate, oral spines and the form of the frontal shield which does not descend steeply at the edges. The zoarial fragments were probably encrusting, but the specimens are broken and no longer attached. Two worn avicularia of type (*b*) are present in D.40548.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *G.* [*A.*] *quadrata*.

SPECIMENS. D.40546, holotype and D.40547–48, two paratypes—see above.

4. ***Ubaghsia langi***[1] sp. nov.

(Pl. 21, figs. 4, 5 ; Text-figs. 121–123)

HOLOTYPE. N.C.M.247.956 (4/1). Incomplete zoarium encrusting a test of *Conulus albogalerus*. Senonian, zone of M. *cor-anguinum*. Northfleet, Kent. A. W. Rowe Collection.

DIAGNOSIS. *Ubaghsia* with closely placed zooecia (Lz = 0·73 to 0·80 mm., lz = 0·50 to 0·60 mm.) ; secondary orifice fairly large, wider than high or of equal height and width, smaller than the frontal shield ; oral spines 4, much thickened and enlarged ; frontal shield flat, very steeply descending at the outer edges ; costae

---

[1] After W. D. Lang.



122



121



123

FIGS. 121–123. *Ubaghsia langi* sp. nov. N.C.M.247.956 (4/1).

(121) Holotype. Adult zooecium and part of a proximally adjacent zooecium: *a*, proximal-lateral oral avicularia forming proximal oral shield; *ia*, interzooecial avicularia; *os*, distal oral spines.

(122) Young zooecium; the rim of the secondary orifice is damaged.

(123) Three interzooecial avicularia.

7–10 ; lateral costal fusions 2 or 3 ; avicularia of two kinds : (*a*) a prominent, large, proximal-lateral oral pair, very close to the orifice, raised on stout stalks and united above the orifice to produce a strong proximal shield, (*b*) a prominent, frequent, distally pointed and directed, interzooecial form.

Description. *Zoarium* unilaminar, encrusting. *Adult zooecia* broadly oval, closely spaced. *Secondary orifice* wider than high, its form obscured by the proximal and distal oral shields. The rim of the secondary orifice is much thickened by secondary tissue which incorporates the much enlarged oral spines. The orifice is rounded distally with straight sides and a straight proximal margin. *Oral spines* in two pairs, much enlarged and thickened. *Distal oral shield* formed by the pair of distal oral spines and secondary tissue. *Proximal oral shield* very prominent, produced by the upgrowth and median fusion above the orifice of the very large lateral oral avicularia, often fused with the distal-lateral pair of oral spines. Occasionally the proximal shield may be enlarged by the additional incorporation of an enlarged interzooecial avicularium (*ia.* in Text-fig. 121). *Frontal shield* flat within the outermost lateral costal fusions, but outwards from them the costae descend very steeply to meet the obscured gymnocyst. *Costae* 7–10, stout, each with a very prominent *pelma* nearer the outer end, and, towards the inner end, one, or often two, smaller *pelmata*. *Lateral costal fusions* arise at the levels of the pelmata producing very irregular *intercostal spaces* in the wide median area. *Apertural bar* prominent, straight, with a pelma near each of the outer ends and another pelma near the mid-line. *Gymnocyst* obscured by fairly abundant *interzooecial secondary tissue* which almost fills the interzooecial valleys and contains irregular shallow *lacunae*. The secondary tissue forms a *tertiary front wall* which connects the proximal and distal oral shields of the zooecia, and often encroaches on to the edges of the frontal shields or stands up above their general level as a ridge.

Avicularia of two kinds : (*a*) a very large, thickened, heavily calcified lateral oral pair, raised on stout stalks, very close to, and one on either side of, the orifice. These unite above the orifice to produce the very prominent proximal oral shield ; (*b*) an interzooecial form, fairly prominent, frequent, unpaired, distally pointed and directed, lacking a transverse bar, but laterally constricted, producing a semicircular proximal portion and a triangular *rostrum*. Secondary thickening of avicularia of type (*b*) is variable. *Ovicells* not seen.

Young zooecia small. *Orifice* of equal height and width. *Oral spines* 4, thickened, the distal pair not so enlarged as the distal-lateral pair. *Frontal shield* very well arched but flat on top. *Costae* 9 or 10, not so closely placed as in the adult zooecia, each with a very large, prominent, *pelma*, and, inwards from this, one, occasionally two, *pelmatidia*. The pelmata form a conspicuous ring round the flat median area. Outwards from them the costae descend very steeply to meet the gymnocyst. *Lateral costal fusions* occur at the levels of the pelmata and pelmatidia, producing very irregular *intercostal spaces*. *Apertural bar* thickened, prominent, relatively straight, with two pelmata near the mid-line. *Gymnocyst* smooth ; widely exposed, especially proximally to the frontal shield. There is some *interzooecial secondary tissue* but this does not fill, or extend from, the interzooecial valleys as in the adult stages. *Avicularia* of two kinds : (*a*) a lateral oral pair, raised on stout stalks, and apparently not uniting to form a complete proximal oral shield ; (*b*) an occasional, sporadic, unpaired, rounded, small, interzooecial form.

MEASUREMENTS. *Adult zooecia* Lz = 0·73 to 0·80 mm.
lz = 0·50 to 0·60 mm.
hr = 0·20 mm.
lr = 0·20 to 0·23 mm.
Avicularia of type (*a*) Lz = 0·20 mm.
lz = 0·11 mm.
Young zooecia Lz = 0·53 to 0·62 mm.
lz = 0·30 to 0·32 mm.
hr = 0·12 mm.
lr = 0·12 mm.

REMARKS. *Ubaghsia langi* is distinguished from *U. gasteri* particularly by its more numerous, larger, distally pointed, interzooecial avicularia, and by its smaller size and generally fewer costae. The holotype encrusts a test of *Conulus albogalerus* from the Senonian, zone of *M. cor-anguinum.* On the base of the same echinoid is a well-preserved zoarial fragment of *Lagynopora horsleyensis* Lang, N.C.M.247.956 (4/2), a species recorded previously from the higher *M. cor-anguinum* Chalk. *Ubaghsia langi* is, as yet, the earliest known *Ubaghsia.* It does not have the features which Lang (1922 : 395) regarded as " primitive " for the genus, i.e., a large number of costae and a small size compared with the Maastrichtian species.

STRATIGRAPHICAL DISTRIBUTION. Senonian, higher part of zone of *M. cor-anguinum.*

MATERIAL. N.C.M.247.956 (4/1). Holotype—see above.

5. ***Ubaghsia crassa*** (Lang)

(Pl. 20, fig. 5 ; Pl. 21, fig. 7 ; Text-fig. 124)

1916*a* *Batrachopora crassa* Lang, pp. 110, 111.
1919*d* *Batrachopora crassa* Lang : Lang, pp. 206, 208, 217, 219, 221, text-fig. 83.
1922 *Batrachopora crassa* Lang : Lang, p. 365, pl. 8, fig. 3 ; text-fig. 116.
1925*b* *Batrachopora crassa* Lang : Voigt, p. 104.
1930 *Batrachopora crassa* Lang : Voigt, p. 551.

HOLOTYPE. D.16674. Poorly preserved zoarial fragment encrusting a cheilostome polyzoan ; only one worn zooecium is complete, in all others the frontal shield is destroyed. Labelled by Lang " *Batrachopora crassa.* Senonian, zone of *B. mucronata*, Rügen "—probably Maastrichtian. Agnes Laur Collection.

EMENDED DIAGNOSIS. *Ubaghsia* with fairly closely placed large zooecia (Lz = 1·00 to 1·12 mm., lz = 0·66 to 0·82 mm.) ; secondary orifice relatively large, wider than high (hr = 0·22 to 0·26 mm., lr = 0·28 to 0·30 mm.), smaller than the frontal shield ; oral spines at least 4 ; frontal shield flat, descending at the outer edges ; costae 7, each with 2 (very occasionally 3) lateral costal fusions and pelmata ; interzooecial secondary tissue fairly abundant ; avicularia of two kinds : (*a*) a large lateral oral pair, (*b*) an occasional, small, sporadic, interzooecial form.

Description. *Zoarium* encrusting, unilaminar. *Zooecia* broadly oval, less closely placed than in *U. gasteri*. *Secondary orifice* large, but smaller than the frontal shield, wider than high, shape variable from square or oblong to rounded distally with straight sides and a straight proximal margin. *Oral spines* probably at least 4, thickened but not lengthened, a distal pair visible in the rim of secondary tissue and a distal-lateral pair apparently fused on to the distal sides of the lateral oral avicularia. *Distal oral shield* indistinct. *Proximal oral shield* formed by the distal-lateral oral spines fused with the lateral oral avicularia which probably united above the orifice. *Frontal shield* occasionally somewhat arched, but commonly flat over most of its width, descending steeply at the outer edges. *Costae* of unusually constant number, 7, each with 2 (very occasionally 3) stout *lateral costal fusions* arising at the levels of large, prominent *pelmata*. A distinct ridge surrounds the pelmata and is apparently continuous on the upper surface of each costa between the pelmata. *Pelmatidia* may occur irregularly very close to the mid-line. *Apertural bar* broad, straight, with the outer ends bent distally, widening near the mid-line, with a straight median furrow and bearing two very large, elongate pelmata. *Gymnocyst* smooth, partly concealed by fairly abundant, irregularly



Fig. 124. *Ubaghsia crassa* (Lang), D. 16674. Holotype. Very worn adult zooecium—the only complete zooecium in the holotype zoarial fragment.

distributed *interzooecial secondary tissue,* which may in places expand on to the edges of the frontal shields. Irregular, shallow *lacunae* occur in the interzooecial tissue.

Avicularia of two kinds : (*a*) a very prominent, large lateral oral pair, well raised on stout stalks, with secondarily thickened walls, occurring very close to, and one on either side of, the orifice ; (*b*) an occasional, small, sporadic, interzooecial form. Avicularia of type (*a*) probably united above the orifice to produce a complete proximal oral shield. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 1·00 to 1·12 mm.
lz = 0·66 to 0·82 mm.
hr = 0·22 to 0·26 mm.
lr = 0·28 to 0·30 mm.
Avicularia Not measurable.

REMARKS. Lang (1922 : 368) based his description of *Batrachopora crassa* on the holotype, D.16674, the only specimen available. Unfortunately, it is badly damaged and worn, the details of the pelmata, oral spines and avicularia being particularly uncertain. The dimensions of its only complete zooecium are : Lz = 1·10 mm., lz = 0·75 mm., hr = 0·22 mm., lr = 0·30 mm. There are 7 costae, each with at least two pelmata and lateral costal fusions. These characters compare very closely with those found in N.C.M.76.937 (30), which is a better preserved zoarial fragment encrusting a piece of echinoid test.

He suggested that interzooecial secondary tissue is scanty and the gymnocyst widely exposed. This opinion was based on the very worn holotype where most of the frontal shields and much of the interzooecial tissue have been destroyed. N.C.M.76.937 (30), however, has fairly plentiful interzooecial secondary tissue. Avicularia are of two kinds, the oral pair being large and prominent, not, as Lang has stated, " apparently confined to a comparatively small, somewhat raised, apertural [oral] pair ".

N.C.M.76.937 (30) from the *B. mucronata* Chalk of Catton, Norfolk, extends the downward range of *Ubaghsia crassa* from the Rügen horizon.

STRATIGRAPHICAL DISTRIBUTION. Probably Maastrichtian, and Senonian, zone of *B. mucronata.*

SPECIMEN. D.16674. Holotype—see above.

OTHER MATERIAL. N.C.M.76.937 (30). Zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *B. mucronata.* Catton, near Norwich, Norfolk. R. M. Brydone Collection.

6. ***Ubaghsia coaxans*** (Lang)

(Pl. 20, fig. 6 ; Pl. 21, fig. 6 ; Text-fig. 125)

1916*a* *Batrachopora coaxans* Lang, p. 111.
1919*d* *Batrachopora coaxans* Lang : Lang, pp. 206, 208, 221, text-fig. 84.
1922 *Batrachopora coaxans* Lang : Lang, p. 377, pl. 8, fig. 5 ; text-fig. 120.
1930 *Batrachopora coaxans* Lang : Voigt, p. 551.

HOLOTYPE. D.14209. Fragment of zoarium, now unattached, with two complete zooecia. Labelled by Lang "*Batrachopora coaxans*. Upper Senonian, Rügen" —probably Maastrichtian. Agnes Laur Collection.

EMENDED DIAGNOSIS. *Ubaghsia* with zooecia 1·07 to 1·15 mm. long, 0·78 to 0·83 mm. wide, secondary orifice wider than high (hr = 0·25 to 0·26 mm., lr = 0·31 to 0·35 mm.); costae 11–14, each with 3 or 4 lateral costal fusions and pelmata; avicularia apparently of two kinds: (*a*) a large lateral oral pair, (*b*) a somewhat smaller, occasional, unpaired interzooecial form.



FIG. 125. *Ubaghsia coaxans* (Lang), D. 14209. Holotype. Adult zooecium with broken proximal oral shield.

DESCRIPTION. *Zoarium* unilaminar, now unattached but probably encrusting. *Zooecia* very broadly oval. *Secondary orifice* wider than high, much smaller than the frontal shield, rounded distally, sides and proximal margin straight. *Distal oral shield* indistinct. *Proximal oral shield* formed by the paired, large lateral oral avicularia which possibly united above the orifice. *Oral spines* probably at least 4; a distal pair of thickened and slightly enlarged, but not markedly lengthened, oral spines is almost completely immersed in the secondary tissue of the rim of the orifice, and a distal lateral pair probably fused on to the distal sides of the lateral oral avicularia, thus forming part of the proximal oral shield. *Frontal shield* wide, flat except at the margins. *Costae* 11–14, narrow. Near the outer end of each costa is a relatively small *pelma*, outwards from this each costa slopes down more steeply to meet the gymnocyst; inwards from this pelma each costa is level and carries another two, more commonly three, *smaller pelmata*. *Lateral costal fusions* 3 or 4 (corresponding in number to the pelmata), narrow, those at the level of the outermost pelmata arcuate. *Intercostal spaces* regularly arranged. *Apertural bar* wide, flat, straight, with 3 or 4 small pelmata. *Gymnocyst* smooth, well exposed, particularly proximally to the frontal shield, not obscured by *interzooecial secondary tissue* which is absent or only slightly developed.

Avicularia apparently of two kinds : (*a*) a large, prominent, lateral, oral pair with much thickened walls, (*b*) a somewhat smaller, occasional, unpaired, sporadic, rounded interzooecial form. Avicularia of type (*a*) were raised to some extent on thickened, stout stalks, and possibly fused above the orifice, but in the available material they are all worn down to the level of the secondary orifice. *Ovicells* not seen.

MEASUREMENTS. *Zooecia* Lz = 1·07 to 1·15 mm.
lz = 0·78 to 0·83 mm.
hr = 0·25 to 0·26 mm.
lr = 0·31 to 0·35 mm.

REMARKS. *Ubaghsia coaxans* is clearly distinguished by its numerous costae, pelmata and lateral costal fusions, and its dimensions. It is not possible to decide, on the available evidence, whether or not the oral avicularia united above the orifice.

STRATIGRAPHICAL DISTRIBUTION. Specimens labelled " Upper Senonian, Rügen ", but probably boreal Maastrichtian.

SPECIMENS. D.14209. Holotype—see above.

D.15041. Paratype. Zoarial fragment, now unattached. Labelled by Lang " *Batrachopora coaxans* ". Horizon, locality and collection as for the holotype.

7. ***Ubaghsia perforata*** (Marsson)

1887	*Cribrilina perforata* Marsson, pp. 98, 109, pl. 10, fig. 11.
1895	*Cribillina* [*sic*] *perforata* Marsson : Deecke, p. 80.
1900*b*	*Cribrilina perforata* Marsson : Canu, p. 445.
1916*a*	*Batrachopora perforata* (Marsson) Lang, pp. 110, 111.
1919*d*	*Batrachopora perforata* (Marsson) : Lang, pp. 206, 219, 221.
1921	*Batrachopora perforata* (Marsson) : Lang, p. xci.
1922	*Batrachopora perforata* (Marsson) : Lang, p. 362.
1925	*Cribrilina brachiata* Levinsen, p. 377, pl. 8, fig. 24.
1927	*Cribrilina brachiata* Levinsen : Voigt, p. 102.
1930	*non Stichocados ordinatus* Lang : Voigt, p. 510, pl. 30, fig. 14.

LECTOTYPE. (Chosen by Lang, 1922 : 362.) The specimen figured by Marsson, 1887, pl. 10, fig. 11. " Schreibkreide of Rügen " (? boreal Maastrichtian).

EMENDED DIAGNOSIS. *Ubaghsia* with encrusting or unattached zoarium ; secondary orifice wider than high, much smaller than the frontal shield ; costae 10–13, each with at least 3 lateral costal fusions ; avicularia apparently of two kinds : (*a*) a prominent lateral oral pair (*b*) ? occasional interzooecial forms ; [length of zooecia given by Lang (1922 : 362) as about 0·73 mm.].

REMARKS. Although Levinsen (1925 : 377, pl. 8, fig. 24) introduced *Cribrilina brachiata* as a new species, he himself suggested by its synonymy that it might be the same as *C. perforata* Marsson. I accept their synonymy, and although Levinsen's figure is obscure, the diagnosis of *Ubaghsia perforata* may be expanded as above to include Levinsen's species. I do not accept Voigt's (1930 : 510) placing of the species as synonymous with *Stichocados ordinatus* Lang. Levinsen's specimen was encrusting; Marsson's material was unattached. Lang (1922 : 362) regarded the species as " primitive in most of its features ".

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian, possibly extending downwards into the Senonian, upper part of the zone of *B. mucronata*.

SPECIMENS. None in the collection.

8. *Ubaghsia ocellata* (Jullien)

(Pl. 22, figs. 1, 2)

1886 *Steginopora ocellata* Jullien, p. 614, pl. 19, figs. 1–3.
1900*b* *Steginopora ocellata* Jullien : Canu, pp. 453, 454, text-fig. 66 [a reduced copy of Jullien, 1886, pl. 19, figs. 1–3].
1916*a* *Ubaghsia ocellata* (Jullien) Lang, p. 99.
1920 *Steginopora ocellata* Jullien : Canu & Bassler, p. 282, text-fig. 81S [a reduced copy of Jullien, 1886, pl. 19, fig. 1].
1921 *Ubaghsia ocellata* (Jullien) : Lang, p. xciii.
1922 *Ubaghsia ocellata* (Jullien) : Lang, p. 226.
1929 *Steginopora ocellata* Jullien : Canu & Bassler, p. 243.

LECTOTYPE. (Chosen by Lang, 1922 : 226.) The specimen figured by Jullien, 1886, pl. 19, fig. 1. " Etage sénonien. Port Brehay ", north-east of Gourbesville, north-west of Carentan, Manche, France. [Horizon probably middle or lower parts of zone of *B. mucronata*.]

EMENDED DIAGNOSIS. (Based on Jullien's figures and description of the species.) *Ubaghsia* with unilaminar, apparently unattached zoarium; distal oral spines bifurcate; orifice large, transversely elongate, almost oblong to rounded oblong, with straight proximal margin, much smaller than the convex frontal shield; number of costae, lateral costal fusions, and pelmata uncertain; avicularia of two kinds : (*a*) a large prominent, well-raised, lateral oral pair, (*b*) a single, occasionally paired, small, vicarious form, directed variously but always outwards from the zooecium it accompanies, distally pointed and divided by a straight transverse bar into a triangular rostrum and a smaller semicircular proximal portion; tertiary front wall complete, with a median depression containing two or three lacunae, and two or three more lacunae symmetrically placed near the margin : other lacunae, of about the size of the orifice, occur interzooecially.

REMARKS. Lang (1922 : 226) stated " having simpler apertural [oral] spines than *Ubaghsia demorgani* and *M.* [*sic*] *meudonensis*, *M.* [*sic*] *ocellata* is probably more primitive than these; its relationship to *M.* [*sic*] *reticulata* is doubtful ". However, *Ubaghsia ocellata* may be readily distinguished from *U. reticulata*, which has unbranched distal oral spines and a very irregular tertiary front wall. *U. meudonensis* and *U. demorgani* have more complex oral spines, each being branched from three to six times ; this feature particularly distinguishes these two species from *U. ocellata.* The tertiary front walls of the latter and of *U. demorgani* are, however, very similar.

STRATIGRAPHICAL DISTRIBUTION. Senonian, ? lower or middle part of zone of *B. mucronata.*

SPECIMENS. None in the collection.

9. *Ubaghsia meudonensis* (Jullien)

(Pl. 22, figs. 3, 4)

1886 *Steginopora meudonensis* Jullien, p. 614, pl. 17, figs. 1–5 ; pl. 18, figs. 1–3.
1916*a* *Ubaghsia meudonensis* (Jullien) Lang, pp. 99, 100.
1921 *Ubaghsia meudonensis* (Jullien) : Lang, p. xci.
1922 *Ubaghsia meudonensis* (Jullien) : Lang, p. 228.

LECTOTYPE. (Chosen by Lang, 1922 : 228). The specimen figured by Jullien, 1886, pl. 17, fig. 3. " Etage sénonien." Les Moulineaux, near Meudon, south-west of Paris. Zone of *B. mucronata.*

EMENDED DIAGNOSIS. (Based on Jullien's figures and description of the species.) *Ubaghsia* with unilaminar, encrusting zoarium ; distal oral spines with from two to six branches ; orifice large, as wide as high, rounded distally, sides and proximal margin straight, smaller than the frontal shield ; number of costae, lateral costal fusions and pelmata unknown ; avicularia of two kinds : (*a*) a large, prominent, well-raised, lateral oral pair, (*b*) a smaller, interzooecial form, usually paired but occasionally single or absent, often raised on a stout stalk, variously directed, with a small, lunate, proximal portion divided by a curved transverse bar from a triangular rostrum ; tertiary front wall apparently much thicker than in *U. ocellata* and *U. demorgani,* containing irregular large and small lacunae and well-marked secondary orifices.

REMARKS. *U. meudonensis* possesses a more robust tertiary front wall than *U. demorgani,* but differences in this character alone may be insufficient to keep the two species distinct. However, until specimens are available for study these two species are maintained as separate.

See also remarks on *U. ocellata,* (p. 255).

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *B. mucronata.*

SPECIMENS. None in the collection.

10. *Ubaghsia demorgani* (Jullien)

(Pl. 22, figs. 5, 6)

1886 *Steginopora demorgani* Jullien, p. 615, pl. 19, figs. 4, 5 ; pl. 20, fig. 1.
1916*a* *Ubaghsia demorgani* (Jullien) Lang, p. 99, 100.
1921 *Ubaghsia demorgani* (Jullien) : Lang, p. xci.
1922 *Ubaghsia demorgani* (Jullien) : Lang, p. 227.

LECTOTYPE. (Chosen by Lang, 1922 : 227.) The specimen figured by Jullien, 1886, pl. 19, fig. 4. " Etage sénonien, craie de Meudon (Les Moulineaux)." Meudon, south-west of Paris. Zone of *B. mucronata.*

Emended diagnosis. (Based on Jullien's figures and description of the species.) *Ubaghsia* with bilaminar zoarium ; distal oral spines with three or four branches ; orifice large, transversely elongate-oval or almost oblong, or rounded distally with straight sides and a straight proximal margin, much smaller than the convex frontal shield ; number of costae, lateral costal fusions, and pelmata not known ; avicularia of two kinds : (*a*) a large, prominent, well-raised, lateral oral pair, (*b*) a small, vicarious form, commonly paired but occasionally single or absent, directed variously, but always outwards from the zooecium it accompanies, distally pointed and divided by a straight transverse bar into a triangular rostrum and a smaller, semicircular proximal portion ; tertiary front wall complete, with the orifice markedly transversely elongate, and the lacunae and median depression similar to those of *U. ocellata* in number and arrangement, though the interzooecial lacunae are apparently much smaller than those of that species.

Remarks. See remarks on *U. ocellata* (p. 255) and *U. meudonensis*, p. 256).

Stratigraphical distribution. Senonian, zone of *B. mucronata.*

Specimens. None in the collection.

11. *Ubaghsia reticulata* (Ubaghs)

(Pl. 21, figs. 1–3 ; Pl. 22, figs. 7, 8 ; Text-figs. 126–128)

1865	*Steginopora reticulata* Ubaghs, p. 55, pl. 2*a*, fig. 7*a–d*.
1886	*Ubaghsia reticulata* (Ubaghs) Jullien, pp. 610, 612, 616, pl. 18, figs. 4, 5.
1916*a*	*Ubaghsia reticulata* (Ubaghs) : Lang, p. 99.
1916*a*	*Batrachopora hyla* Lang, pp. 111, 112.
1919*d*	*Batrachopora hyla* Lang : Lang, pp. 205, 208, 219, 221, text-fig. 82.
1921	*Ubaghsia reticulata* (Ubaghs) : Lang, pp. lxvi, xc.
1921	*Batrachopora hyla* Lang : Lang, p. xc.
1922	*Ubaghsia reticulata* (Ubaghs) : Lang, p. 227.
1922	*Batrachopora hyla* Lang : Lang, p. 364, pl. 8, fig. 2 ; text-fig. 115.
1922	*Batrachopora ornata* (Goldfuss) : Lang, p. 370 [*partim*—pl. 8, fig. 4, specimen D.1391 only].
1929	*Steginopora reticulata* Ubaghs : Canu & Bassler, p. 244.
1930	*Ubaghsia reticulata* (Ubaghs) : Voigt, p. 513 [*partim*], pl. 31, fig. 19 only, *non* fig. 20.
1930	*Batrachopora ornata* (Goldfuss) : Voigt, p. 515, pl. 31, fig. 18.
1930	*Batrachopora hyla* Lang : Voigt, p. 515.
1935	*Steginopora reticulata* Ubaghs : Bassler, p. 222.
1953	*Steginopora reticulata* Ubaghs : Bassler, p. G193.
1953	*Ubaghsia reticulata* (Ubaghs) : Bassler, p. G193, text-fig. 144, 30.

Lectotype. (Chosen by Lang, 1922 : 227.) The specimen figured by Ubaghs, 1865, pl. 2*a*, fig. 7*c*. Maastrichtian. Ober Bryozoenschichte (horizon *f* of Ubaghs). Valkenburg (Fauquement), east of Maastricht, or Geulem, north-east of Maastricht, Holland.

Emended diagnosis. *Ubaghsia* with unbranched distal oral spines ; zooecia 0·80 to 0·85 mm. long 0·60 to 0·72 mm. wide ; secondary orifice wider than high (hr = 0·20 to 0·25 mm., lr = 0·26 to 0·35 mm.) ; costae 4–7 (often 5 or 6) each with 3 (occasionally 2) lateral costal fusions and pelmata ; avicularia of two kinds : (*a*) a large lateral oral pair, (*b*) small, rounded, occasionally paired, interzooecial forms.



126



127



128

DESCRIPTION. *Zoarium* unilaminar, probably encrusting. *Zooecia* broadly oval, smaller than in *U. ornata*. *Secondary orifice* very large, wider than high, almost as large as the frontal shield, rounded distally and laterally with a straight proximal margin. The orifice is surrounded by complete distal and proximal oral shields. The *distal oral shield* is formed by the amalgamation of the pair of distal oral spines which are enlarged and joined by secondary tissue. The *proximal oral shield* is composed of the very large, paired oral avicularia, which fuse above the apertural bar and orifice, and which incorporate on their distal sides the distal-lateral oral spines. *Oral spines* originally 4, rapidly enlarged but unbranched, united by secondary tissue to form parts of the distal and proximal oral shields. *Frontal shield* generally flat, broadly oval. *Costae* 4–7 (often 5–6), broad, irregular, tapering inwards ; near the outer end of each costa is a large *pelma*, outwards from which the ends of the costae bend down abruptly to meet the gymnocyst, while inwards there is one other *pelma* of large size and commonly, nearer the mid-line a third *pelma*. No other definite pelmata or pelmatidia have been seen. *Lateral costal fusions* occur opposite the pelmata, so that there are normally at least three pairs on each costa producing irregular, somewhat lacunate, *intercostal spaces*. *Apertural bar* not prominently thickened, straight, with at least two, often three, pairs of pelmata. *Gymnocyst* smooth, particularly exposed proximally to some frontal shields.

Avicularia of two kinds : (*a*) a very large lateral oral pair raised on much thickened stalks, one on either side, of, and close to, the lateral margins of the orifice, (*b*) a small, rounded, interzooecial form which is apparently sometimes paired, one near each end of the apertural bar and a little proximal to it. Avicularia of type (*a*) make up an essential part of the proximal oral shield ; the unmodified form of these avicularia has not been seen.

Tertiary front wall ultimately developed. The distal and proximal oral shields, above the secondary orifices, increase in size by the virtual sealing of the space directly above the orifice by a secondary plug of tissue. In addition, the huge, complete oral shields of adjacent zooecia join up by further accumulation of secondary calcareous tissue to form a very irregular cover pierced by deep *lacunae* and raised above the levels of the underlying frontal shields on the very stout stalks of the oral avicularia and on the ramifications of the distal oral shields. *Ovicells* not seen.

FIGS. 126–128. *Ubaghsia reticulata* (Ubaghs), D.1391. Labelled by Lang " *Batrachopora ornata* (Goldfuss) ".

(126) Adult zooecium and part of proximally adjacent zooecium. The tertiary front wall has been destroyed by wear.

(127) Adult zooecium with complete proximal oral shield supporting a fragment of tertiary front wall.

(128) Proximal oral shields. (*a*) Front view of orifice with broken proximal oral shield ; *fd*, fused distal oral spines ; *ia*, interzooecial avicularia ; *la*, lateral oral avicularia. (*b*, *c*) Side and top views respectively of orifice of adult zooecium with complete proximal oral shield ; *ab*, apertural bar ; *cp*, central plug of secondary tissue ; *fd*, fused distal oral spines ; *la*, lateral oral avicularia.

MEASUREMENTS. *Zooecia* Lz = 0·80 to 0·85 mm.
lz = 0·60 to 0·72 mm.
hr = 0·20 to 0·25 mm.
lr = 0·26 to 0·35 mm.
Avicularia Not measurable.

REMARKS. Voigt (1930 : 513) recognized that *Ubaghsia reticulata* (Ubaghs) and *Batrachopora hyla* Lang are synonymous. He stated " *Batrachopora hyla* Lang is merely a *Ubaghsia reticulata* in which the lamina peristomica [tertiary front wall] is missing " [translation]. Discussing other species of *Batrachopora*, he also remarked (1930 : 515) that " this [i.e., loss of the tertiary front wall in fossilization] is also true for other species, for example, in the case of the closely related *Batrachopora ornata* (Goldfuss), or even for the whole genus. But until there is more definite proof of this I retain the other species in *Batrachopora* " [translation].

Voigt's suggestions are here regarded as valid and the two genera are merged, for the frontal shields of *Batrachopora* and *Ubaghsia* are completely similar in structure, while the tertiary front wall was obviously loosely attached and easily lost in fossilization. Comparison of Jullien's figures (1886, pl. 18, figs. 4, 5) of *Ubaghsia reticulata* with D.1391, in which part of the tertiary front wall is preserved, make the synonymy particularly convincing.

See also remarks under *Ubaghsia ornata* (Goldfuss), (p. 263).

STRATIGRAPHICAL DISTRIBUTION. Maastrichtian.

SPECIMENS. D.11852. Holotype of *Batrachopora hyla* Lang. Small fragment of zoarium, now unattached. Maastrichtian, Maastricht, Holland. van Breda Collection.

D.1391. Zoarial fragment labelled by Lang " *Batrachopora ornata* (Goldfuss) ". Part of the tertiary front wall is preserved. Maastrichtian, Maastricht, Holland. G. R. Vine Collection.

D.40674–78. Five zoarial fragments labelled by Lang " *Batrachopora hyla* ". Horizon, locality and collection as for D.11852.

12. ***Ubaghsia ornata*** (Goldfuss)

(Pl. 23, figs. 1–4 ; Text-fig. 129)

1826 *Cellepora ornata* Goldfuss, pp. 26, 248, pl. 9, fig. 1*a*, *b*.
1828 *Cellepora ornata* Goldfuss : Morren, p. 34.
1832 *Cellepora ornata* Goldfuss : Dumont, p. 360.
1834 *Cellepora ornata* Goldfuss : von Klöden, pp. 266, 341, pl. 5, fig. 3.
1836 *Discopora ornata* (Goldfuss) Edwards, p. 253.
1846 *Cellepora ornata* Goldfuss : Geinitz, p. 612.
1848 *Cellepora ornata* Goldfuss : Bronn, p. 255.
1848 *Discopora ornata* (Goldfuss) : Bronn, p. 432.
1849 *Discopora ornata* (Goldfuss) : Bronn, p. 130.
1849 *Cellepora ornata* Goldfuss : Geinitz, p. 252.
1850 *Escharina ornata* (Goldfuss) d'Orbigny, p. 262.
1851 *Cellepora* (*Dermatopora*) *ornata* Goldfuss : von Hagenow, p. 98, pl. 10, fig. 16.
1851 *Cellepora* (*Dermatopora*) *ornata* Goldfuss : Bronn and Römer, p. 103 ; 1852, pl. 29^2, fig. 11 [copy of von Hagenow, 1851, pl. 10, fig. 16].

1853 *Semiescharipora ornata* (Goldfuss) d'Orbigny, p. 480.
1857 *Semiescharipora ornata* (Goldfuss) : Pictet, p. 112.
1862 *Cellepora ornata* Goldfuss : Goldfuss, p. 24, pl. 9, fig. 1*a*, *b*.
1879 *Cellepora ornata* Goldfuss : Quenstedt, p. 313, pl. 154, fig. 42.
1879 *Semiescharipora ornata* (Goldfuss) : Ubaghs, p. 217.
1881 *Semiescharipora ornata* (Goldfuss) : Mourlon, p. 116.
1885 *Cellepora ornata* Goldfuss : Vine, p. 161.
1885 *Cellepora* (*Dermatopora*) *ornata* Goldfuss : Vine, p. 164.
1889 *Semiescharipora ornata* (Goldfuss) : Ubaghs, p. 52.
1900*b* *Cribrilina* (*Costula*) *ornata* (Goldfuss) Canu, p. 450.
1916*a* *Batrachopora ornata* (Goldfuss) Lang, pp. 111, 112.
1916*a* *Cellepora ornata* Goldfuss : Lang, p. 169.
1919*d* *Batrachopora ornata* (Goldfuss) : Lang, pp. 205, 206, 217, 219, 221.
1921 *Batrachopora ornata* (Goldfuss) : Lang, p. xc.
1922 *Batrachopora ornata* (Goldfuss) : Lang, p. 370 [*partim*], text-fig. 117 [*non* pl. 8, fig. 4 = *Ubaghsia reticulata* (Ubaghs)].
1930 *non Batrachopora ornata* (Goldfuss) : Voigt, pp. 515, 543, 561, pl. 31, fig. 18.
1930 *Ubaghsia reticulata* (Ubaghs) : Voigt, p. 513 [*partim*], pl. 31, fig. 20, only—*non* fig. 19.

LECTOTYPE. (Chosen by Lang, 1922 : 372). The specimen figured by Goldfuss, 1826, pl. 9, fig. 1*b*. Maastrichtian, St. Petersburg, near Maastricht, Holland.

EMENDED DIAGNOSIS. *Ubaghsia* with unbranched distal oral spines ; zooecia large (Lz = 1·00 to 1·25 mm., lz = 0·70 to 0·90 mm.) ; secondary orifice usually of equal height and width, as large as or larger than, the frontal shield ; costae massive, 3–7 (commonly 5), each with two lateral costal fusions and two very large pelmata ; avicularia of two kinds : (*a*) a very large lateral oral pair, (*b*) commonly paired, small, distally pointed interzooecial avicularia directed obliquely outwards from the zooecium they accompany.

DESCRIPTION. *Zoarium* unilaminar, probably encrusting. *Zooecia* broadly oval, very large. *Secondary orifice* very large, generally of equal height and width, as large as, or larger than, the frontal shield, rounded distally and laterally with a relatively straight proximal margin, or circular. The orifice is surrounded by complete distal and proximal oral shields. *Distal oral shield* probably formed by the amalgamation of the distal and distal-lateral pairs of oral spines. All the available specimens are generally worn, with only occasional zooecia with undamaged distal and proximal oral shields. Apparently the distal-lateral oral spines do not fuse on to the distal sides of the large oral avicularia as in *Ubaghsia reticulata*. *Proximal oral shield* formed by the very large, paired, lateral oral avicularia, which are raised on very thick stalks. The oral avicularia are distally pointed, proximally rounded and directed distally and obliquely inwards. They very probably fuse above the secondary orifice by the addition, mainly on their upper, inner sides, of secondary calcareous tissue. *Oral spines* 4, enlarged, unbranched, and immersed in much secondary tissue forming the distal oral shield. *Frontal shield* generally flat, transversely oval or longitudinally oval. Costae 3–7 (commonly 5), often very coarse and wide : the occurrence of only 3 very large costae in an adult zooecium of normal size is exceptional (D.8529). Each costa bears two very large *pelmata* at the levels of the *lateral costal fusions*, which are of variable thickness producing irregular *intercostal spaces*. Outwards from the very large proximal pelma each costa bends downwards abruptly to meet the *gymnocyst* which is apparently smooth but seldom visible. The *apertural bar* is about as wide as the costae, straight or slightly concave, and carrying two large pelmata which may be confluent in one large depression.

Avicularia of two kinds: (*a*) a very large lateral oral pair, one on either side of, and close to, the lateral margins of the orifice, (*b*) a small, commonly paired interzooecial form. Avicularia of type (*a*) are proximally rounded and distally pointed, directed distally and obliquely inwards, raised on much thickened stalks, and probably connected by a bar-like accumulation of secondary tissue between them. Avicularia of type (*b*) are much smaller, commonly paired, one near each end of the apertural bar and a little proximally to it; they are pointed, rounded proximally and directed distally and obliquely outwards from the zooecium they accompany. These small interzooecial avicularia are often worn and partly obscured by secondary tissue expanding from the proximal oral shield. *Tertiary front wall* probably present and formed as in *Ubaghsia reticulata*, but eroded in the specimens available. *Ovicells* not seen.



Fig. 129. *Ubaghsia ornata* (Goldfuss). D.19356. Labelled by Lang "*Batrachopora ornata* (Goldfuss)". Adult zooecium with typically very large orifice and two pointed interzooecial avicularia. The proximal oral shield and tertiary front wall destroyed by wear.

Measurements. *Zooecia* Lz = 1·00 to 1·25 mm.
lz = 0·70 to 0·90 mm.
hr = 0·40 to 0·42 mm.
lr = 0·40 to 0·45 mm.
Lateral oral avicularia Lz = 0·16 mm.
lz = 0·10 mm.
Interzooecial avicularia Not measurable.

Remarks. Lang (1922 : 372) emphasized the importance of the distal and proximal oral shields as distinguishing features of the species, and compared the distal shields with those of *Ichnopora denticulata*. He figured (1922, pl. 8, fig. 4), as representative of *Ubaghsia ornata*, four zooecia of D.1391, which is a fairly well-preserved zoarial fragment retaining part of its tertiary front wall. This specimen clearly belongs to *Ubaghsia reticulata* (Ubaghs), for its measurements and other characters agree with those of that species and differ consistently from those of the other specimens (D.8529, D.19356–57) which Lang also placed in *ornata*. I agree that the last three specimens should be retained in *Ubaghsia ornata*.

Ubaghsia ornata (Goldfuss) is distinguished from *U. reticulata* (Ubaghs) by its fewer costae, its pelmata (which are equal in number to the lateral costal fusions), its larger zooecia, and its larger and differently shaped orifice. In addition, the lateral oral and interzooecial avicularia of *U. ornata* differ from those of *U. reticulata*. The species can be distinguished without placing too much reliance on the characters of the distal and proximal oral shields ; not only are these structures easily eroded, but they may also vary considerably in form and size, depending on the amount of secondary tissue accumulated round them. The tertiary front wall is extremely variable and irregular, and is seldom preserved at all in the material available ; it is of diagnostic value generically but does not aid in separation of the species.

D.28526 is doubtfully assigned to *U. ornata*. The measurements and other characters of three of the complete adult zooecia in the specimen (e.g., Lz, lz and the number of lateral costal fusions) are more in accord with those of *U. ornata* than with those of *U. reticulata*. But the dimensions of the orifice (hr = 0·30 mm., lr = 0·35 mm.) are generally smaller than those of *U. ornata* and it is somewhat wider than high.

Ubaghsia ornata (Goldfuss) and *U. reticulata* (Ubaghs) are closely related but distinguishable species. As yet they have been recorded only from the Maastrichtian of Maastricht.

Lang (1922 : 365) has suggested that they were derived respectively from *U. crassa* and *U. ranunculus*, two species described by him from the Rügen Chalk. Apart from the possibility that the Rügen horizon of these specimens is equivalent to the littoral Maastrichtian of the type area, the geographical relationship of the two areas makes such a relationship very unlikely.

If Lang's suggested lineages for *Ubaghsia ornata* and *U. reticulata* are not accepted, it does not mean that the " alternative hypothesis " mentioned by him (1922 : 366), that *U. ornata* is derived from *U. reticulata*, must be adopted. There is no evidence to decide whether *U. ornata* developed from *U. reticulata* or vice versa.

Stratigraphical distribution. Maastrichtian.

Specimens. D.8529. A large zoarial fragment, now unattached, labelled by Lang " *Batrachopora ornata* (Goldfuss) ". Maastrichtian. Fauquement, east of Maastricht, Holland. W. Gamble Collection.

D.19356–57. Two zoarial fragments, now unattached, labelled by Lang " *Batrachopora ornata* (Goldfuss) ". Maastrichtian. Maastricht, Holland. G. R. Vine Collection.

?D.28526. A fragmentary zoarium, labelled by Lang "*Batrachopora ornata* (Goldfuss)". Horizon, locality and collection unknown, but very probably Maastrichtian of Maastricht, Holland. [The specimen is here only doubtfully assigned to *Ubaghsia ornata* (Goldfuss).]

SPECIES OF UNCERTAIN SYSTEMATIC POSITION PREVIOUSLY ASSIGNED TO *BATRACHOPORA*

(1) ***Cellepora (Discopora) signata*** von Hagenow

1851 *Cellepora* (*Discopora*) *signata* von Hagenow, p. 96, pl. 10, fig. 17*a*, *b*.

REMARKS. Although Lang (1922 : 361) chose the specimen figured by von Hagenow as lectotype, he also remarked "from the evidence available, it is impossible to diagnose this species or more than tentatively to refer it to *Batrachopora*". There is no evidence to suggest it might be a *Batrachopora* ; the genus and species are indeterminate.

(2) ***Semiescharipora ovalis*** d'Orbigny

1852 *Semiescharipora ovalis* d'Orbigny, pl. 719, figs. 13–16.
1853 *Semiescharipora ovalis* d'Orbigny : d'Orbigny, p. 488.
1854 *Semiescharipora ovalis* d'Orbigny : d'Orbigny, p. 1098.

REMARKS. Lang (1916*a* : 110, 111) placed *Semiescharipora ovalis* d'Orbigny in *Batrachopora* and later (1922 : 367) selected the specimen figured by d'Orbigny (1852, pl. 719, fig. 14) as lectotype. Again there is no definite evidence to suggest that the material figured by d'Orbigny or described by him (1853 : 488) belongs to *Batrachopora*. Canu (1900*b* : 457) has listed the species as worn and doubtful. All that may be deduced with certainty from the figure and description is that *Semiescharipora ovalis* d'Orbigny is a cribrimorph polyzoan from the Maastrichtian of Royan. It has a well-developed frontal shield with fairly numerous costae and lateral costal fusions, and paired, large, lateral oral avicularia.

(3) ***Reptescharipora convexa*** d'Orbigny

1852 *Reptescharipora convexa* d'Orbigny, pl. 720, figs. 1–3.
1853 *Reptescharipora convexa* d'Orbigny : d'Orbigny, p. 492.
1854 *Reptescharipora convexa* d'Orbigny, : d'Orbigny, p. 1098.

REMARKS. Lang (1916*a* : 111, 112) regarded *Reptescharipora convexa* d'Orbigny as a species of *Batrachopora* and later (1922 : 368) selected the specimen figured by d'Orbigny (1852, pl. 720, fig. 2) as lectotype.

There is, if anything, even less evidence for regarding *Reptescharipora convexa* as a *Batrachopora* than in the case of *Semiescharipora ovalis* d'Orbigny. It is a cribrimorph polyzoan, from the Senonian of Meudon, with apparently about 10 costae and about 5 lateral coatal fusions. The paired lateral oral avicularia are not large. Canu (1900*b* : 449) referred the species to *Cribrilina*. He gave the dimensions of the species as Lz = 1·00 to 1·14 mm., lz = 0·57 to 0·71 mm., hr = 0·17 to 0·22 mm., lr = 0·28 mm., and mentioned the occasional presence of two marginal [oral] spines.

(4) ***Batrachopora royanensis*** Lang

1916*a* *Batrachopora royanensis* Lang, pp. 110, 111.
1919*d* *Batrachopora royanensis* Lang : Lang, pp. 205, 206, 221.
1922 *Batrachopora royanensis* Lang : Lang p. 374, text-fig. 118.

REMARKS. Lang (1916*a*) based his diagnosis of this species on a photograph of the holotype, which is in the F. Canu Collection. It can be seen from the photograph that the orifices of the zooecia are much smaller than the frontal shields, wider than high, rounded distally with straight sides, which begin to converge proximally, and with a straight proximal margin. There are apparently no lateral oral avicularia. Apparently 12–14 costae are present, each with at least two lateral costal fusions and pelmata and pelmatidia. The number of oral spines is indeterminate. Interzooecial avicularia of two kinds appear to be present : (*a*) a small, rounded, paired form just distal to each orifice, (*b*) a larger, laterally constricted, occasional, sporadic form of uncertain shape and orientation. Ovicells cannot be seen. There is no indication that the zoarium is necessarily " erect [and] bilaminar " as Lang (1922 : 374) has stated.

There is not sufficient evidence to place the specimen with certainty in any cribrimorph genus. It is clear, however, that the specimen lacks the large, well-defined, lateral oral avicularia characteristic of *Batrachopora*, and that its frontal shield is different from those of species of *Ubaghsia* (which here includes *Batrachopora*).

(5) ***Batrachopora ultima*** Voigt

1925*b* *Batrachopora ultima* Voigt, pp. 99, 103, pl. 3, figs. 7, 8 ; text-fig. 4.
1930 *Batrachopora ultima* Voigt : Voigt, pp. 515, 561, pl. 31, fig. 16 (copy of part of pl. 3, fig 8, 1925*b*).
1939 *Batrachopora ultima* Voigt : Voigt, p. 93, text-fig. 4*c*.

REMARKS. Voigt (1925*b* : 103) described the species as follows : " zoarium unilaminar, erect. Zooecia arranged in regular quincuncial lines, elongate-oval, sometimes twice as long as broad but usually somewhat shorter. The ' anterior wall ' is formed only from the frontal shield, which, on the average, consists of from 7–10 exceptionally broad costae which fuse laterally and at their inner ends along the mid-line. Thus the frontal shield is pierced generally by 4, or more rarely 5, regular, parallel, transverse, rows of pores [intercostal spaces], usually 3 or 4 to a row. The broad ' prolongations ' [outermost ends] of the costae alternate at the margin [of the frontal shield] with the narrower inter[-costal] spaces. Under the microscope, delicate ' vertically crossing ' lines may be seen on the costae. *Pelmata were presumed in view of the ' sporadic punctures '* [?pores or intercostal spaces] *rather than observed ; the figure* [Voigt, 1925*b*, text-fig. 4] *is to be considered in this light.*[1] Orifice semicircular . . . surrounded usually by 5 [oral] spines. In all probability there is a small, elongate avicularium on either side of the orifice. I consider this to be the case because of the small protuberances on either side of the orifice, each of which seems to have possessed a pore. Several other species of *Batrachopora* support this, as they always have avicularia on either side of the orifice. *But I have*

---

[1] My italics.

not observed the avicularia for certain because of the condition of preservation of the specimen.[1]

" *Batrachopora ultima* is the youngest *Batrachopora*. *B. crassa* Lang is similar (also with about 7 costae) but it is distinctly larger (1·25 mm.). *B. perforata* (Marsson), from Rügen, which is distinguished by two large lateral avicularia, is more distantly related " [translation].

It is apparent that Voigt described *B. ultima* without fully determining the character of the frontal shield, particularly the presence or absence of definite pelmata. He also seemed uncertain of the nature of the oral avicularia, or even whether they were present or not.

I do not accept *B. ultima* Voigt as a definite species of *Ubaghsia*, because of the uncertainty of the true character of the type material. There is one specimen, D.31990, in the British Museum (Natural History), presented by Prof. E. Voigt, and labelled by him " *Batrachopora ultima* Voigt, Daniendrift, Blaue Berge, near Rosslau, Anhalt ". This specimen is a cribrimorph polyzoan with frequent, prominent, hyperstomial ovicells, and is quite unlike any species of *Batrachopora* [=*Ubaghsia*]. Lang's label with the specimen shows that he doubtfully retained it in *Batrachopora*.

(6) ***Escharoides peltata*** Roemer

1840 *Escharoides peltata* Roemer, p. 14, pl. 5, fig. 7.

REMARKS. Lang (1916 : 410 ; 1921 : 13) has already listed this as a species of doubtful systematic position. However, Voigt (1949 : 41, pl. 10, figs. 4, 5) recorded *Batrachopora* [*Escharoides*] *peltata* (Roemer) from the Senonian, zone of *G.* [*A.*] *quadrata* of Oberg and Misburg, Hanover, Germany. I do not agree with Voigt's placing of his specimens as a species of *Batrachopora*. Roemer's figure is difficult to interpret, and Voigt's very clear figures (1949, pl. 10, figs. 4, 5) show a species which may belong to *Escharoides peltata* Roemer, but definitely not to *Batrachopora* Lang.

Genus ***ANDRIOPORA*** Lang

1921 *Andriopora* Lang, p. 89—for earlier references.

The new species described below are assigned to *Andriopora* which has not been completely revised. The described material has been compared with all available specimens of *Andriopora* in the collections of the British Museum (Natural History).

1. ***Andriopora minor***[2] sp. nov.

(Pl. 23, fig. 6 ; Text-fig. 130)

HOLOTYPE. D.39417. Zoarial fragment encrusting the dorsal surface of a fragment of a unilaminar zoarium of *Pelmatopora* cf. *palmata* Lang. Senonian, zone of *G.* [*A.*] *quadrata*. Pit 17 of Gaster, large pit, in two sections, in Cote Bottom, Durrington, Sussex.

---

[1] My italics.

[2] With reference to the small size of the zooecia.

DIAGNOSIS. *Andriopora* with small, elliptical, non-caudate zooecia ; orifice wider than high ; oral spine-bases 6 ; costae 15–20, each with about 4 lateral costal fusions ; apertural bar unmodified ; gymnocyst narrowly exposed round the frontal shield ; avicularia generally paired.

DESCRIPTION. *Zoarium* encrusting, uniserial, tending to appear multi-serial through an irregular decrease in the angle of branching and through the tendency for different series of zooecia to overlap. *Zooecia* small, elliptical, non-caudate. *Orifice* wider than high, rounded distally and laterally or tending to be straight-sided, proximal margin straight. *Oral spine-bases* 6, very minute, hardly visible. *Frontal shield* well arched, slightly flattened on top. *Costae* 15–20, very narrow, with a variable number, often 4, very fine *lateral costal fusions*. There are no median pores or slits on the costae, nor do the costae tend to be up-turned at their inner ends. *Median area of fusion* narrow. *Apertural bar* unmodified, but tending to be somewhat wider than the costae. *Gymnocyst* narrowly exposed laterally, and more widely exposed proximally to the frontal shield and round the avicularia.



FIG. 130. *Andriopora minor* sp. nov. D.39417. Holotype. Five zooecia ; one with single avicularium and one with paired avicularia. One zooecium with slightly damaged ovicell.

Avicularia large, prominent, paired, one just beyond each distal lateral margin of the orifice, or occurring singly on one side or the other, directed distally away from the mid-line of the orifice they accompany. The avicularian aperture is elongate-oval, or tending to be pointed distally, and is divided by a narrow *transverse bar* into a more-or-less triangular *rostrum*, and a wider, semicircular proximal portion.

Ovicells hyperstomial, very prominent, large, globular, smooth, somewhat compressed laterally and standing up above the zooecium on which they are developed.

MEASUREMENTS. *Zooecia* Lz = 0·27 to 0·33 mm.
lz = 0·15 to 0·18 mm.
hr = 0·035 mm.
lr = 0·060 mm.
Avicularia Lz = 0·06 mm.
lz = 0·03 mm.

REMARKS. *Andriopora minor* is characterized by its small size and its closely placed non-caudate zooecia. The oral spine-bases are very minute and not at all thickened.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *G.* [*A.*] *quadrata*.

SPECIMENS. D.39417. Holotype—see above.

2. ***Andriopora major***[1] sp. nov.

(Pl. 23, fig. 5 ; Text-figs. 131, 132)

HOLOTYPE. D.39418. Small zoarial fragment encrusting a piece of echinoid test. Senonian, zone of *B. mucronata*. Norwich, Norfolk. A. W. Rowe Collection.

DIAGNOSIS. *Andriopora* with large, widely elliptical zooecia, each with a long narrow cauda ; orifice higher than wide, with proximal-lateral constrictions ; oral spine-bases 6, prominent, somewhat thickened ; costae 15–19, without lateral costal fusions ; apertural bar sharply bent proximally ; gymnocyst well exposed ; avicularia apparently absent.

DESCRIPTION. *Zoarium* encrusting, uniserial, the angle of branching is apparently maintained, keeping the individual series of zooecia clearly separate. *Zooecia* broadly elliptical, with long, straight *caudae*. *Orifice* higher than wide, rounded distally and laterally with proximal-lateral constrictions, the proximal margin " U "-shaped. *Oral spine bases* 6, somewhat enlarged, tending to be arranged laterally in two groups of three. *Frontal shield* well arched. *Costae* 15–19 (occasionally 13), usually fairly wide. No *lateral costal fusions*. The inner ends of the costae are not up-turned near the mid-line, nor are any pores or costal slits visible. *Median area of fusion* fairly wide. *Apertural bar* with rounded outer ends, narrowed and bent proximally along the mid-line, otherwise not flattened in a vertical plane nor substantially thickened. *Gymnocyst* widely exposed laterally and particularly proximally to the frontal shield where it is prolonged into a straight, rapidly tapering, narrow *cauda*. The gymnocyst also appears on either side of the orifice and tapers rapidly distally giving a rounded distal margin to the zooecium from which the cauda of the distally adjacent zooecium often arises.

Avicularia not seen. There is no evidence of broken avicularia or other heterozooecia, and avicularia are presumed absent in this species. *Ovicells* hyperstomial, globular, smooth, apparently somewhat compressed laterally.

---

[1] With reference to the unusually large size of the zooecia.

MEASUREMENTS. *Zooecia* Lz = 0·72 mm.
lz = 0·42 mm.
hr = 0·15 to 0·18 mm.
lr = 0·12 mm.
Lc = 0·33 to 0·50 mm.
lc = 0·02 mm.

REMARKS. The large size of the zooecia, the widely exposed gymnocyst, and the apparent absence of avicularia are particularly distinctive characters of *Andriopora major*. The orifice is more complex than in *A. minor* and the oral spine-bases more distinct than in that species. The ovicells of *A. major* are described from one broken example present on an isolated zooecium.



131 132

FIGS. 131, 132. *Andriopora major* sp. nov. D.39418. Holotype.
(131) Single adult zooecium ; the cauda worn away.
(132) Part of zoarium showing mode of branching and long, straight, narrow caudae.

STRATIGRAPHICAL DISTRIBUTION. Senonian, zone of *B. mucronata.*

SPECIMEN. D.39418. Holotype—see above.

VIII. SYSTEMATIC BASIS, STRATIGRAPHY AND EVOLUTION OF THE GENERA AND SPECIES REVISED

Recognition of distinct species in the Cretaceous cribrimorph Polyzoa is made difficult by the wide variation in morphology which may occur between adult zooecia of the same and different zoaria of a single species, and between young and adult zooecia of a single zoarium. It is important to assess this range of variation, especially as Cretaceous cribrimorph polyzoan colonies are often broken, so that the extremes of variation—originally part of the same zoarium—become separated and form apparently distinct species. This fragmentation also gives a false idea of the abundance of a particular species at any one horizon or locality.

The present assessment of the species-unit in Cretaceous cribrimorph Polyzoa is an attempt to use all available zooecial characters rather than distinguishing species by the modifications which may be apparent in any one character. Divisions based solely on measurements of the length and width of zooecia, or on zoarial growth habit alone, have been avoided, and, generally, where Lang has distinguished species by these methods they have proved to be synonymous.

In addition to comparative morphology Lang used stratigraphical and astogenetic " criteria of relationship " in his systematic assessment of species. Thus species which differ only very slightly in one or more characters were maintained as distinct if they occurred at well-separated horizons. Species may be maintained on this basis when little material is available for study. The principle has been followed to some extent in the present revision, with the reservation that the slight, but distinct, differences must be essentially structural rather than incidental differences in size or growth habit.

The astogenetic basis on which Lang occasionally distinguished some species has not been followed. He generally used astogenetic differences as indications of the possible course of evolution.

As several of the species established by Lang are here merged it has been necessary to revise the stratigraphical ranges of some species, and the large amount of new material examined has extended the known ranges of most species.

The stratigraphical ranges of the species revised and described here are shown in the accompanying distribution table and discussed subsequently. The range of some species is shown in parts as a broken line indicating that the species has not yet been recorded from that horizon, but that it is thought to range through.

Lang's very extensive studies (1916*a*, *b*, 1919*a–d*, 1921, 1922, 1925, 1929) of Cretaceous Polyzoa, and other groups, led him to adopt an orthogenetic view of evolution, formulated in what has since been termed the " Theory of Trends ". This concept required that, by means of an internal compulsive force, groups of related organisms may follow a predetermined and irreversible course of evolution to a pitch of disharmony with their environment, developing apparently functionless structures ; and that they may undergo periodic, parallel evolution or recapitulation independent of environmental effects. Lang applied this vitalist view of evolution to the Cretaceous cribrimorph Polyzoa (1919*c* : 105 ; 1919*d* : 196 ; 1921 : xvii ; 1922).

This application of the Theory of Trends produced much lively criticism which laid increasing emphasis on environmental adaptations as a more acceptable explanation of the apparent sequence of evolutionary changes (e.g., Bather, 1921 : 18 ; Brydone, 1929 : 6 ; Withers, 1935 : 8).

In the genera revised here, and in other genera of Cretaceous cribrimorph Polyzoa, evolution in the detailed sequence of forms has taken place in a diverging plexus of

Stratigraphical Distribution of Genera and Species Revised



variation, rather than along the confined and determinate, distinct lines of development which Lang has postulated. The present revision merges several of the species, and some genera, recognized by Lang so that his suggested detailed, and even more general, lines of evolution break down. This is particularly so in the case of *Pelmatopora*.

AEOLOPORA

Two species, *A. distincta* and *A. nebulosa*, are here regarded as comprising the genus. By an increase in zooecial size, a thickening of the apertural bar, a slight reduction in the number of costae and an apparent loss of avicularia *A. nebulosa* has evolved from *A. distincta*. The stratigraphical occurrence of both species is discussed on p. 64.

LAGYNOPORA

The stratigraphical ranges of the twelve species of *Lagynopora* recognized here are almost all brief. Only *L. horsleyensis* is recorded from the zone of *M. coranguinum* as well as from the zone of *G.* [*A.*] *quadrata*, and no specimens are as yet recorded from the intervening zones. *L. pustulosa* sp. nov. is stratigraphically the highest species known from the Senonian.

The evolutionary relationships of the twelve species are uncertain. Probably *L. horsleyensis* may be excluded from the main plexus of evolution of the genus as it is the only species with lateral costal fusions. Similarly the small zooecia and other distinctive characters of *L. saltdeanensis* and *L. pustulosa* suggest that the evolution of these two has been somewhat different from that of the remaining nine species of *Lagynopora*. These are generally related, but it is very uncertain that the individual lineage linkages shown by Lang (1921 : 54) are correct.

HEXACANTHOPORA

Only two species are recognized here; *H. sexspinosa*, from the zones of ?*M. coranguinum* to *G.* [*A.*] *quadrata* of the English Chalk, and *H. viginticostata* from the zones of *G.* [*A.*] *quadrata* and *B. mucronata* in Germany. *H. sexspinosa* is not yet recorded from the *Echinocorys scutata* var. *cincta* zubzone of the zone of *O. pilula*.

H. viginticostata may have evolved from *H. sexspinosa* by an increase in the number of costae, a marked widening of the median area of fusion, and by a reduction in the number and type of avicularia.

LEPTOCHEILOPORA

Only four species are recognized here. *L. tenuilabrosa* is the earliest, and *L. filliozati* has the longest range in the Senonian and is represented by the greatest number of specimens. *L. magna* ranges from the Senonian, zone of *B. mucronata*, in England, to the Lower Maastrichtian, in Germany, but is not yet recorded from the Chalk with *Ostrea lunata* in England. *L. vulnerata* is known from only three specimens, one from the Senonian, zone of *B. mucronata*, and two from the Lower Maastrichtian, zone of *Belemnella lanceolata*, from the Chalk with *Ostrea lunata*. *Leptocheilopora* is unrecorded from the upper part of the zone of *Marsupites*.

The sequence *L. tenuilabrosa–L. filliozati–L. vulnerata* probably indicates the main course of evolution in the genus. There is a slight reduction in zooecial size and in the number of costae and a loss of avicularia. By contrast, there is an increase in the number of oral spines from four in *L. tenuilabrosa* and *L. filliozati* to seven in *L. vulnerata*. The structure of the ovicell is simplified from the costate type in *L. tenuilabrosa* to more usual smooth ovicells in the remaining species. In *L. magna*, the youngest species of the genus, there is a pronounced increase in zooecial size and very marked proximal-lateral constrictions in the orifice. The four oral spine bases are very small, but, like the preceding species, there are no avicularia. *L. magna* possibly diverged from the main course of evolution in early *B. mucronata* times.

PELMATOPORA

Lang (1922 : 242) has stated that " though almost certainly arising in the Turonian (since *Sandalopora*, an obvious derivative, has occurred in the Turonian of France) the earliest known forms of *Pelmatopora* are found at the base of the Senonian, and immediately appear in great numbers both of species, and in some species, of individuals ". While it is true to say that several species of *Pelmatopora* make an abrupt appearance at the base of the Senonian it is an overstatement that they are present in great numbers. They occur frequently, but they are by no means so abundant as in the upper part of the Senonian, particularly in the zones of *O. pilula* and *G.* [*A.*] *quadrata*.

The twenty-two species of *Pelmatopora* recognized here fall stratigraphically into two well-defined groups. There is an early Senonian development in the zones of *M. cortestudinarium* and *M. cor-anguinum* of sixteen species, which, with the exception of *P. brydonei* and *P. quadrivolucris*, all have unmodified oral spine-bases. A second, very abundant, group of six species characterizes the higher Senonian zones of *Marsupites* to *G.* [*A.*] *quadrata*. With the exception of *P. simplex*, these all have modified and enlarged distal oral spines. No species of *Pelmatopora* are yet recorded definitely from the *Uintacrinus* Subzone of the *Marsupites* Zone. Recently a definite *Pelmatopora* specimen has been recorded from the Danian of Herfølge, Denmark (see p. 202).

The ranges of certain species of *Pelmatopora* are apparently limited within some zones of the Senonian. Thus *P. pero* and *P. brydonei* are recorded only from the upper part of the zone of *M. cor-anguinum*, while *P. calceata*, *P. crepidaria*, *P. solearis*, *P. gasteri*, *P. fragilis*, *P. quadrata*, *P. suffulta*, *P. plantaris* and *P. quadrivolucris* are most frequent in the lowest quarter of the same zone. *P. chrysalis*, *P. larva*, *P. striata*, *P. d'orbignyi* and *P. fecampensis* apparently characterize the middle part of the zone of *M. cortestudinarium*. *P. simplex* occurs frequently in the zone of *O. pilula*. The remaining five species of *Pelmatopora* are very abundant from the upper part of the zone of *Marsupites* to the upper part of the zone of *G.* [*A.*] *quadrata*, but, unlike the earlier Senonian species, they do not appear to be limited to distinct horizons of these zones.

The broad course of evolution in *Pelmatopora* may be summarized as a development from a Lower Senonian group of smaller, less complex, species with unmodified oral spine bases to an Upper Senonian group of larger, more complex, species with modified, enlarged distal oral spines. Several of the species which Lang recognized in the higher group are synonymous. The almost total absence of *Pelmatopora* species in the *Uintacrinus* Subzone of the *Marsupites* Chalk, and in fact the general absence of Polyzoa and other fossils at this level in the Senonian, may correlate with contemporary deeper water conditions. Calcareous shells of other organisms, such as echinoids and lamellibranchs, are noticeably scarce in the *Uintacrinus* Chalk thus reducing the occurrence of substrates suitable for polyzoan growth. Where larger shelled organisms are more abundant there is a clearly marked increase in the polyzoan fauna. Such an effect would account for the apparent periodicity of evolution which Lang regarded as having occurred during the development of *Pelmatopora*.

The suggested evolution from *P. pero* of the higher group of more complex *Pelmatopora* species (Lang, 1922 : 250) seems less likely than the course of evolution indicated here. *P. quadrivolucris* and *P. brydonei*, in contrast to other species of the Lower Senonian group, both possess distinctly enlarged oral spines and more complex frontal shields, therefore I regard them as fore-runners of the Upper Senonian group of *Pelmatopora* species in which the distal oral spines form a prominent distal oral shield.

P. simplex, which lacks enlarged distal oral spines, is the only species of its kind above the zone of *Marsupites*. Within the Upper Senonian group of more complex species there was, apparently, a tendency to evolve increasingly larger, flatter and more bifurcated distal oral spines. *P. palmata*, in which this tendency is most marked, is closely related to the species-group of *P. marsupitorum*, *P. gregoryi* and *P. somptingensis*.

CASTANOPORA

Fourteen Cretaceous species of *Castanopora* are described here. Only one of these, *C. retrorsa*, represented by an isolated specimen from the *Uintacrinus* Subzone of the *Marsupites* Zone, occurs below the top part of the zone of *B. mucronata*. All other species are of Maastrichtian age—many coming from the boreal Maastrichtian of Rügen. *C. castanea*, *C. dibleyi* and *C. magnifica* have been recorded from the Senonian, zone of *B. mucronata*, but are more common in the Maastrichtian, in the zone of *Belemnella lanceolata* (*Ostrea lunata* Chalk) in England, and in that zone abroad. The appearance of several new species of *Castanopora* in the *Ostrea lunata* Chalk of Trimingham, Norfolk, is a significant indicator of the general faunal change, shown particularly by the foraminifera at this level, between the Senonian, and the Maastrichtian.

Canu & Bassler (1933 : 53, 54) have assigned two species from the Vincentown Limesand to *Rhiniopora* thus extending the upward range of the genus into the Palaeocene.

The main evolutionary spread of *Castanopora* took place at the close of the Senonian, so that *Castanopora* characterizes the Lower Maastrichtian. The species recognized exhibit considerable variation in the number and type of avicularia.

UBAGHSIA

Twelve species of *Ubaghsia* are recognized here. Of these, only *U. gasteri* sp. nov. and *U. langi* sp. nov. occur below the zone of *Belemnitella mucronata*. There is a marked increase in the number of species in the Maastrichtian, most specimens coming either from the boreal Maastrichtian of Rügen, or from the littoral Maastrichtian of Maastricht, Holland.

The two new species, *U. gasteri* and *U. langi*, from the English Senonian (zones of *G.* [*A.*] *quadrata* and *M. cor-anguinum* respectively), extend the downward range of the genus. All the material previously described by Lang, and Jullien's original specimens, were from continental localities, either from the Senonian zone of *Belemnitella mucronata* of Meudon [*U. meudonensis* and *U. demorgani*] and Port Brehay [*U. ocellata*], or from the boreal Maastrichtian of Rügen [*U. perforata*, *U. ranunculus*, *U. crassa* and *U. coaxans*], from the littoral Maastrichtian of Maastricht [*U. ornata* and *U. reticulata*], and from Royan, France [*U. aurita*].

The later species of *Ubaghsia* fall into two main groups. *U. ranunculus*, *U. aurita*, *U. crassa*, *U. coaxans* and (?) *U. perforata* all lacking the very heavy tertiary front wall which characterizes the other group of *U. ocellata*, *U. meudonensis*, *U. demorgani*, *U. reticulata* and *U. ornata*. There is some indication that the marked accumulation of calcium-carbonate, in the tertiary front walls of the second group, may be related to habitat. As with some Cretaceous cirripedes (Withers, 1935 : 11, 13) there seems to be some correlation between a shallow water habitat and marked skeletal development of calcium-carbonate, since the species of the second group occur in littoral Maastrichtian sediments of the type locality, but other species from the boreal Chalk of Rügen are noticeably less robust.

It is thus evident, from the material revised here alone, that the Cretaceous cribrimorph polyzoan faunas of the Chalk contain a number of species and genera with well-marked stratigraphical ranges. Some species apparently have a limited range within the existing recognized zones. Some division of the very thick zones of *M. cor-anguinum* and *B. mucronata* should be possible, on the basis of their contained cribrimorph polyzoan faunas, when further systematic revision of other genera is completed and when new material has been examined from collections made more recently.

IX. REFERENCES

BARNARD, T. & BANNER, F. T. 1953. Arenaceous Foraminifera from the Upper Cretaceous of England. *Quart. J. Geol. Soc. Lond.*, **109** : 173–216, pls. 8, 9.

BASSLER, R. S. 1935. *Bryozoa. Fossilium Catalogus, I. Animalia. Pars* 67. 229 pp. 's-Gravenhage.

—— 1953. *In* MOORE, R. C. Ed. *Treatise on Invertebrate Palaeontology, G. Bryozoa* : i–xiii, G1–253, figs. 1–175. Kansas.

BATHER, F. A. 1907. *A Guide to the Fossil Invertebrate Animals in the Department of Geology and Palaeontology in the British Museum* (*Natural History*). ix + 182 pp., 7 pls. British Museum (Nat. Hist.), London.

—— 1921. Fossils and Life. *Rep. Brit. Ass.*, London, **1920** : 61–68.

BÖHM, J. 1924. Uber eine senone Fauna von Misol. *In* WANNER, J. *Paläontologie von Timor*, **14** : 83–103, pls. 207–211. Stuttgart.

Boswell, P. G. H. 1929. *In* Evans, J. W. & Stubblefield, C. J. *Handbook of the Geology of Great Britain.* 556 pp. London.

Bower, C. R. & Farmery, J. R. 1910. The Zones of the Lower Chalk of Lincolnshire, with a List of New Records from the Red Chalk of the County. *Proc. Geol. Ass. Lond.*, **21** : 333–359, pl. 27.

Bronn, H. G. 1848. *Index Palaeontologicus, A. Nomenclator palaeontologicus.* lxxxiv + 1381 pp. Stuttgart.

—— 1849. *Index Palaeontologicus, B. Enumerator palaeontologicus.* 980 pp. Stuttgart.

Bronn, H. G. & Roemer, F. 1851–52. *Lethaea Geognostica.* 3rd edit., **2**, 5, Kreide-Gebirge. 412 pp. Stuttgart.

Brown, D. A. 1952. *The Tertiary Cheilostomatous Polyzoa of New Zealand.* xii + 405 pp., 296 figs. British Museum (Nat. Hist.), London.

Brydone, R. M. 1900. *The Stratigraphy and Fauna of the Trimmingham Chalk.* 16 pp. London.

—— 1906. Further Notes on the Stratigraphy and Fauna of the Trimmingham Chalk. *Geol. Mag. Lond.* (5) **3** : 13–22, 72–78, 124–131, 289–300, pls. 2–5, 8, 9.

—— 1908*a*. On the Subdivision of the Chalk at Trimmingham (Norfolk). *Geol. Mag. Lond.* (5) **5** : 134–135.

—— 1908*b*. On the Subdivisions of the Chalk at Trimingham (Norfolk). *Quart. J. Geol. Soc. Lond.*, **64** : 401–412, pls. 47, 48.

—— 1910. Notes on new or imperfectly known Chalk Polyzoa. *Geol. Mag. Lond.* (5) **7** : 4–5, 76–77, 145–147, 258–260, 390–392, 481–483, pls. 3, 8, 14, 21, 30, 36.

—— 1912. *The Stratigraphy of the Chalk of Hants.* 116 pp., 3 pls. London.

—— 1913. Notes on new or imperfectly known Chalk Polyzoa. *Geol. Mag. Lond.* (5) **10** : 97–99, 196–199, 248–250, 436–438, pls. 4, 7, 8, 14.

—— 1914*a*. The Zone of *Offaster pilula* in the South English Chalk. *Geol. Mag. Lond.* (6) **1** : 359–369, 405–411, 449–457, 509–513.

—— 1914*b*. Notes on new or imperfectly known Chalk Polyzoa. *Geol. Mag. Lond.* (6) **1** : 97–99, 345–347, 481–483, pls. 4, 24, 25.

—— 1915. The *Marsupites* Chalk of Brighton. *Geol. Mag. Lond.* (6) **2** : 12–15.

—— 1917. Notes on new or imperfectly known Chalk Polyzoa. *Geol. Mag. Lond.* (6) **4** : 49–53, 145–148, 492–496, pls. 3, 11, 32.

—— 1929. *Further notes on new or imperfectly known Chalk Polyzoa.* **1.** 40 pp., 14 pls. London.

—— 1932*a*. The Course of *Marsupites* and *Uintacrinus* across Norfolk (with a note on the Basal Norwich Chalk). *Trans. Norfolk Norw. Nat. Soc.*, **13** : 115–118.

—— 1932*b*. *Uintacrinus* in North Suffolk. J. *Ipswich Dist. Nat. Hist. Soc.*, **1** : 158–161.

—— 1938. *On Correlation of some of the Norfolk Exposures of Chalk with* Belemnitella mucronata. 16 pp. London.

—— 1939. *The Chalk Zone of* Offaster pilula. 8 pp. London.

Butler, E. A. & Cheetham, A. 1958. A new Cretaceous Cribrimorph Bryozoan from Louisiana. *J. Paleont.*, Chicago, **32** : 1153–1157.

Canu, F. 1900*a*. Note preliminaire sur les Bryozoaires [Sénonien] de Tours. *C.R. Ass. franç. Av. Sci.*, Paris, **28** : 406–411.

—— 1900*b*. Révision des Bryozoaires du Crétacé figurés par d'Orbigny. Pt. 2, Cheilostomata. *Bull. Soc. géol. Fr.*, Paris (3) **28** : 334–463, pls. 4–7.

—— 1910. *In* Douville, H. La Craie et le Tertiaire des environs de Royan : 3. Liste des Bryozoaires de la Craie de Royan. *Bull. Soc. géol. Fr.*, Paris (4) **10** : 62–65

—— 1911. Iconographie des Bryozoaires Fossiles de l'Argentine, 2. *An. Mus. nac. B. Aires* (3) **14** : 215–291, pls. 1–12.

Canu, F. & Bassler, R. S. 1920. North American Early Tertiary Bryozoa. *Bull. U.S. Nat. Mus.*, Washington, **106** : x + 878 pp., 162 pls.

—— 1927. Classification of the Cheilostomatous Bryozoa. *Proc. U.S. Nat. Mus.*, Washington, **14** : 1–4, pl. 1.

CANU, F. & BASSLER, R. S. 1929. Contributions to the Biology of the Philippine Archipelago and Adjacent Regions. Bryozoa of the Philippine Region. *Bull. U.S. Nat. Mus.*, Washington, **100**, 9 : xi + 685 pp., 94 pls.

—— 1933. The Bryozoan Fauna of the Vincentown Limesand. *Bull. U.S. Nat. Mus.*, Washington, **165** : 1–108, pls. 1–21.

COQUAND, H. 1860. *Description physique, géologique, paléontologique et minéralogique de département de la Charente*, **2**. 420 pp. Marseille.

DEECKE, W. 1895. Die Mesozoischen Formationen der Provinz Pommern. *Mitt. naturw. Ver. Griefswald*, Berlin, **26** : 1–115.

DIDON, J. 1958. Bryozoaires Cribrimorphes récoltés dans la craie santonienne de Normandie. *Rev. Soc. Sav. Haute-Normandie Sci.*, Rouen, **9** : 79–87, figs. 1–12.

DINES, H. G., HOLMES, S. C. A. & ROBBIE, J. A. 1954. Geology of the Country around Chatham (One-inch Geological Sheet 272, New Series). vii + 157 pp., 5 pls. *Mem. Geol. Surv. Gt. Britain*, London.

DUMONT, A. H. 1832. Mémoire sur la constitution géologique de la province de Liège. *Mém. Acad. R. Belg.*, Bruxelles, 8, 7 : 1–374, pls. 1–3.

—— 1849. Rapport sur la carte géologique du Royaume. *Bull. Acad. R. Belg.*, Bruxelles, **16**, 2 : 351–373.

EDWARDS, H. M. 1836. *In* LAMARCK, J. P. B. A., de MONNET DE. *Histoire naturelle des Animaux sans Vertèbres.* 2nd edit., **2**. 684 pp. Paris.

FERUGLIO, E. 1949. Descripción geologica de la Patagonia. *Min. Ind. Comerc. nac. Direc. Gen. Yacimientos Petrol. Fisc.* (*Rep. Argentina*) **2** : 1–349, pls. 61–80. Buenos Aires.

FRIČ, A. 1889. Studien im Gebiete der bohmischen Kreideformation, 4. Der Teplitzer-Schichten (1). *Arch. naturw. LandesForsch. Böhm.*, Prag, **7**, 2 : 1–120.

GASTER, C. T. A. 1920. An undescribed species of *Trochiliopora*. *Geol. Mag. Lond.*, **57** : 526 (letter).

—— 1924. The Chalk of the Worthing District, Sussex. *Proc. Geol. Ass. Lond.*, **35** : 89–110, pl. 8.

—— 1930. Chalk Zones in the neighbourhood of Shoreham, Brighton and Newhaven, Sussex. *Proc. Geol. Ass. Lond.*, **40** : 328–340, pl. 27.

—— 1932. The Zones of the Chalk of the Arun Gap, Sussex : with a description of new species of *Bicavea*. *Proc. Geol. Ass. Lond.*, **43** : 212–223, pl. 9.

—— 1937. The Stratigraphy of the Chalk of Sussex. Part 1. West Central Area—Arun Gap to Valley of the Adur, with Zonal Map. *Proc. Geol. Ass. Lond.*, **48** : 356–373, pl. 34.

—— 1939. The Stratigraphy of the Chalk of Sussex. Part 2. Eastern Area—Seaford to Cuckmere Valley and Eastbourne, with Zonal Map. *Proc. Geol. Ass. Lond.*, **50** : 510–526.

—— 1941. The Chalk Zones of *Offaster pilula* and *Actinocamax quadratus*. *Proc. Geol. Ass. Lond.*, **52** : 210–215.

—— 1944. The Stratigraphy of the Chalk of Sussex. Part 3. Western Area—Arun Gap to the Hampshire Boundary, with Zonal Map. *Proc. Geol. Ass. Lond.*, **55** : 173–188, pl. 9.

—— 1951. The Stratigraphy of the Chalk of Sussex. Part 4. East Central Area—Between the Valley of the Adur and Seaford, with Zonal Map. *Proc. Geol. Ass. Lond.*, **62** : 31–64, pl. 3.

GEINITZ, H. B. 1849–50. *Das Quadersandsteingebirge oder Kreidegebirge in Deutschland.* iv + 293 pp., 12 pls. Freiberg.

GILLARD, P. A. 1940. Sur les bryozoaires néocrétacé des environs de Royan, 1. Cheilostomata. *C.R. Soc. géol. Fr.*, Paris, **1940** : 14–16.

—— 1942. Les bryozoaires du forage de Montbazon (feuille de Loches au 1/80,000°). *Bull. Serv. Carte géol. Fr.*, Paris, **43** : 89–92.

—— 1943. Sur quelques Bryozoaires cribrimorphes du Campanien supérieur des environs de Royan. *Bull. Soc. géol. Fr.*, Paris (5) **12** : 185–188.

GOLDFUSS, G. A. 1826. *Petrefacta Germaniae*, **1**, 1 : 1–76, pls. 1–25. Düsseldorf.

—— 1862. *Petrefacta Germaniae*, **1**. 2nd edit. iv + 234 pp., atlas (1862–63), 201 pls. Leipzig.

GREGORY, J. W. 1894. On some Jurassic Species of Cheilostomata. *Geol. Mag. Lond.* (4) **1** : 61–64.

—— 1896. *Catalogue of the Fossil Bryozoa in the Department of Geology, British Museum (Natural History). The Jurassic Bryozoa.* 239 pp., 11 pls. British Museum (Nat. Hist.), London.

GRIFFITH, C. & BRYDONE, R. M. 1911. *The Zones of the Chalk in Hants.* 36 pp., 4 pls. London.

HAGENOW, K. F. VON. 1846. *In* GEINITZ, H.B. *Grundriss der Versteinerungskunde.* Pp. 586–635. Dresden & Leipzig.

—— 1851. *Die Bryozoen der Maastrichter Kreidebildung.* xvi + 111 pp., 12 pls. Cassel.

HARMER, S. F. 1902. On the Morphology of the Cheilostomata. *Quart. J. Micr. Sci.*, London (N.S.) **46** : 263–350, pls. 15–18.

—— 1926. The Polyzoa of the Siboga Expedition, 2. Cheilostomata, Anasca. *Siboga Exped.*, Leiden, **28**B : viii, 181–501, pls. 1–22.

—— 1930. Polyzoa. *Proc. Linn. Soc. Lond.*, **1928–29** : 68–118, pl. 1.

—— 1931. Recent work on Polyzoa. *Proc. Linn. Soc. Lond.*, **1930–31** : 113–168.

HAUG, E. 1910. *Traité de Géologie, II. Les Périodes géologiques,* **2** : 929–1396, pls. 101–119. Paris.

HEIDE, S. VAN DER. 1954. The Original Meaning of the Term Maastrichtian (Dumont, 1849). *Geol. en Mijnb.*, Den Haag (N.S.) **16** : 509–511.

HILTERMANN, H. 1952. Stratigraphische Fragen des Campan und Maastricht unter besonderer Berüchsichtigung der Mikropaläontologie. *Geol. Jber.*, Berlin, **67** : 47–66.

HILTERMANN, H. & KOCH, C. 1950. Taxonomie und Vertikalverbreitnung von *Bolivinoides*—Arten im Senon Norwestdeutschlands. *Geol. Jber.*, Berlin, **65** : 595–632.

ILLIES, G. 1953. Variationsstatische Untersuchungen an *Rhiniopora cacus* (Bryd.). Bryoz. Cheil. aus der Oberkreide von Henmoor/Niederelbe. *Mitt. min.-geol. (St) Inst. Hamburg,* **22** : 76–101, pls. 15, 16.

JELETZKY, J. A. 1951. The Place of the Trimingham and Norwich Chalk in the Campanian-Maestrichtian Succession. *Geol. Mag. Lond.*, **88** : 197–208.

JUKES-BROWNE, A. J. 1904. The Cretaceous Rocks of Britain, 3. The Upper Chalk of England. x + 566 pp., 9 pls. *Mem. Geol. Surv. Gt. Britain*, London.

JULLIEN, J. 1883. Dragages ud *Travailleur*. Bryozoaires. Espèces draguées dans l'ocean Atlantique en 1881. *Bull. Soc. zool. Fr.*, Paris, **7** : 497–529, pls. 13–17.

—— 1886. Les Costulidées, nouvelle Famille des Bryozoaires. *Bull. Soc. zool. Fr.*, Paris, **11** : 601–620, pls. 17–20.

KLÖDEN, K. F. VON. 1834. *Die Versteinerungen der Mark Brandenburg, insonderheit diejenigen welche sich in den Rollsteinen und Blöchen der südbaltischen Ebene finden.* x + 378 pp., 10 pls. Berlin.

KÜHN, O. 1930. Das Danien der äusseren Klippenzone bei Wien. *Geol. paläont. Abh.*, Jena, **21** : 495–576, pls. 1, 2.

LAGAAIJ, R. 1952. The Pliocene Bryozoa of the Low Countries and their bearing on the marine stratigraphy of the North Sea Region. *Meded. geol. Sticht.*, 'sGravenhage (N.S., C), **5**, 5 : 1–233, pls. 1–25.

LAMARCK, J. P. B. A. & DE MONNET DE. 1816. *Histoire naturelle des Animaux sans Vertèbrés,* **2**. 586 pp. Paris.

LANG, W. D. 1913. Report of a visit to the exhibits of Polyzoa and Corals in the Geological Dept., of the B.M. (Nat. Hist.). *Proc. Geol. Ass. Lond.*, **24** : 169–173.

—— 1916*a*. A Revision of the "Cribrimorph" Cretaceous Polyzoa. *Ann. Mag. Nat. Hist.*, London (8) **18** : 81–112, 381–410.

—— 1916*b*. Calcium carbonate and Evolution in Polyzoa. *Geol. Mag. Lond.* (6) **3** : 73–77.

—— 1917. The Genotypes of certain Polyzoan Genera. *Geol. Mag. Lond.* (6) **4** : 169–174.

—— 1919*a*. The Kelostominae : a Subfamily of Cretaceous Cribrimorph Polyzoa. *Quart. J. Geol. Soc. Lond.*, **74** : 204–220.

—— 1919*b*. The Evolution of Ammonites. *Proc. Geol. Ass. Lond.*, **30** : 49–65.

LANG, W. D. 1919*c*. Old Age and Extinction in Fossils. *Proc. Geol. Ass. Lond.*, **30** : 102–113.

—— 1919*d*. The Pelmatoporinae, and Essay on the Evolution of a Group of Cretaceous Polyzoa. *Philos. Trans.*, London (B) **209** : 191–228.

—— 1921. *Catalogue of the Fossil Bryozoa (Polyzoa) in the Department of Geology, British Museum (Natural History). The Cretaceous Bryozoa (Polyzoa), 3. The Cribrimorphs, Part 1.* cx + 269 pp., 8 pls. British Museum (Nat. Hist.), London.

—— 1922. *Catalogue of the Fossil Bryozoa (Polyzoa) in the Department of Geology, British Museum (Natural History). The Cretaceous Bryozoa (Polyzoa), 4. The Cribrimorphs, Part 2.* xii + 404 pp., 8 pls. British Museum (Nat. Hist.), London.

—— 1925. Persistence in Fossils. *Proc. Geol. Ass. Lond.*, **36** : 227–239, pls. 20–22.

—— 1929. Form in Fossils. *Proc. Geol. Ass. Lond.*, **39** : 429–444.

LEVINSEN, G. M. R. 1907. Sur la régénération totale des Bryozoaires. *Overs. danske Vidensk. Selsk. Forh.*, Kjøbenhavn, **4** : 151–159, pl. 1.

—— 1909. *Morphological and Systematic Studies on the Cheilostomatous Bryozoa.* vii + 431 pp., 24 pls. Copenhagen.

—— 1925. Undersøgelser over Bryozoerne ; den danske Kridtformation Efter Forfattersen, Død vdg. af Dr. K. Brünnich Nielsen og Dr. T. Mortensen. *K. danske Vidensk. Selsk.*, Kjøbenhavn, **7** : 283–445, pls. 1–8.

LOEBLICH, A. R. & TAPPAN, H. 1957. Correlation of the Gulf and Atlantic Coastal Plain Paleocene and Lower Eocene Formations by means of Planktonic Foraminifera. *J. Paleont.*, Chicago, **31** : 1109–1137.

MARCUS, E. 1926. Beobachtungen und Versuche an lebenden Meeresbryozoen. *Zool. Jber*, Berlin, **52** : 1–102, pls. 1, 2.

MARSSON, T. F. 1887. Die Bryozoen der weissen Schreibkreide der Insel Rügen. *Paläont. Abh.*, Berlin, **4** : 1–112, pls. 1–10.

MORREN, C. F. A. 1828. Descriptio coralliorum fossilium in Belgio repertorum. *Ann. Acad. Groningen*, **1828** : 1–76, pls. 1–7.

MOURLON, M. F. 1881. *Géologie de la Belgique*, **2**. xvi + 392 pp. Brussels.

NORMAN, A. M. 1903. Notes on the Natural History of East Finmark. Polyzoa. *Ann. Mag. Nat. Hist.*, London (7) **12** : 87–128, pls. 8, 9.

NOVÁK, O. 1877. Beitrag zur Kenntniss der Bryozoen der böhmischen Kreideformation. *Denkschr. Akad. Wiss. Wien*, **37** : 79–126, pls. 1–10.

ORBIGNY, A. D'. 1850. *Prodrome de Paléontologie stratigraphique*, **2**. iii + 428 pp. Paris.

—— 1851–54. *Paléontologie Française. Description des Animaux Invertèbrés : Terrain Crétacé, 5. Bryozoaires.* 1192 pp., pls. 600–800. Paris.

OSBURN, R. C. 1940. Bryozoa of Porto Rico with a resume of the West Indian Bryozoan Fauna. *Sci. Surv. P. Rico*, **16** : 321–486, pls. 1–9.

—— 1950. Bryozoa of the Pacific Coast of America, II. Cheilostomata—Anasca. *Allen Hancock Pacif. Exped.*, Los Angeles, **14** : 1–270, pls. 1–29.

OSSWALD, VON. 1890. Die Bryozoen der mecklenbergischen Kreidegeschiebe. *Arch. Ver. Naturg. Mecklenb.*, Güstrow, **43** : 101–110.

PEAKE, N. B. & HANCOCK, J. M. 1961. The Upper Cretaceous of Norfolk. *Trans. Norfolk Norw. Nat. Soc.*, **19** : 293–339, 9 figs.

PERGENS, E. 1889. Zur fossilen Bryozoenfauna von Wola Lu'zanska. *Bull. Soc. belge Géol. Pal. Hydr.*, Bruxelles, **3** (Mems.) : 59–72.

—— 1893. Bryozoaires du Sénonien de Sainte-Patern, de Lavardin et de la Ribochère. *P. V. Soc. belge Géol.*, Bruxelles, **6** : 200–217.

PETTIT, N. E. 1954. *Rhynchonellidae of the British Chalk*, **2** : 27–52, pl. 1. Palaeontogr. Soc. [Monogr.] London.

PHLEGER, F. B., PARKER, F. L. & PEIRSON, J. F. 1953. Sediment Cores from the North Atlantic Ocean (1) North Atlantic Foraminifera. *In* PETTERSSON, H. Reports of the Swedish Deep-Sea Expedition, 1947–1948. *Göteborgs VetenskSamh. Handl.*, **7** : 1–122, pls. 1–12.

PICTET, F. J. 1857. *Traité de Paléontologie.* 2nd edit. **4** : xvi + 768 pp., atlas (1853–57), 77 pp., 110 pls. Paris.

QUENSTEDT, F. A. 1879. *Petrefactenkunde Deutschlands,* **6** : 218–363, atlas (1878), pls. 153–155.

REUSS, A. E. VON. 1874. *In* GEINITZ, H. B. Das Elbthalgebirge in Sachsen, 2. Die Bryozoen des oberen Planers. *Palaeontographica,* Stuttgart, **20** : 127–138, pls. 24–26.

ROEMER, F. A. VON. 1840. *Die Versteinerungen des Nordeutschen Kreidegebirge.* iv + 145 pp., 16 pls. Hannover.

ROWE, A. W. 1900. The Zones of the White Chalk of the English Coast : Part 1. Kent and Sussex. *Proc. Geol. Ass. Lond.,* **16** : 289–368, pls. 8–10.

—— 1904. The Zones of the White Chalk of the English Coast : Part IV. Yorkshire. *Proc. Geol. Ass. Lond.,* **18** : 193–296, pls. 17–40.

—— 1908. The Zones of the White Chalk of the English Coast : Part V. The Isle of Wight. *Proc. Geol. Ass. Lond.,* **20** : 209–352, pls. 8–23.

SILEN, L. 1944. The Main Features of the Development of the Ovum, Embryo, and Ooecium in the Ooeciferous Bryozoa Gymnolaemata. *Ark. Zool.,* Uppsala, **35***A*, 17 : 1–34.

SPATH, L. F. 1926. On New Ammonites from the English Chalk. *Geol. Mag. Lond.,* **63** : 77–83.

STEPHENSON, L. W., KING, P. B., MONROE, W. H. & IMLAY, R. W. 1942. Correlation of the Outcropping Cretaceous Formations of the Atlantic and Gulf Coastal Plain and Trans-Pecos Texas. *Bull. Geol. Soc. Amer.,* Rochester. N.Y., **53** : 435–448. pl. 1.

SPENCER, W. K. 1913. The Evolution of the Cretaceous Asteroidea. *Philos. Trans.,* London (B) **204** : 99–177, pls. 10–16.

UBAGHS, J. C. 1865. Die Bryozoen-Schichten der Maastrichter Kreidebildung, nebst einige neuen Bryozoen-Arten aus der Maastrichter Tuff-Kreide. *Verh. narturh. Ver. preuss. Rheinl.,* Bonn, **22**, 3, 2 : 31–62, pls. 2, 2*a*, 3.

—— 1879. *Description géologique et paléontologique du Sol du Limbourg.* ii + 276 pp., 7 pls. Ruremonde.

—— 1889. Quelques considérations sur l'âge de la Craie tuffeau de Folx-les-Caves. *Bull. Soc. belge Géol. Pal. Hydr.,* Bruxelles, **2** (Méms.) : 49–57.

VINE, G. R. 1885. Fifth and Last Report of the Committee appointed for the completion of a Report on the Cretaceous Polyzoa. *Rep. Brit. Ass.,* London, **1884** : 97–219.

—— 1893. Report of the Committee appointed for the completion of a Report on the Cretaceous Polyzoa. *Rep. Brit. Ass.,* London, **1892** : 301–337.

VOIGT, E. 1925*a*. Uber das Vorkommen von Bryozoen in Diluvialgeschieben und die Grundzüge ihrer Systematik. *Z. Geschiebeforsch.,* Berlin, **1** : 13–28, pl. 1.

—— 1925*b*. Neue Cribrimorphe Bryozoen aus der Familie der Pelmatoporidae in Kreidegeschieben Anhalts. *Z. Geschiebeforsch.,* Berlin, **1** : 97–104, pl. 3.

—— 1925*c*. Gehört das Danien zum Tertiär ? *Z. Geschiebeforsch.,* Berlin, **1** : 172–186.

—— 1927. Review of Levinsen (1925). *Z. Geschiebeforsch.,* Berlin, **3** : 101–103.

—— 1930. Morphologische und stratigraphische Untersuchungen über die Bryozoenfauna der oberen Kreide, 1. Die cheilostomen Bryozoen der jüngeren Oberkreide in Nordwestdeutschland, im Baltikum und in Holland. *Leopoldina,* Halle a.S., **6** : 379–579, pls. 1–39.

—— 1939. Über die Dornenspezialisation bei cheilostomen Bryozoen und die Nichtumkehrbarkeit der Entwicklung. *Paläont. Z.,* Berlin, **21** : 87–107.

—— 1949. Cheilostome Bryozoen aus der Quadratenkreide Nordwestdeutschlands. *Mitt. min.-geol. (St) Inst. Hamburg,* **19** : 1–49, pls. 1–11.

—— 1951. Das Maastricht-Vorkommen von Ilten bei Hannover und seine Fauna mit besonderer Berüchsichtigung der Gross-Foraminifera und Bryozoen. *Mitt. min.-geol. (St) Inst. Hamburg,* **20** : 15–109, pls. 1–10.

WATERS, A. W. 1923. Mediterranean and other Cribrilinidae, together with their Relationship to Cretaceous forms. *Ann. Mag. Nat. Hist.,* London (9) **12** : 545–573, pls. 17, 18.

WHITAKER, W. 1872. The Geology of the London Basin. Part 1. The Chalk and the Eocene Beds of the Southern and Western Tracts. xi + 619 pp. *Mem. Geol. Surv. Gt. Britain,* London.

WHITE, H. J. O. 1909. The Geology of the Country around Basingstoke. Explanation of Sheet 284. v + 119 pp. *Mem. Geol. Surv. Gt. Britain*, London.

—— 1910. The Geology of the Country around Alresford. Explanation of Sheet 300. iv + 102 pp. *Mem. Geol. Surv. Gt. Britain*, London.

—— 1912. The Geology of the Country around Winchester and Stockbridge. Explanation of Sheet 299. iv + 89 pp. *Mem. Geol. Surv. Gt. Britain*, London.

—— 1913. The Geology of the Country near Fareham and Havant. Explanation of Sheet 316. iv + 96 pp. *Mem. Geol Surv. Gt. Britain*, London.

—— 1924. The Geology of the Country near Brighton and Worthing. Explanation of Sheets 318 and 333. vi + 114 pp., 4 pls. *Mem. Geol. Surv. Gt. Britain*, London.

WICHER, C. A. 1949. On the Age of the Higher Upper Cretaceous of the Tampico Embayment Area in Mexico, as an example of the worldwide existence of micro-fossils and practical consequences arising from this. *Bull. Mus. Hist. Nat. Pays Serbe* (A) **2** : 76–105, pls. 2–8.

—— 1953*a*. Mikropaläontologische Beobachtungen in der höheren borealen Oberkreide, besonders im Maastricht. *Geol. Jber.*, Berlin, **68** : 1–26.

—— 1953*b*. Beobachtungen im borealen Maastricht. *Paläont. Z.*, Berlin, **27** : 233–236.

WILLIAMS-MITCHELL, B. 1948. The Zonal Value of Foraminifera in the Chalk of England. *Proc. Geol. Ass. Lond.*, **59** : 91–112, pls. 8–10.

WITHERS, T. H. 1935. *Catalogue of Fossil Cirripedia in the Department of Geology, British Museum (Natural History)*, **2**. *Cretaceous*. xiii + 433 pp., 50 pls. British Museum (Nat. Hist.), London.

WOODWARD, H. 1897. *A Guide to the Fossil Invertebrates and Plants in the Department of Geology and Palaeontology*. xvi + 158 pp., 182 figs. British Museum (Nat. Hist.), London.

WRIGHT, C. W. 1947. *In* ARKELL, W. J. The Geology of the Country around Weymouth Swanage, Corfe and Lulworth. Explanation of Sheets 341, 342, 343, and with small portions of Sheets 327, 328, 329. xii + 386 pp., 19 pls. *Mem. Geol. Surv. Gt. Britain*, London.

WRIGHT, C. W. & E. V. 1942. The Chalk of the Yorkshire Wolds. *Proc. Geol. Ass. Lond.*, **53** : 112–127.

—— 1951. *A Survey of the Fossil Cephalopoda of the Chalk of Great Britain*. 40 pp. Palaeontogr. Soc. [Monogr.] London.

INDEX

New taxonomic names and the page numbers of the principal references are printed in **Clarendon** type. An asterisk (*) indicates a figure.

Aeolopora 15, 23, 34, 29*, 30, 40, 52, 53, **54**, 272
—*distincta* 33*, 43, 44, 54, **55**, 56*, 57*, 58*, 64, 272 ; Pl. 2, figs. 1–7
—*nebulosa* 15, 33*, 55, **62**, 63*, 272 ; Pl. 2, figs. 8, 9
—*stellata* 58, 59
Amphiblestrum 36
—*cervicorne* 37*
ancestrulae 45
Andriopora 23, **266**
—*frequens* 55
—*gallica* 55
—**major** 33*, **268**, 269* ; Pl. 23, fig. 5
—**minor** 32, 33*, **266**, 267*, 269 ; Pl. 23, fig. 6
Anornithopora 54
apertural bars 32
avicularia 41

Barroisina galeata 106
Batrachopora 52, **240**, 241, 244, 264
—*coaxans* 253
—*crassa* 250, 252, 253, 266
—*hyla* 260
—*ornata* 260
—*peltata* 266
—*perforata* 266
—*ranunculus* 242
—*royanensis* 242, **265**
—*ultima* 242, **265**, 266
Belemnella lanceolata, Zone of 12
Belemnitella mucronata, Zone of 11
Brydone, R. M. 19, 20

Canupora prima 66, 67
Carydiopora 54, 170
Castanopora 15, 24, 29*, 40, 41, 43, 52, 53, 202, **203**, 274
—**armata** 41, 205, **206**, 207* ; Pl. 16, fig. 1
—*aviculosa* 15, 206, **226**, 227* ; Pl. 19, fig. 3
—*castanea* 15, 33*, 39, 203, 205, **211**, 212*, 274 ; Pl. 16, figs. 3–5
—*dibleyi* 15, 39, 205, **215**, 218*, 274 ; Pl. 16, figs. 6–8, Pl. 17, figs 1, 2
—*faujasi* 15, 206, **236**
Castanopora glandulosa 15, 206, **225**, 226* ; Pl. 18, fig. 1
—*guascoi* 15, 205, **219**
—*juglans* 214
—*jurassica* 15, 206, **223**, 224* ; Pl. 17, figs. 3–5
—*labiata* 15, 41, 42*, 205, **208**
—*magnifica* 15, 25, 33*, 206, **228**, 230*, 274 ; Pl. 18, figs. 2–7, Pl. 19, figs. 1, 2, 4–6 ; Pl. 20, fig. 1
—*multicostata* 16, 206, **221**
—*nucifera* 214
—*retrorsa* 205, **209**, 210, 217, 274 ; Pl. 16, fig. 2
—*spooneri* 16, 206, **220**
—**voigti** 16, 22, 206, **236**
Cellepora perforata 206, **237**
—(*Discopora*) *signata* 242, **264**
Cenomanian 8
Chaperia 36
—*cervicornis* 37*
—*galeata* 37*
—*quadrispinosa* 36*
Colletosia bellula 111
Corymbopora 23
costae 26
Cribrilina 264
—*alcicornis* 36*
—*annulata* 27*
—*balzaci* 27*
—*brachiata* 255
—*cacus* 234
—*fragilis* 139, 144
—*furcifera* 82
—*galanthis* 163
—*gregoryi* 163
—*innominata* 27*
—*insignis* 155, 156
—*labiata* 209
—*perforata* 255
—*radiata* 26
—*repleta* 153, 154, 163
—*suffulta* 154, 163
—*tumuliformis* 110, 111
—*valida* 209

Davis, A. G. 13
Decurtaria 54
Diacanthopora 30
Dishelopora 24, 30
Disteginopora 54, 241
—*horrida* 241

Electra pilosa 23
Escharipora chrysalis 133
—*guascoi* 220
—*magnifica* 220
—*mumia* 137
—*pretiosa* 220
—*striata* 133, 137
Escharoides peltata 242, **266**
Eucheilopora 55

Figularia 111
—*philomela* 111
—*spatulata* 111
frontal shield 25
—wall 25

Gaster, C.T. A. 5, 19
Gephyrotes nitido-punctata 27*
Gonioteuthis [*Actinocamax*] *quadrata*, Zone of 11
Granulatensenon 19
gymnocyst 25

Haplocephalopora 42
Hastings, Anna B. 7
Hesperopora 54
heterozooecia other than avicularia 43
Hexacanthopora 16, 24, 25, 30, 31, 40, 52, 53, 65, **97**, 272
—*brightonensis* 99, 100
—*kintburiensis* 99, 100
—*sexspinosa* 16, 25, 32, 33*, 39, 43, 97, **98**, 272 ; Pl. 5, figs. 3–6
—*viginticostata* 16, 97, **102**, 272
Hippiopora 34
Histricopora 30
Holaster planus, Zone of 9
—*subglobosus*, Zone of 8
Hyphantoceras reussianum, Subzone of 9

Ichnopora 54
—*denticulata* 263
Inoceramus labiatus, Zone of 8
intercostal spaces 31
interzooecial secondary tissue 40

Lagynopora 16, 24, 25, 30, 31, 32, 35, 39, 40, 43, 52, 53, **65**, 66, 97, 272
—*amphora* 39, 66, **77*** ; Pl. 3, fig. 9
—*ampulla* 33*, 39, 66, 78, **79*** ; Pl. 3, figs. 7, 8
—*birlingensis* 39, 66, 74, **75*** ; Pl. 3, fig. 6
—*furcifera* 16, 39, 66, **81*** ; Pl. 4, figs. 1, 2
—*horsleyensis* 16, 66, **68**, 69*, 250, 272 ; Pl. 3, figs. 2, 3
—*lagena* 16, 39, 65, 66, **86**, 87* ; Pl. 4, figs. 4–6
—*ollula* 39, 66, 82, **83**, 84*, 95, 97 ; Pl. 4, fig. 3
—*praecursor* 39, 66, **71*** ; Pl. 3, fig. 5
—*prima* 16, 67, **96**, 97
—**pustulosa** 33*, 66, **67***, 272 ; Pl. 3, fig. 1
—*saltdeanensis* 66, **73***, 76, 272 ; Pl. 3, fig. 4
—*urceolus* 39, 66, **90**, 91*, 92* ; Pl. 4, fig. 7, Pl. 5, fig. 1
—*vasculum* 39, 66, 82, 83, **93**, 94* ; Pl. 5, fig. 2
Lang, W. D. 5, 7
lateral costal fusions 30
Lepralia otto-mulleriana 26, 28*, 30
—*pediculus* 66, 67, 95
Leptocheilopora 16, 25, 30, 31, 40, 43, 52, 53, 97, **102**, 272
—*arcuata* 114
—*filliozati* 16, 32, 33*, 34, 35, 39, 103, **112**, 113*, 115*, 272, 273 ; Pl. 7, figs. 3, 4, Pl. 8
—*gasteri* 109
—*longuessensis* 106, **118**
—*magna* 16, 25, 30, 35, 39, **103**, 104*, 105*, 113, 115, 272, 273 ; Pl. 6, figs. 1–7
—*regularis* 119
—*tenuilabrosa* 35, 41, 43, 44, 103, **107**, 108*, 109*, 110*, 111*, 115, 272, 273 ; Pl. 6, figs. 8–11, Pl. 7, figs. 1, 2
—*vulnerata* 33*, 39, 97, 103, 115, **116**, 117*, 272, 273 ; Pl. 9, figs. 1, 2
Lithothamnienkalk of Haidhof 19
lumen pores 27

Maastrichtian 12
Mammillatensenon 19
Marsupites testudinarius, Zone and Subzone of 10
median area of fusion 31
Membranipora 223
—*crenulata* 119
—*meudonensis* 255
—*nitida* 26, 31

Membranipora ocellata 255
—*reticulata* 255
Micraster cor-anguinum, Zone of 9
—*cortestudinarium*, Zone of 9
Møen Chalk 19
Monoceratopora 41
Murinopsia 54

Offaster pilula, Zone of 10
oral spines 35
—shields 39
orifice, primary 34
—secondary 35
Ostrea lunata, Chalk with 13
ovicells 43

Pachydera 54
Pancheilopora 55
pelmata 27
pelmatidia 27
Pelmatopora 17, 22, 25, 30, 32, 34, 35, 36, 39, 40, 41, 43, 52, 53, **119**, 272, 273
—*bidens* 181
—*brydonei* 17, 21, 38, 123, 150, **164**, 165*, 166, 273, 274; Pl. 12, fig. 3
—*calceata* 40, 41, 42*, 122, **123**, 124*, 129, 273; Pl. 9, fig. 3
—*chrysalis* 17, 122, **132**, 133*, 136, 137, 273; Pl. 9, fig. 7
—*collium* 181
—*coryli* 38, 123, **168**, 169*; Pl. 12, fig. 4
—*crepidaria* 40, 122, **125**, 126*, 273; Pl. 9, fig. 4
—*damicornis* 198
—*daniensis* 17, 200, **202**
—*danktonensis* 192
—*d'orbignyi* 17, 33, 122, **140**, 141*, 142*, 273; Pl. 10, fig. 3
—*fecampensis* 17, 40, 44, 122, **145**, 146*, 147*, 148*, 273; Pl. 10, fig. 6
—*filliozati* 139, 144
—*fragilis* 17, 122, **138***, 273; Pl. 10, figs. 1, 2
—*gasteri* 122, **130**, 131*, 273; Pl. 9, fig. 8
—*grandiporosa* 201
—*gregoryi* 21, 38, 40, 42*, 44, 123, 146, 154, 166, **175**, 178*, 179*, 180*, 192, 193, 198, 200, 274; Pl. 13, Pl. 14, figs. 1–4
—*gyrinoides* 181
—*incerta* 200
—*insignis* 154, 155
—*interrupta* 136
—*lacuum* 181
—*lancingensis* 181
Pelmatopora larva 17, 22, 122, **134***, 142, 273; Pl. 9, fig. 9
—*marsupitorum* 17, 38, 42*, 44, 123, **171**, 172*, 192, 274; Pl. 12, figs. 2, 5
—*palmata* 33*, 38, 123, 193, **196**, 197*, 274; Pl. 15, figs. 3, 5, 6
—*pauciclavia* 139
—*pero* 120, 123, **148**, 149*, 167, 273, 274; Pl. 10, fig. 7
—*plantaris* 123, **159**, 160*, 273; Pl. 11, fig. 4
—*promontoriorum* 181
—*quadrata* 17, 44, 122, **143**, 144*, 273; Pl. 10, figs. 4, 5
—*quadrivolucris* 38, 123, **161**, 162*, 273, 274; Pl. 11, figs. 5, 6, Pl. 12, fig. 1
—*ranunculoides* 181
—*repleta* 153, 154, 155
—*roedeanensis* 173
—*saltdeanensis* 181
—*simplex* 123, **157**, 158*, 161, 273, 274; Pl. 11, figs. 2, 3
—*solearis* 33*, 40, 122, **127***, 128*, 273; Pl. 9, figs. 5, 6
—*somptingensis* 21, 24, 31, 33*, 38, 41, 42*, 44, 123, 146, **188**, 190*, 191*, 274; Pl. 14, figs. 5, 6, Pl. 15, figs. 1, 2, 4
—*spp.* **199**, 201; Pl. 15, fig. 7
—*striata* 17, 122, **136**, 273; Pl. 9, fig. 10
—*suffulta* 17, 123, **151**, 152*, 163, 273; Pl. 10, figs. 8, 9, Pl. 11, fig. 1
Pelmatoporidae 51
Pelmatoporinae 52
Phrynopora 54, 241
Pliophloea 30, 43
Pnictopora 22, 40
Polycephalopora 21
Prodromopora 52, 66, 76, 96
—*praecursor* 66, 72

Reptescharella radiata 202, 206, 237
Repteschariporа convexa 242, **264**
—*ornata* 206, **238**
Reptoporella 103, 119
—*regularis* 103, **119**
Rhabdopora 22
Rhiniopora 52, 202, **204**, 237, 274
—*aspera* 231, 232, 233, 234
—*asperula* 231, 232, 233
—*aviculosa* 227, 228
—*cacus* 231, 232, 233, 234
—*hispida* 206, **236**, 237, 238
—*horrida* 232, 233

Rhiniopora jurassica 223, 224
—*scabra* 223, 224
—*spooneri* 221
" *Rhynchonella* " *cuvieri*, Zone of 8
Rocanean 19
Rügen Chalk 19

Sandalopora 22, 54, 273
Schloenbachia varians, Zone of 8
Scrupocellaria 36
—*frondis* 37*
—*securifera* 37*
Semiescharipora interrupta 133
—*ovalis* 242, 264
Senonian 9
Steganoporella buski 22, 233
Steginopora 54, 241
—*ornata* 241
—*reticulata* 240
Stichocados 54
—*ordinatus* 255

Terebratulina lata, Zone of 9
tertiary frontal wall 33
Thomas, H. D. 7
Tricephalopora 30, 33, 35
Trochiliopora Bed 10
Turonian 8

Ubaghsia 18, 19*, 33, 35, 36, 39, 52, 53, **238**, 275
—*arcifera* 241
Ubaghsia aurita 18, 241, **244**, 275 ; Pl. 20, fig. 3
—*coaxans* 18, 241, **252**, 253*, 275 ; Pl. 20, fig. 6, Pl. 21, fig. 6
—*crassa* 18, 241, **250**, 251*, 263, 275 ; Pl. 20, fig. 5, Pl. 21, fig. 7
—*demorgani* 18, 36, 241, 242, 255, **256**, 275 ; Pl. 22, figs. 5, 6
—**gasteri** 241, **245**, 246*, 250, 275 ; Pl. 20, figs. 4, 7
—**langi** 241, **247**, 248*, 275 ; Pl. 21, figs. 4, 5
—*meudonensis* 18, 36, 242, 255, **256**, 257, 275 ; Pl. 22, figs. 3, 4
—*ocellata* 18, 36, 241, **255**, 256, 257, 275 ; Pl. 22, figs. 1, 2
—*ornata* 18, 31, 33*, 242, **260**, 262*, 275 ; Pl. 23, figs. 1–4
—*perforata* 18, 241, 244, **254**, 275
—*ranunculus* 18, 42, 241, **242**, 243*, 263, 275 ; Pl. 20, fig. 2
—*reticulata* 18, 36, 240, 242, 255, **257**, 258*, 262, 263, 275 : Pl. 21, figs. 1–3, Pl. 22, figs. 7, 8

Uintacrinus westphalicus, Subzone of 10

young zooecia 45

zoarial budding 23
—growth habit 21
zooecial walls other than the frontal walls 24
—size 24
—shape 24

BRITISH MUSEUM
12 JUN 1962
NATURAL HISTORY